



ISSN 0971-6777 (Print)  
ISSN 2277-8934 (Online)

# JOURNAL OF CAMEL PRACTICE AND RESEARCH

[www.camelsandcamelids.com](http://www.camelsandcamelids.com) • [www.indianjournals.com](http://www.indianjournals.com)

Volume 31

August 2024

Number 2

## In This Issue

The camel in the United Kingdom:

-monuments, memorials and municipalities

Camelology

The trends of camel research in North America:

-A bibliometric approach

Camel milk and its safety for human consumption-  
an Overview

Spleen - morphology and function

Prostate gland- histology during ante and postnatal  
period of ontogenesis

Haematological profile

-during dry period, early pregnancy,  
lactogenesis and galactopoiesis

Gastrointestinal nematodes

- in Sonid Bactrian camels

Blood parameters

-during different physiological states

Radiographic evaluation of impacted dulaa

Soft palate diverticulectomy

Weight, growth and genetics in Shami camels



# JOURNAL OF CAMEL PRACTICE AND RESEARCH

## EDITOR

**TARUN KUMAR GAHLOT**

Camel Publishing House

Bikaner - 334001, INDIA

Email : [tkcamelvet@yahoo.com](mailto:tkcamelvet@yahoo.com)

Website : [www.camelsandcamelids.com](http://www.camelsandcamelids.com) • [www.tkgahlotcamelvet.com](http://www.tkgahlotcamelvet.com) • [www.indianjournals.com](http://www.indianjournals.com)

## Members of the Editorial Board

Adel I Alsheik-Mubarak	Saudi Arabia	Mehta SC	India
Amir Niasari-Naslaji	Iran	Moosavi-Movahedi AA	Iran
Ashraf M. Abu-Seida	Egypt	Muyldermans Serge	Belgium
Binoy, S. Vettical	UAE	Nagy P	U.A.E.
Eerdunmutu	China	Rollefson IK	Germany
Faye B	France	Saber AS	Egypt
Hasi Surong	China	Sabry Mohamed Bahy El-Bahr	Saudi Arabia
Jirimutu	China	Schuster RK	U.A.E.
Kataria AK	India	Tinson A	U.A.E.
Kataria N	India	Wael M. EL-Deeb	Saudi Arabia
Kinne J	U.A.E.	Wani Nisar Ahmad	U.A.E.
Mahmoud Kandeel	Saudi Arabia	Wasfi Ibrahim	U.A.E.
		Wernery U	U.A.E.

## Assistant Editors

Sakar Palecha

Mahendra Tanwar

Kapil Kachwaha

Kachwaha Subhash



**CAMEL PUBLISHING HOUSE**

Bikaner - 334001, INDIA

**Manuscripts and other related correspondence may be made to :**

Dr. Tarun Kumar Gahlot  
Editor, Journal of Camel Practice and Research  
67, Gandhi Nagar West  
Near Lalgarh Palace  
Bikaner-334001, INDIA

**Mobile** : 0091-9414137029

**Email** : tkcamelvet@yahoo.com

**Website** : www.camelsandcamelids.com • www.tkgahlotcamelvet.com • www.indianjournals.com

### **Scope of Journal of Camel Practice and Research**

Journal of Camel Practice and Research (JCPR) publishes only research and clinical manuscripts related to the Camelids (Old and New World camelids), hence published contents are consistent with the title and scope of the journal. Review articles on emerging research are invited and published. JCPR also publishes the news related to the New or Old World Camelids, specially those related to new products, conferences, books, trainings or workshops etc.

## **SUBSCRIPTION RATE - 2024**

ANNUAL

Rs. 4500/- or US \$ 450

*Note : Subscription in Rupees is applicable to Indian subscribers only.*

**Publisher** : The **Journal of Camel Practice and Research** (Triannual) is published by the “**Camel Publishing House**” 67, Gandhi Nagar West, Near Lalgarh Palace, Bikaner-334001, India. email: tkcamelvet@yahoo.com

**Cover Design:** Tarun Kumar Gahlot

**Cover Photo:** The camel group representing Africa at the northeast corner of the Albert Memorial. Source: [https://en.wikipedia.org/wiki/Albert\\_Memorial#/media/File:Albert\\_Memorial\\_statue.JPG](https://en.wikipedia.org/wiki/Albert_Memorial#/media/File:Albert_Memorial_statue.JPG)

**Courtesy:** R Trevor Wilson

**Printer:** Sankhla Printers, Vinayak Shikhar, Near Polytechnic College, Bikaner-334003, India.

**Phone:** 0091 - 151 - 2242023

CONTENTS		
Volume 31	August 2024	Number 2
S.No.	Title of Contents and Authors	Page No.
1.	The one-humped camel in the United Kingdom: monuments, memorials and municipalities R. TREVOR WILSON	105-117
2.	Camelology : definitions, history and scientific challenges Bernard Faye and Tarun Kumar Gahlot	119-130
3.	The trends of camel research in North America: A bibliometric approach Ahmed Alsaleem and Mahmoud Kandeel	131-145
4.	Some aspects of camel milk and its safety for human consumption-An overview Shehadeh Kaskous and Michael W. Pfaffl	147-161
5.	Morphology and function of the spleen in camel: A comparative review A.M. Marwa-Babiker and Nawal M. Elkhair	163-169
6.	Histological structure of the prostate gland during ante and postnatal period of ontogenesis in dromedary camel in Algeria Hafiza Aidoudi, Djallal Eddine Rahmoun and Tarek Khenenou	171-177
7.	Haematological profile variations in female dromedary camels during dry period, early pregnancy, lactogenesis and galactopoiesis in algerian dromedary camels Maria Chikha, Tarek Khenenou, Djalel Eddine Gherissi, Soundes Akriche, Mohammed Saci Ferdia, Rabah Mayouf and Sabry Mohamed El-Bahr	179-182
8.	Prevalence investigation of gastrointestinal nematodes in Sonid Bactrian camels Bin Hou, Mila Tai and Surong Hasi	183-188
9.	Determination and identification of reference intervals for selected blood parameters during different physiological states in female dromedary camels Nawal Mohamed Elkhair	189-200
10.	Radiographic evaluation and soft palate diverticulectomy of impacted dulaa in two camels ( <i>Camelus dromedarius</i> ) Ruchi Patwa, Mahendra Tanwar, P. Bishnoi, A.K. Bishnoi, Shivangi Diwedi and Balkrishan Yadav	201-204
11.	Weight, growth and genetics in Shami camels ( <i>Camelus dromedarius</i> ) Khaled A Al-Najjar, Moammar Dayoub, Azzam N. Al-Yacoub and Widya P.B. Putra	205-210
12.	News	118, 146, 162, 170, 178
13.	Instructions to Contributors	211-215



# EXPLORATION OF CAMELS IN ARCHIVES AND HERITAGE IN DIFFERENT COUNTRIES BY

## R. TREVOR WILSON

R. Trevor Wilson's maiden contribution to the world of camels was a minor paper published in 1978 in the journal *Tropical Animal Health and Production* as part of a series on the domestic livestock of Southern Darfur in Sudan where he worked and lived with his family in 1972-1974 and 1976-1978. His reference book "The Camel" was published in the year 1984. The book proved a plethora of scientific information about camels in a classified manner. The book received a big opening with great appreciations because at that time there was a big void of camel related books. In the International Year of Camelids 2024, it is important to recall the outstanding research done by R Trevor Wilson on camel archives and heritage in different countries through several monuments, pictures, statues, stamps, archives, old literature and other relevant evidences. He reviewed the presence and performance of the one-humped camel - exotic to Tanzania<sup>1</sup>. Accordingly, camels appeared in annual veterinary reports in 1926-1934: numbers varied from 26 (1926) to 67 (1930) and 5 (1934). In this pre-independence period they occurred mostly in the coastal provinces and since independence some 340 camels have been imported privately and by NGOs.

Later, Dr Wilson and his co-researchers explored camels in Turkey<sup>4</sup> and found that camels were first known to have been in Turkey some 2600 years ago when they were used as animals of war. There has been continued presence ever since and the country is unusual as it is home to both one-humped and bactrian types. During the nineteenth and as late as the early twentieth century there was an important exchange of breeding animals between Turkey and Syria. Numbers declined rapidly after the 1930s and at the end of the first decade of the twenty-first century the country's camel population was little more than 1000 animals. These are now mainly used for sport and as a tourist attraction with meat being a minor product.

He gave first comprehensive account of the presence of camels and of camel production in Uganda. The dromedary camel started to appear in the arid northeast of the country during the 1960s/1970s. In 2008 the national camel population was about 31 000 but in 2017 could go as high as or more than 40 000. Traditional pastoral tribes have become camel keepers for the production of milk, some meat, some transport and for medical uses. It was inferred that these camels contribute to food security and to the livelihoods of pastoralists in some of the remote parts of Uganda<sup>2</sup>.

His further research focused on Somaliland which is a self-proclaimed republic and had about 1.7 million camels and the camel has always been an icon in the culture and customs of the ethnic Somali. The British administration organised a Camel Corps to help in the maintenance of law and order in the then Protectorate but it also distinguished itself against the troops of Italian Somaliland in the World Wars of 1914-1918 and 1939-1945 before it was disbanded in 1946<sup>3</sup>.

His further research revolved around Eritrea and Ethiopia, among the poorest in the world<sup>5</sup>. Ethiopia has the greatest number of livestock in Africa at an estimated 120 million ruminant animals of which 1.1 million are camels. In contrast Eritrea's livestock population is under 10 million in which camels may number 320 000. The one-humped camel possibly arrived in the area of study about 1900 years ago. In Eritrea camels were owned by Beja tribes near the border with Sudan, by Tigre clans in the north and by Afar and some Somali in the east along the Red Sea littoral. Ownership in Ethiopia was mainly by the Somali people and by the Afar in their respective Regional States in eastern Ethiopia and by the Boran in the south.

His most recent publication unfolds the use of camels in the British military and in transport of trade goods for British trading companies from their areas of origin and have been recognised as

making considerable economic contributions to Britain in these roles. In recognition of this there are several monuments and memorials to the animal around the country. Camel is also a word used in British geography for some British towns and villages and as names for geographic features such as rivers and hills. This paper provides a repertoire of such monuments, memorials and place names<sup>6</sup>.

This editorial is specially dedicated to R. Trevor Wilson with a big salute because he chose the Journal of Camel Practice and Research as a platform for publication of his research on this aspect.



(Dr. Tarun Kumar Gahlot)  
Editor


### References:

1. Wilson, Richard. The one-humped camel and the environment in northern Tanzania. Journal of Camel Practice and Research. 2011; 18:25-29.
2. R. Trevor Wilson. The one-humped camel in Uganda. Journal of Camel Practice and Research. 2017; 24(1):1-7. DOI : 10.5958/2277-8934.2017.00001.7
3. R. Trevor Wilson. The one-humped camel in Somaliland. Journal of Camel Practice and Research. 2000; 27(1):1-12. DOI : 10.5958/2277-8934.2020.00001.
4. Yilmaz Orhan, Ertugrul Mehmet, Wilson R Trevor. The domestic livestock resources of Turkey: Camels. Journal of Camel Practice and Research. 2011; 18(1):21-24.
5. R. Trevor Wilson. The one-humped camel in Eritrea and Ethiopia: a critical review of the literature and a bibliography. Journal of Camel Practice and Research. 2020; 27(3):229-262. DOI : 10.5958/2277-8934.2020.00034.X
6. R. Trevor Wilson. The one-humped camel in the United Kingdom: monuments, memorials and municipalities. Journal of Camel Practice and Research. 2024; 31(2):105-117.

# SELECTED RESEARCH ON CAMELID IMMUNOLOGY

(Hard Bound, 392 pages, few figs coloured, Edition 2016)

In 1989 a group of biologists led by Raymond Hamers at the Free University Brussels investigated the immune system of dromedaries. This discovery was published in Nature in 1993. Based on their structure, these peculiar camelid antibodies have been named Heavy Chain Antibodies (HCAb), as they are composed of heavy chains only and are devoid of light chains. Sera of camelids contain both conventional heterotetrameric antibodies and unique functional heavy (H)-chain antibodies (HCAs). The smaller size and monomeric single domain nature make these antibodies easier to transform into bacterial cells for bulk production, making them ideal for research purposes. Camelid scientists world over were greatly fascinated by a new field of research called "Camelid Immunology". Significant research has been done on camelid immunology in recent decade. In order to benefit future camelid immunology researchers, this book was planned in the series of "Selected Topics" by Camel Publishing House with a title- "Selected Research on Camelid Immunology" edited by T.K. Gahlot, U. Wernery and Serge Muyldermans. This book is a unique compilation of research papers based on "Camelid Immunology" and published in Journal of Camel Practice and Research between 1994-2015. Research on this subject was done in 93 laboratories or institutions of 30 countries involving about 248 scientists. In terms of number of published papers in JCPR on the immunology the following countries remain in order of merit (in parenthesis), i.e. Iran (1), India and UAE (2), China and Saudi Arabia (3), Sudan (4), Kenya and Belgium (5), USA (6), Germany (7) and so on. The book contains 11 sections and is spread in 384 pages. The diverse sections are named as overview of camel immune system; determinates of innate immunity, cells, organs and tissues of immune system; antibodies; immunomodulation; histocompatibility; seroprevalence, diagnosis and immunity against bacteria, viruses, parasites and combination of other infections; application of camel immunoglobulins and applications of immune mechanisms in physiological processes. The camelid immunology has to go a long way in its future research, therefore, this reference book may prove quite useful for those interested in this subject. Book can be seen on [www.camelsandcamelids.com](http://www.camelsandcamelids.com).



Editor:

T.K. Gahlot

U. Wernery

Serge Muyldermans

Edition: 2016

© Camel Publishing House



Publisher:

**Camel Publishing House**

67, Gandhi Nagar West, Near Lalgargh Palace,  
Bikaner-334001 Rajasthan, India

email: [tkcamelvet@yahoo.com](mailto:tkcamelvet@yahoo.com)

Website:

[www.camelsandcamelids.com](http://www.camelsandcamelids.com)

[www.tkgahlotcamelvet.com](http://www.tkgahlotcamelvet.com)

Price: US \$ 475

INR 12500

ISBN 81-903140-4-1



# SELECTED RESEARCH ON CAMELID PARASITOLOGY

Hard bound, 291 pages, few figures coloured

New research and experience always broaden our knowledge, and help us adopting new diagnostic methods and treatments. Camel Publishing House has taken a step forward to compile this knowledge in form of a book and this Herculean task was accomplished with the help of dedicated editors. viz. Drs. T.K. Gahlot and M.B. Chhabra. *Selected Research on Camelid Parasitology* is most comprehensive guide to Camelid Parasitology. The classic reference book serves as a one stop resource for scientific information on major aspects of Camelid Parasitology. Featuring abundant photographs, illustrations, and data, the text covers camelid protozoa, helminths, and arthropods of dromedary and New World camelids. This hard bound book of 304 pages contains seroepidemiological studies, immunological and other diagnostic procedures, and new treatments of parasitic diseases. There are at least 17 countries involved in camelid parasitology research, viz. Ethiopia, France, India, Iran, Jordan, Kenya, Libya, Mauritania, Nigeria, Sultanate of Oman, Pakistan, Saudi Arabia, Sudan, Sweden, United Arab Emirates, Uganda and U.S.A. As per published papers in Journal of Camel Practice and Research (JCPR), 173 authors have contributed 72 manuscripts which are appropriately placed in 5 sections. The text of each manuscript published previously in JCPR remains the same except the pattern of numbering the references in the body of text. This book indicates a swing of camelid research during period 1994-2008 and will help identifying the missing links of research in this subject.

Editors:

T.K. Gahlot and M.B. Chhabra

Edition: 2009

© Camel Publishing House

Publisher: **Camel Publishing House**

67, Gandhi Nagar West,  
Near Lalgarh Palace  
Bikaner 334001 Rajasthan,  
India

email: [tkcamelvet@yahoo.com](mailto:tkcamelvet@yahoo.com)

website: [www.camelsandcamelids.com](http://www.camelsandcamelids.com)

Price: US\$ 200 (Abroad)

INR 3000 (India)

ISBN: 81-903140-0-9

## SELECTED RESEARCH ON CAMELID PARASITOLOGY

Editors

T.K. Gahlot  
M.B. Chhabra



# SUBSCRIPTION - 2024

FOR

## **JOURNAL OF CAMEL PRACTICE AND RESEARCH**

*(Triannual In English Language, April, August and December Issue Every Year)*

### SUBSCRIPTION RATE - 2024

ANNUAL

Rs. 4500/- or US \$ 450

*Note : Subscription in Rupees is applicable to Indian subscribers only.*

### Subscription Form

I want to become annual subscriber of the **Journal of Camel Practice and Research**, for/from the year 2024 ..... For this purpose I am enclosing herewith a cheque / demand draft number ..... dated ..... for Rs./US \$. ..... in favour of "**Camel Publishing House**". The cheque or D.D. should be payable at State Bank of India, Code No. 7260, Bikaner. Payment may be made through payment portal of website [www.camelsandcamelids.com](http://www.camelsandcamelids.com) or money transfer to bank account.

**Name** : .....

**Permanent Address** : .....

: .....

**Country** : .....

**Signature** : .....

Mail to :

**Camel Publishing House**

67, Gandhi Nagar West

Near Lalgah Palace

Bikaner - 334001, INDIA

email : [tkcamelvet@yahoo.com](mailto:tkcamelvet@yahoo.com)

website : [www.camelsandcamelids.com](http://www.camelsandcamelids.com)



# THE ONE-HUMPED CAMEL IN THE UNITED KINGDOM: MONUMENTS, MEMORIALS AND MUNICIPALITIES

R. TREVOR WILSON

Bartridge Partners, Bartridge House, Umberleigh, EX37 9AS, UK

## ABSTRACT

The one humped camel (*Camelus dromedarius*) has never been numerous in the United Kingdom and in British law is classed as a dangerous wild animal for which a licence is needed by the keeper. It is now illegal to use camels for entertainment, for example in circuses. There have been, nonetheless, both one humped and two humped (*Camelus bactrianus*) camels in Britain since the early 12<sup>th</sup> century, 900 years ago. Camels have seen service in the British military and in transport of trade goods for British trading companies from their areas of origin and have been recognised as making considerable economic contributions to Britain in these roles. In recognition of this there are several monuments and memorials to the animal around the country. Camel is also a word used in British geography for some British towns and villages and as names for geographic features such as rivers and hills. This paper provides a repertoire of such monuments, memorials and place names.

**Key words:** Albert Memorial, Bactrian camel, Cleopatra's Needle, General Charles Gordon, Imperial Camel Corps, Kitchener of Khartoum

Most of the world's countries class the camel as a domesticated animal but the UK is among a small minority that considers camels as wild. Camels – unspecified as to being Old World or New World – are classified in British Law as 'Dangerous Wild Animals' for which a licence is required in order for the general public to keep them (LGU, 1976). There are separate regulations for zoos, circuses and some other bodies but live performances in circuses have been banned since January 2020 (LGU, 2019). Up until 2019 two circuses in the UK held wild animal licences but the Wild Animals in Circuses Act 2019 banned their use in the entertainment industry from January 20, 2020. Several British zoos and wildlife parks continue to exhibit Bactrian camels but only two 'safari' parks keep the one humped camel. It seems there is only one commercial herd of one humped camels in the UK. The herd, with animals obtained from Europe and maintained under a Dangerous wild animal licence, comprised 17 animals aged from 2 to 25 years old in May 2024. The unit was founded as a pseudo-safari business in the 2<sup>nd</sup> decade of the 21<sup>st</sup> century but converted to a dairy herd, milked by a mobile machine, as a result of the Covid pandemic in 2020 when movements were not allowed. Milk was not being produced in May 2024 because

imports of pregnant females were stopped, as with all ruminants, due to the risk of contracting BTV-3 (bluetongue disease) (Joseph and Rebecca Fossett, personal communication 13 May, 2024).

There has never been more than a very small number of camels in the United Kingdom. There is, however, solid evidence for the presence of both one- and two-humped camels in England, Scotland and Ireland at least as far back as the early 12<sup>th</sup> century (900 years ago). Henry I of England (king from 1100 to 1135 AD) is known to have had camels in his menagerie at Woodstock near Oxford (Mynors *et al*, 1998) with potential hints of an even earlier presence. Later, in 1466 in the mid 15<sup>th</sup> century, the Patriarch of Antioch (a city in Greece) presented four "dromedaries" and two "camels" (i.e. Bactrians) to the English King Edward IV (Searle, 1902). The tomb of a John Camel in the Church of St John the Baptist in Glastonbury in Somerset, southwest England dated 1487 has a carving of a Bactrian camel: the history of this camel does not appear to be known but it may be a pun on the occupier's name.

Several geographical place names contain the word camel in addition to Camelford in Cornwall (see below). There is a West Camel village on the River Cam in Somerset and another nearby village

---

SEND REPRINT REQUEST TO R. TREVOR WILSON [email: trevorbart@aol.com](mailto:trevorbart@aol.com)

known as Queen Camel: none of these places has any connection with the camel as an animal but have been derived from earlier indigenous place names. The Sopwith Camel was a single engine biplane fighter aircraft of the First World War whose name was said to derive from the hump at the front of the aircraft that housed the machine guns.

In spite of there only ever having been very few camels in the United Kingdom their various contributions to military operations and to international trade the country has several “hard” memorials to their role in these contexts. This paper provided a partial repertoire of such mementos.

## Methodology

This paper is based on the author’s own knowledge and experience and on a thorough review of the literature. In view of the rather unusual subject formal sources are relatively few. Much of the text is thus based on grey literature and the ever-growing number of internet sources usually under the epithet of “blog”

## Monuments, Memorials and Municipalities

### *Prince Consort National Memorial (The Albert Memorial)*

The Albert Memorial, north of the Royal Albert Hall in Kensington Gardens, London, was commissioned by Queen Victoria in memory of her husband Prince Albert, who died in 1861. Designed by Sir George Gilbert Scott in the Gothic Revival style it is an ornate canopy or pavilion 176 feet (54 m) tall, like a ciborium over a church high altar with a statue of the prince facing south below (Risebero, 1985) (Fig 1). It took more than ten years to complete, the £120,000 cost (the equivalent of about £19,000,000 in 2024) met by public subscription. The memorial was opened in July 1872 by Queen Victoria although Albert’s statue was not ceremonially “seated” until 1876. It has been Grade I listed since 1970 (Appendix A).<sup>1</sup>

Each of the four corners of the central area has an allegorical sculpture depicting Victorian industrial arts and sciences, these being agriculture, commerce, engineering and manufacturing. At the outer corners are four more groups representing the traditional four

continents: Africa, the Americas, Asia and Europe. Each of these groups comprises several ethnographic figures and a large animal: a bison is represented for the Americas, an elephant for Asia, a bull for Europe and a one-humped camel for Africa (Fig 2).

The memorial was planned by a committee of well-known architects that was led by Sir George Gilbert Scott. The sculptural work was coordinated by Henry Hugh Armstead who recruited many of the artists from the Royal Academy. William Theed, who was a favourite of Prince Albert, was chosen to carve the “Africa” group.

### *Imperial Camel Corps Memorial, Victoria Embankment Gardens, London WC2*

The beginning of what eventually became the Imperial Camel Corps was four Company-strength units raised in Egypt in January 1916. The companies comprised, respectively, Australian troops that had returned from the disastrous Gallipoli Campaign, New Zealand, British and Indian troops. A Company of 130 men comprised a small Headquarters and four Sections, each of 7 groups of 4 men armed with Lee-Enfields, the then standard British Army bolt action rifle. The first camels were heavy baggagers from the Bikaner Camel Corps but these were later replaced by lighter Egyptian camels for riding mounts (Falls and MacMunn, 1930; Clayton and Killingray, 1989). Six new companies were raised from British yeomanry regiments in March 1916, four additional Australian companies were raised in June from reinforcements intended for the Australian Light Horse regiments and reinforcements intended for the New Zealand Mounted Rifles Brigade formed two companies, one in August and the other in November (NZMCH, n.d.).

Initial activities were confined to long range patrols around the Suez Canal and the Sinai Desert. In August 1916, a transition to a more active combat role took place when each Company added a machine gun Section of 15 men with three Lewis guns. Additional staff on the Headquarters section brought each Company’s strength to 184 men. Companies operated independently as cavalry whilst on the march but dismounted to fight as infantry in actual battle and thereafter, the Corps (as it eventually became) continued to fight as mounted infantry.

The unit eventually expanded to Battalion size. Four such entities were eventually formed, one each from Great Britain (also known as the Imperial) and New Zealand (ANZAC) and two from Australia, each of 770 men and 922 camels. When Brigade status was achieved on 19 December 1916 support troops

1. Buildings, scheduled monuments, parks and gardens, battlefields and shipwrecks judged of national importance in the United Kingdom in terms of architectural or historic interest are “listed” in one of three grades. These are: Grade I (buildings of exceptional interest); Grade II\* (particularly important buildings of more than special interest); and, Grade II (buildings of special interest).

included a mountain artillery battery (Hong Kong and Singapore (Mountain) Battery), machine gun squadron (265<sup>th</sup> (Camel) Machine Gun Squadron), Royal Engineers (10<sup>th</sup> (Camel) Field Troop), field ambulance (Australian (Camel) Field Ambulance) and 97<sup>th</sup> Australian Dental Unit, this last comprising only four men!, the Mobile Veterinary Section and – inevitably – an administrative train including a Brigade Ammunition Column and the Brigade Train which carried five days of supplies (Underwood, 2003). Brigade strength at this stage was about 4150 men and 4800 camels (NZMCH, n.d.).

The Imperial Camel Corps (ICC) eventually became part of the Egyptian Expeditionary Force (EEF).<sup>2</sup> It fought in many battles in the Senussi Campaign (November 1915-February 1917), the Sinai and Palestine Campaign (January 1915-October 1918) (Reid, 1934; Robertson, 1938) and in the Arab Revolt (June 1916-October 1918). The ICC was disbanded in May 1919 after the end of the war having had more than 240 men killed over the period of its existence.

A memorial to the Imperial Camel Corps was unveiled in the Victoria Gardens on the Thames Embankment in London on 22 July 1921 (Ward-Jackson, 2012) and listed as Grade II in 1958 (Appendix B). The bronze effigy of a rider carrying a rifle was sculpted by Major Cecil Brown (who had himself served in the Corps) in 1919-1920 and unveiled in July 1921 (Fig 3). The monument is mounted on a two-stage pedestal of Portland stone with the upper part bearing bronze bas-relief images of various actions. The lower part has bronze plaques, the one with a sentiment commemorating those men who made the supreme sacrifice:

To the Glorious and Immortal Memory of the Officers, N.C.O.s and Men of the Imperial Camel Corps -- British, Australian, New Zealand, Indian -- who fell in action or died of wounds and disease in Egypt, Sinai, and Palestine, 1916, 1917, 1918. (Fig 4).

The other sides carry the names of all the members of the Corps who were killed or died of their wounds during the war and all the battles and engagements fought by the Corps;

1916: Romani, Baharia, Mazar, Dakhla, Maghara, El Arish, Maghdaba.

1917: Rafa, Hassana, Gaza 1, Gaza 2, Sana Redoubt, Beersheba, Bir Khu Weilfe, Hill 265.

1918: Amman, Jordan Valley, Mudawar (Hedjaz).

2. It is not clear how the camel fighting force became a Corps. In the British Army a Corps (commanded by a Lieutenant General) comprises at least two Divisions (each commanded by a Major General) made up of several Brigades (each commanded by a Brigadier)

## ***Statue of General Gordon at the Royal School of Military Engineering, Brompton Barracks***

Charles George Gordon was born into a military family in Woolwich on 28 January 1833. He obtained a commission as Second Lieutenant in the Royal Engineers in 1852. After fighting in the Crimean War in 1853-1856 he was then in China in 1860 with the Anglo-French forces. At 30 years of age in 1863 Gordon commanded a force established in Shanghai to protect Europeans from Chinese rebels and within two years the revolt known as the Taiping Rebellion was put down. As a result, the Chinese Emperor honoured him with the yellow jacket and peacock feather of a mandarin after which he became known by the soubriquet of “Chinese” Gordon although, he was promoted only to Captain in the British Army (Barnes, 1885; Trench, 1979).

Promoted Colonel in 1872 Gordon accepted an offer from the Egyptian Khedive in 1874 – with British Government consent – to join the Egyptian army. He was shortly appointed Governor of Equatoria Province (now mainly South Sudan and northern Uganda), where he attempted to suppress the slave trade and where he stayed until 1876. Gordon then returned to England but was eventually persuaded by the Khedive to accept the Governor-Generalship of the entire Sudan which also carried the Ottoman title of Pasha (Sparrow, 1962).

Gordon tried hard to suppress the slave trade but was hindered in this by not only the corruption of his Egyptian official but also of his British colleagues (Moore-Harell, 1998). He resigned as Governor General in 1879. Offered positions in the Belgian Congo and in the Cape Colony, Gordon refused but then accepted a post as Private Secretary to the Governor-General of India before returning to China (Galbraith, 1971; Hsu, 1964). He spent much of 1881 in Mauritius and the Seychelles before moving to Palestine. Gordon was promoted to Major General in the British Army in March 1882 (Boulger, 1897).

In January 1884, Gordon was again sent to the Sudan by the British Government. At this time, the country was in turmoil as a result of the carnage being created by a religious fanatic known as the Mahdi and his followers who were winning battles against the Egyptian army over most of the country.<sup>3</sup>

3. Mahdi is an Arabic word that means “guided to the right path” or “rightly guided”. In Islam, Mahdi is the prophesied messianic figure who will appear at the end of the world to rid humanity of evil and injustice. Mahdi is also a popular and deeply spiritual choice given by Muslim parents to their children (<https://www.thebump.com/b/mahdi-baby-name>).



Gordon's brief was to evacuate the Egyptian garrisons endangered by the revolt of the Sudanese rebels and then to leave the country himself. He successfully evacuated several expatriate families but refused to leave himself and sought to hold the central areas but was besieged in Khartoum. The city held out for 10 months before it fell on 26<sup>th</sup> January 1885 as the rebels stormed the Governor's Palace. The whole garrison was slaughtered, Gordon being killed, supposedly through a spear thrust from one of the attackers. A relief expedition sailing up the Nile, for which the British public had been demanding for a considerable time but over which the Government had prevaricated, reached Khartoum two days later. It was at this point that Gordon was hailed as a martyr and a national hero. There were calls for revenge but this did not come until 13 years later when another British General, Herbert Kitchener, recaptured Khartoum and reinstalled a kind of peace (Ziegler, 1973).

There was a general feeling that a permanent memorial to Gordon in the form of a statue should be created but it was Gordon's own Corps, the Corps of Royal Engineers that took the initiative. Officers and other ranks contributed money to a fund to commission a statue of Gordon. The original specification was for a statue on foot in the uniform of an Egyptian General. The selected sculptor, Edward Onslow Ford, considered it more appropriate for Gordon to be seated on a camel.<sup>4</sup> Ford studied the camels at London Zoo and made casts of a dead camel (said to be a one humped beast). The resulting monument, of bronze and stone is of a one humped camel that is correct in most details. The exception is that the beast has a horse's tail rather than that of a camel intended as a sop to late Victorian (and perhaps especially Queen Victoria's) morality (Droth *et al*, 2014; Darke, 1991). The statue was first exhibited and unveiled by the Prince of Wales at the Royal Academy in 1890. It was later moved to Brompton Barracks, the home of the Royal School of Military Engineering in Chatham, where it still stands (Fig 5). It became Grade II\* listed in 1998 (Appendix C).

### ***Statue of General Gordon at Gordon's School***

A considerable grounds well persisted in favour of a statue to Gordon being put on public display at the site of his death in Khartoum. The hero of the

recapture of Khartoum, General Kitchener together with Algernon Borthwick, the owner of the Morning Post newspaper, persuaded the paper's readers to subscribe to a fund for this to happen. By 1902, there was sufficient money for a second statue to be cast. The new effigy was erected in St Martin's Place in London (Fig 6). In 1904, it was shipped for Khartoum although it survived two major mishaps, the first being that of the Cedarden which sank in the Thames and the second being the SS Lesbian which was submerged in the Nile! Finally it arrived in Khartoum where it was placed at the intersection of Gordon and Victoria Avenues, some 200 metres south of the new palace that had been built in 1899 after the former edifice had been destroyed (Fig 7).

The Anglo-Egyptian Sudan, a Condominium ruled jointly by the United Kingdom and Egypt, gained its independence on 1 January 1956 to become the Republic of Sudan. In 1958, the Sudanese government removed the Gordon statue and a second one of Kitchener which had been erected later. Both statues were offered to the UK, the offer being accepted and the statues were shipped back to England.

The UK government then offered the Gordon statue to Gordon's School, founded as the Gordon Boys' Home in 1885, in Woking in the county of Surrey. The statue was erected at the School in April 1959 and unveiled in May 1959 (Fig 8). Time has wreaked its worst on the statue and eventually the Portland stone plinth and the bronze statue were degraded and tassels and reins were missing. The statue was eventually given a make over and HRH The Earl of Wessex unveiled a plaque to mark the restoration of the statue on 10 October 2014. The front of the plinth now bears the inscription "Charles George Gordon". On the left side the inscription reads "Born Woolwich 28<sup>th</sup> January 1833, killed Khartoum 26<sup>th</sup> January, 1855"; the right side is inscribed with "This Statue was Erected at Khartoum in 1904. Removed and presented to the School in 1959" with the remaining side bearing the words "This plinth has been presented in memory of Major-General Sir Hubert J. Huddleston, G.G.M.C., G.G.B., C.B, D.S.O., M.C., who served the Sudan for 28 years and was Governor-General 1940-47." The statue was listed as Grade II on 29 April 2015 (Appendix D).

### ***Camel benches on Victoria Embankment of River Thames, London WC2***

Following British victories at the Battle of the Nile in 1798 and the Battle of Alexandria in 1801 the

4. Edward Onslow Ford RA (27 July 1852 - 23 December 1901) was an English sculptor whose early success was with portrait heads or busts of prominent people. Several of Ford's monumental commissions celebrate the British Empire either by promoting imperial values or as memorials to military figures.



**Fig 1.** Full view of the Albert Memorial in front of the Royal Albert Hall, London. Source: Photo by the Author.



**Fig 2.** The camel group representing Africa at the northeast corner of the Albert Memorial. Source: [https://en.wikipedia.org/wiki/Albert\\_Memorial#/media/File:Albert\\_Memorial\\_statue.JPG](https://en.wikipedia.org/wiki/Albert_Memorial#/media/File:Albert_Memorial_statue.JPG).

Khedive of Egypt, Muhammad Ali, sought to reward the victors with some unwanted pieces of Egyptian antiquity. One was an obelisk which became known as Cleopatra's Needle presented to the United Kingdom in 1819. The British Government, however, were unable to justify the cost of its transport to England at that time. It therefore remained in




**Fig 3.** Memorial to Imperial Camel Corps in Victoria Gardens on Thames Embankment. Source: [https://en.wikipedia.org/wiki/File:Statues\\_in\\_Victoria\\_Embankment\\_Gardens\\_-\\_geograph.org.uk\\_-\\_1729996.jpg](https://en.wikipedia.org/wiki/File:Statues_in_Victoria_Embankment_Gardens_-_geograph.org.uk_-_1729996.jpg).

Alexandria for more than 50 years until anatomist and dermatologist Sir William James Erasmus Wilson provided the money for the obelisk – of red Aswan granite, 68 feet (21 metres) tall and weighing more than 220 tons – to be dug out of the sand and transported to London. Encased in an iron cylinder and towed by a ship the obelisk left Alexandria on 21 September 1877 and after a perilous journey through stormy waters and via a harbour in northwest Spain it arrived at Gravesend in southeast England on 21 January 1878. Transported to London, Cleopatra's Needle was erected on the Victoria Embankment of the River Thames on 12 September 1878 (Wilson, 1877; Dickens, 1885).<sup>5</sup>

5. Two other "Cleopatra's Needles" are extant: one in New York which is the precise image of the London one in 1881 and one in Paris which is of a different provenance to the London and New York obelisks.





**Fig 4.** Detail of inscription on ICC statue. Source: enlarged from Fig 3.



**Fig 5.** Gordon's statue at the Brompton Barracks, School of Military Engineering in Chatham. Source: [https://en.wikipedia.org/wiki/Statue\\_of\\_General\\_Gordon#/media/File:Gordon\\_Memorial,\\_New\\_Brompton,\\_England-LCCN2002708002.jpg](https://en.wikipedia.org/wiki/Statue_of_General_Gordon#/media/File:Gordon_Memorial,_New_Brompton,_England-LCCN2002708002.jpg).

During the 1860s, some 22 acres (8.9 ha) of marshland along the River Thames were drained and reclaimed, a retaining wall was built and a new sewage system and other infrastructural improvements were constructed. The new – now the Victoria Embankment – terrace thus created needed,



**Fig 6.** The second cast of Gordon on a camel in St Martin's Place, London before shipment to Khartoum in 1902. Source: <https://www.londonremembers.com/memorials/general-gordon-on-a-camel>.



**Fig 7.** General Gordon's statue *in situ* in Khartoum 1936. Source: Library of Congress Prints and Photographs Division Washington, D.C. 20540 USA. <https://hdl.loc.gov/loc.pnp/pp.print>.

to the minds of London's councillors, appropriate street furniture so that people could rest after their perambulations.

George John Vulliamy, architect to the Metropolitan Board of Works, designed an ornamental "dolphin" lamp (the ornament is actually a sturgeon), along with cast iron and slatted wood benches that featured sphinx and camel-shaped armrests that expanded on the perceived historic, patriotic and imperial concepts of the obelisk (Fig 9; Fig 10). The benches were cast and constructed by Z.D. Berry & Son of Regent Street (a well-known and respected manufacturer of cast-iron street furniture and heating and lighting equipment) and installed on the embankment partly in the City of Westminster and partly in the City of London in 1877, one year before the Needle was erected in 1878.





Fig 8. Statue of General Gordon at Gordon's School, Woking, Surrey. Source: [geograph.org.uk - 44414.jpg](https://www.geograph.org.uk/photo/44414).



Fig 9. A camel bench on Victoria Embankment. Source: © Copyright N Chadwick and licensed for reuse under [creativecommons.org/licenses/by-sa/2.0/](https://creativecommons.org/licenses/by-sa/2.0/); <https://www.geograph.org.uk/photo/2714246>.



Fig 10. Detail of bench end from Fig 9 (Note: stamp of manufacturer at base).



Fig 11. Location view of Peek House at 20 Eastcheap, London. Source: Copyright Mike Quinn and licensed for reuse under Creative Commons Licence.

The Westminster ones feature sphinxes whereas the London ones feature camels. Some 21 benches were listed Grade II in 1987 of which only one featured a camel (Appendix E). By the time, the benches were listed, many had been replaced due to wear and tear and replaced by reproductions in 1960-1985. The originals have become “collectors’ items”: one was sold at auction in 2015 by the renowned auctioneers Bonhams at an asking price of £8000-£10,000 (even one of the later reproductions sold in 2019 for £3750) with the description:

**Lot 284: A Victorian cast iron black painted ‘Camel’ bench by Z.D. Berry & Son, designed by George John Vulliamy, circa 1878**

The later slatted scrolling back and seat with ends cast as recumbent laden camels with pierced scrolling leaves





**Fig 12.** Detail of the camel train above the main entrance to Peek House showing latest office tenant. Source: Photograph courtesy of Andrew G Wilson, 7 December 2022.



**Fig 13.** A spandrel with two supporting camels on the former Oriental Bank Corporation building at London's EC2 area. Source: <https://lookup.london/eastcheap-camels-peek-house/>

and flowerheads, on plinth bases, 213 cm wide, 79 cm deep, 83 cm high (83 ½ in wide, 31 in deep, 32 ½ in high).

### ***Camel train relief sculpture in Eastcheap, London EC3 and other London wall sculptures***

A relief of a camel train is sculpted above the entrance to a building that was once a bank after it had been the offices of a tea and coffee importing business at the corner of Eastcheap with Lovat Lane, London EC3.



**Fig 14.** Camels supporting the Coat of Arms of the Merchant Tailors on the wall of this Livery Company's home since 1347 almost opposite the previous Figure. Source: <https://lookup.london/eastcheap-camels-peek-house/>



**Fig 15.** The camel sculpture on the front of the Liberty store in Regent Street, London. Source: <http://www.speel.me.uk/gp/camels.htm>.




**Fig 16.** The Asia Tympanum on the facade of the Manchester Free Trade Hall. Source: (<http://www.speel.me.uk/sculptplaces/manchfreetradehall.htm>).





**Fig 17.** The Obelisk memorial to Scottish Merchant seamen on the Shore, Leith, Scotland. Source: Photograph by the Author.



**Fig 18.** Detail of the bronze casting of camels on the Scottish Merchant seamen memorial. Source: Photograph by the Author.



**Fig 19.** Windvane in the shape of a one humped camel on the old Town Hall, Camelford. Source: Photograph by the Author.



**Fig 20.** The Coat of Arms of the Borough of Camelford showing a two humped camel crossing a stream. Source: Photograph by the Author.

Peek Bros, established in 1823 by three siblings, was a firm of Tea and Coffee Importers. It was a very successful business and imported 5 million pounds weight (2232 tonnes, 2,267,960 kg) of tea to England in 1865. Their office building, known as Peek House, together with much of the street of Eastcheap was demolished in 1882 so that the ground could be excavated to allow the Metropolitan Underground Line to be built. An impressive new 4-storey building was erected at No 20 Eastcheap during 1883-1885 and again named Peek House. The two fronts in Eastcheap and Lovat Lane are joined by a quadrant turning through 90 degrees and it is above the main entrance on this quadrant that the carving is sculpted. The building (Fig 11) is now home to a number of offices including a major branch of the Hongkong and Shanghai Banking Corporation (HSBC).

The carving in high relief (alto-relievo) on Portland stone was created by William Theed the Younger. Theed loved exotic subjects and had already carved the Africa section on the Albert Memorial (see above). The 3 camels and the Arab leading them – effectively the Peek logo – are estimated to be about one quarter of life size and are carrying, respectively, tea, coffee and spices which were the main commodities traded by Peek Bros (Fig 12).

A spandrel on the wall of a building that was the office of the Oriental Bank Corporation between 1845 and 1892 situated where Adam's Court meets Threadneedle Street in London has two camels sculpted alongside two seated figures on each side of the central boss (Fig 13). Geographically, almost opposite the previous sculpture is one that is the Coat of Arms supported by 2 camels of the Merchant Tailors' Company (Fig 14). The Company is one of the twelve great Livery Companies established in Medieval times. It was originally the regulator and trade body of tailoring but over time it has become a philanthropic organisation (<https://www.merchant-tailors.co.uk/>). Liberty is a fashionable store in Regent Street, London W1 which describes itself as a "Designer Department Store Selling Luxury Brands". Liberty was established by the homonymous Arthur Lasenby Liberty in 1875 with the dream of setting up an emporium laden with luxuries and fabrics from distant lands (<https://www.libertylondon.com/uk/information/our-heritage.html>). In keeping with this dream is the camel sculpture on the long frontage of Liberty's in Regent Street, probably indicating trade and one of the exotic sources of Liberty's wares. The camel is seated with two Arabic traders and one naked worker who is pushing the camel possibly urging it to stand up (Fig 15).

### *Manchester Free Trade Hall Tympanum*

In the 19<sup>th</sup> century the city of Manchester and its surrounds were a major industrial area on a world scale. The Manchester Free Trade Hall dates from the Victorian era (1853-1856) and is an important decorated building in a Renaissance style that is 9 bays wide and 2 storeys high. Its glory is the tympani (tympanum -- a half-round recessed space above a window with an arch on top) of decorative allegorical sculpture on the upper level. The sculptor was John Thomas, an esteemed architectural sculptor, whose work was always of consistently high quality, with the Free Trade Hall sculptures showing him at his best. These sculptural groups in the tympani comprise a central classically-draped seated female figure has around her accoutrements appropriate to the idea she represents.

The central group represents Free Trade with 3 others relating to commerce, industry and manufactures, and the arts (<http://www.speel.me.uk/sculptplaces/manchfreetradehall.htm>). The remaining 5 represent the continents with which Manchester trades. Asia is the one of interest to this paper. The female figure is modestly garbed and wears a turban. There is a small treasure chest in her lap, a large cornucopia is held in one arm and other sculptures include the prow of a bird-headed trade ship, a tea chest with exotic fruit on top, an ornamented pot and rolls of fabric. The lady is seated on a camel (Fig 16).

### *Memorial to Scottish Merchant Navy at Leith, Scotland*

On 16 November, 2010 a memorial was unveiled by Her Royal Highness Princess Anne, daughter of her Majesty the Queen Elizabeth II, to sailors of the Merchant Navy who had been killed in or died as result of conflict from the start of World War I in 1914 to the unveiling of the memorial in 2010 (<https://www.iwm.org.uk/memorials/item/memorial/87515>).

The Memorial, located on The Shore in Leith near Edinburgh, is an 18 feet (5.5 m) tall sandstone obelisk featuring seafaring scenes (Fig 17). Each of the four sides has four tiers faced in light-coloured, grained Beryl sandstone and each panel contains a bronze relief. The architect and sculptor was the well-known Jill Watson: the nearby Powderhall Foundry cast the bronzes.

The upper panels represent the variety of merchant ships throughout history, the middle panels



have war scenes including destruction by torpedo, convoys with protection by Coastal Command and danger from rocks and storms at sea, the eye level panels are images of the world's trading routes, the ground level panels commemorating the educational role of Leith Nautical College and its training ship "Dolphin" and an empty lifebuoy evokes the loss of life at sea. One of the World Trade panels includes the Suez Canal and it is here that are shown two camels indicating that the Merchant Navy travelled world wide (Fig 18).

### ***The town and parish of Camelford, North Cornwall, West of England***

Camelford, a town in northeast Cornwall in southwest England, was first given a charter as a borough more than 800 years ago. The town has no real connection to a camel. The Cornish language name for the town, *reskammel*, derives from a combination of the Middle Cornish "rys" (ford) and the local river's Cornish name "kammel" (Ekwall, 1940; Mills, 2003). The English name, Camelford, was formed by Anglicisation of the river's name to Camel plus Ford, giving it an identical meaning to its Cornish counterpart. Due to the river's name sounding similar to the English word "camel" the animal is seen as a symbol of the town.

There are two obvious representations of a camel within the town limits. The most obvious is the model of a one humped camel acting as a weather vane atop the building in the town centre, erected in 1806, that was formerly the Town Hall and is now the Market Hall on the Ground Floor and the Public Library on the First Floor (Fig 19). The building was listed as Grade II in 1952 and then again listed Grade II in an amendment to the original heritage description in 1988 (Appendix F). As an integral part of the listed building the weather vane is protected under the Listing. The second representation comprises a stained glass window on the first floor of the building (and thus also protected) which shows a two humped camel crossing a body of water (Fig 20). The seal of the borough of Camelford is described as "Arg. a camel passing through a ford of water all proper" with the legend "Sigillum Vill de Camillford" (Pascoe, 1979). The logo of the town's Sir James Smith High School also is dominated by a silhouette of a one humped camel.

### **Discussion**

As a generally perceived and legally classed exotic or wild animal little is known by the British people of the contributions the one humped camel

has made to society over an extended time frame. Most of these contributions derive from outside the territorial UK where there has never been more than a few resident animals. This small number is likely to decrease in view of the legislation now in force. For more than two hundred years, however, camels have supported Britain and its economic success in a number of ways. Few memorials celebrate these contributions. In the area of military conflict it has supported conventional forces as a cavalry animal and kept troops in the field through its activities in transport. Its contribution to overseas trade is little understood other than by a minority of people. The few examples of such activities provided in this paper are, however, adequate testament to the position it has occupied and the esteem in which it is still held by a discerning establishment.

### **Acknowledgements**

Thanks to Andrew Wilson for the photo of the camel train on the wall in Eastcheap. Jill Watson, the sculptor of the Scottish Merchant Navy memorial very kindly provided me with information on her work on the obelisk. Joseph and Rebecca Fossett of Joseph's Amazing Camels kindly provided information on their camel herd and its operations.

### **Funding statement**

No external funds were used in the preparation of this work.

### **Conflict of interest disclosure**

The Author declares no conflict of interest.

### **Ethics approval statement**

Not applicable.

### **Patient consent statement**

Not applicable.

### **Permission to reproduce material from other sources**

Not applicable.

### **References**

- Barnes RH. Charles George Gordon, a sketch. MacMillan, London. 1885.
- Boulger DC. The Life of Gordon, Major-General, R. E., C. B.: Turkish Field-Marshal, Grand Cordon Medjidieh, and Pasha; Chinese Titu (Field-Marshal), Yellow Jacket Order [3<sup>rd</sup> Ed.]. 1897. T. Fisher Unwin, London.
- Clayton A and Killingray D. Khaki and Blue: Military and Police in British Colonial Africa. Africa series. Monographs in International Studies. 1989; Volume

51. Ohio University Center for International Studies, Athens, OH.
- Darke J. *The Monument Guide to England and Wales*. Macdonald Illustrated, London. 1991. ISBN 0-356-17609-6.
- Dickens C (jr). *Dickens's Dictionary of the Thames, From its Source to the Nore, An Unconventional Handbook*. 1885. Macmillan & Co., London.
- Droth M, Edwards J and Hatt M . *Sculpture Victorious: Art in the Age of Invention, 1837-1901*. Yale Center for British Art, Yale University Press, Newhaven CT. 2014. ISBN 9780300208030.
- Ekwall E. *The Concise Dictionary of English Place-names* (2nd edition). Clarendon Press, Oxford. 1940.
- Falls C and MacMunn G. *Military Operations Egypt & Palestine from the Outbreak of War with Germany to June 1917*. Official History of the Great War Based on Official Documents by Direction of the Historical Section of the Committee of Imperial Defence. Volume 1. His Majesty's Stationery Office, London. 1930. OCLC 610273484.
- Galbraith J.Gordon, Mackinnon, and Leopold: *The Scramble for Africa, 1876-84*. *Victorian Studies*. 1971; 14(4):369-388.
- Hsu I. *Gordon in China, 1880*. *Pacific Historical Review*. 1964; 33(2): 147-166.
- LGU. *The Dangerous Wild Animals Act 1976 (Modification) (No.2) Order 2007*. UK Statutory Instruments 2007 No. 2465. Legislation Government. UK. 1976 <https://www.legislation.gov.uk/ukxi/2007/2465/memorandum/contents>.
- LGU. *Wild Animals in Circuses Act 2019* UK Public General Acts 2019 c. 24. Legislation Government. UK. 2019. <https://www.legislation.gov.uk/ukpga/2019/24/contents/enacted>.
- Moore-Harell A. *Slave trade in the Sudan in the 19th century and its suppression in the years 1877-80*. *Middle Eastern Studies*. 1998; 34(2):113-128. doi:10.1080/00263209808701225
- Mills AD. *A Dictionary of British Place-Names*. Oxford University Press, Oxford. 2003. ISBN 9780191578472.
- Mynors RAB, Winterbottom M and Thomson RM (eds and translators). *Gesta Regum Anglorum: The History of the English Kings*, vol. 1 by William, of Malmesbury, ca. 1090-1143. Oxford Medieval Texts, Clarendon Press, Oxford. 1998.
- NZMCH. *New Zealand History online*. History Group of the New Zealand Ministry for Culture and Heritage, Wellington. N.d.
- Pascoe WH. *A Cornish Armory*. Lodenek Press, Padstow, Cornwall. 1979. ISBN 0-902899-76-7.
- Reid F. *The Fighting Cameliars* (Number 7 in Series: The Gallant Legion). 1934. Angus & Robertson, Sydney.
- Risebero W. *Modern Architecture and Design: An Alternative History*, MIT Press, Cambridge MA. 1985. ISBN 0-262-68046-7. pp 91.
- Robertson J. *With the Cameliars in Palestine*. Reed Publishing (NZ) Ltd, Dunedin. 1938.
- Searle WG, Christ Church, Canterbury: *I. The Chronicle of John Stone*. Cambridge Antiquarian Society, Cambridge. 1902.
- Sparrow G. *Gordon: Mandarin and Pasha*. Jarrolds: London. 1962.
- Trench CC. *The Road to Khartoum: A Life of General Charles Gordon*. W.W. Norton & Company, New York NY. 1979. ISBN 10: 0393012379.
- Underwood J . *The organisation of the Imperial Camel Brigade 1916-1918*. *Sabretache Journal of the Military Historical Society of Australia*. 2003; 4:5-14.
- Ward-Jackson P . *Public Sculpture of Historic Westminster: Volume 1*. Liverpool University Press, Liverpool. 2011. 355-357.
- Wilson E. *Cleopatra's Needle: With Brief Notes on Egypt and Egyptian Obelisks*. Brain & Company, London. 1877.
- Ziegler P. *Omdurman*. Collins: London. 1973.

## Appendix A

### Prince Consort National Memorial (Albert Memorial), Westminster

Heritage Category: Listed Building // Grade: I // List Entry Number: 1217741 // Date first listed: 14-Jan-1970 // List Entry Name: PRINCE CONSORT NATIONAL MEMORIAL (ALBERT MEMORIAL) // Statutory Address: PRINCE CONSORT NATIONAL MEMORIAL (ALBERT MEMORIAL), KENSINGTON GORE SW7

TQ 2679 NE CITY OF WESTMINSTER KENSINGTON GORE, SW7. 85/2

14.1.70 Prince Consort National Memorial (Albert Memorial). GV I

Memorial. 1862 to 1875. George Gilbert Scott, with sculpture and other decorations by numerous Victorian artists and craftsmen. Granite, marble, limestone and metal. Square decoratively paved podium reached by steps to south; superimposed podia with steps to all sides, surmounted by ciborium-like canopy with clustered granite columns to four corners. Elaborate sculptured spire. Giant seated figure of Prince Albert within. Gables to all sides with pinnacles between. Richly sculptured band to base breaking outward at corners to support sculptured groups representing the industrial arts:- Agriculture, Manufacture, Commerce and Engineering. Further sculptured groups to four corners of stepped podium representing Europe, Asia, Africa and America, and linked by elaborate railings. Monument enriched with much figure and other sculpture and decorated also in numerous other techniques including mosaic, enamel, etc.

## Appendix B

### Imperial Camel Corps Memorial

Heritage Category: Listed Building // Grade: II // List Entry Number: 1238057 // Date first listed: 24-Feb-1958 // List Entry Name: IMPERIAL CAMEL CORPS MEMORIAL // Statutory Address: IMPERIAL CAMEL CORPS MEMORIAL, VICTORIA EMBANKMENT GARDENS WC2

TQ 3080 NW CITY OF WESTMINSTER VICTORIA EMBANKMENT 72/148 GARDENS, WC2 Imperial Camel Corps 24.2.58 Memorial G.V. II .

Memorial. 1920 by Cecil Brown. Bronze statuette of figure mounted on camel set on Portland stone pedestal with bronze bas-relief.

## Appendix C

### Memorial to General Gordon, Brompton Barracks

Heritage Category: Listed Building // Grade: II\* // List Entry Number: 1375610 // Date first listed: 07-Jul-1998 // List Entry Name: MEMORIAL TO GENERAL GORDON, BROMPTON BARRACKS // Statutory Address: MEMORIAL TO GENERAL GORDON, BROMPTON BARRACKS, PASLEY ROAD

TQ 7669 SW GILLINGHAM PASLEY ROAD (east side), Brompton 3/32 Memorial to General Gordon, Brompton Barracks GV II\*

Equestrian statue. 1890. Bronze statue and Portland stone plinth. Moulded plinth, with a bronze cartouche, with a full-sized figure of General Gordon, d.1885, seated on a camel with tasselled saddle. Attached a well-cast and expressive figure with free-hanging tassels; placed symmetrically on the axis with the Institute, memorial arches and barracks (qqv).

## Appendix D

### Statue of General Gordon, Gordon's School

Heritage Category: Listed Building // Grade: II // List Entry Number: 1424607 // Date first listed: 29-Apr-2015 // List Entry Name: Statue of General Gordon, Gordon's School // Statutory Address: Gordons School, Bagshot Road, West End, Woking, GU24 9PT

Statue on plinth. A 1902 second casting of the sculptor Edward Onslow Ford's statue of General Gordon (first cast in 1890). The plinth is of circa 1959. Bronze statue on stone plinth. A thirteen feet high, bronze, full-sized statue of General Gordon in the uniform of an Egyptian general, seated on a tasselled saddle upon a standing camel. The base is inscribed 'GORDON' on the front and has a decoration of olive branches. The classical style rectangular stone plinth has recessed corner piers, a moulded cornice and base. An inscription on one side of the plinth reads 'THIS STATUE WAS ERECTED AT KHARTOUM/ IN 1904 AND PRESENTED TO/ THE SCHOOL IN 1959.' An inscription on the base of the plinth records that the plinth was presented in memory of Major General Sir Hubert J Huddleston who served the Sudan for 23 years. The statue of General Gordon at Gordon's School by the sculptor Edward Onslow Ford RA, cast in 1902 with a 1959 plinth, is listed at Grade II for the following principal reasons: Aesthetic quality: a fine bronze sculpture by a reputable sculptor with the subject unusually mounted on a camel; Historic interest: General Gordon is a figure of national importance and this statue is located at the school founded in his memory; Group value: with other listed buildings and structures at the school.

## Appendix E

### Bench seats on Victoria Embankment

Heritage Category: Listed Building // Grade: II // List Entry Number: 1357348 // Date first listed: 01-Dec-1987 // List Entry Name: 21 BENCH SEATS SET ON EMBANKMENT

PAVEMENT // Statutory Address: 21 BENCH SEATS SET ON EMBANKMENT PAVEMENT, VICTORIA EMBANKMENT SW1

TQ 3079 NW, 3080 SW and NE CITY OF WESTMINSTER VICTORIA EMBANKMENT, 3180 NW SW1 73/57;74/19;83/61;92/12 21 Bench Seats set on Embankment Pavement G.V. II Public seats. 1872-74 designed by Lewis and G F Vulliamy. Cast iron mounted and timber slatted. The seats take the form of long benches with curved backs supported on cast iron brackets, the terminal arm-brackets designed as winged sphinxes, with the exception of the bench opposite the junction with Horseguards Avenue which has the feature of seated camels instead of sphinxes. Included in their own right as original street furniture and as part of the Embankment design.

## Appendix F

### Library, Market Place, Camelford, Cornwall

Heritage Category: Listed Building // Grade: II // List Entry Number: 1138348 // Date first listed: 19-Jan-1952 // Date of most recent amendment 13-Jan-1988 // List Entry Name: LIBRARY // Statutory Address: LIBRARY, MARKET PLACE

CAMELFORD MARKET PLACE, Camelford SX1083 - 1183 8/76 (1/3) Library (previously listed as Town Hall - 19.1.52) GV II. Townhall erected over market house, now library, exhibition hall and meeting room. 1806. Financed by John, 6<sup>th</sup> Duke of Bedford. Slate stone rubble with dressed quoins and dressed stone arches to window openings. Rag slate roof with hipped ends. Rectangular plan with internal partitions altered. Originally the market house on the ground floor with a central entrance on the front and the town hall above with a staircase to a doorway at the rear where the ground is at a higher level. Ground rises to left and to rear. 2 storeys. Symmetrical 3 window front faces onto Market Place; 5 semi-circular headed windows; the window frames replaced in circa mid C20, the central window in an earlier blocked door opening. First floor has 3 semi-circular windows with horned sashes with intersecting glazing bars. Lower side on right has a C20 glazed door with a Venetian window above, the central horned sash with intersecting glazing bars. The higher side on the left has a C20 door on the ground floor and a Venetian window above with stained glass window with Camelford arms. The wall above has been partly rebuilt. The rear elevation has 4 blocked round headed openings on the ground floor and a central double flight of steps which lead up to the first floor entrance with a C19 timber porch with flat roof and moulded cornice. 2 sash windows with round headed arches and intersecting glazing bars to right and left. Timber cupola on roof with clock and weather vane. This cupola originally had 2 bells dated 1807 and 1699. Interior : remodelled in circa mid C20. Bell from cupola in entrance inscribed 'W. Prideaux Mayor, 1699'. The Charter of Camelford was confirmed in 1259 and the rotten borough was created in 1552 to bolster the Earl of Northumberland's government. The rotten borough was abolished in 1831( Kelly's Directory, 1910. Polsue, J. Lake's Parochial History of the County of Cornwall 1872 Volume I).

## **16<sup>th</sup> ANNUAL SOUTHWEST CAMEL CONFERENCE AND TRAINING CLINIC AT LAS VEGAS**

16<sup>th</sup> Annual Southwest Camel Conference and Training Clinic will be organised at Las Vegas, USA in partnership with North American Camel Ranch Owners Association (NACROA) from 25-27<sup>th</sup> October, 2024. The conference will involve guest speakers, training sessions and many aspects of camels. Various presentations in the conference would cover camel welfare, safe transport, international resources, general health assessments and advancing the camel science knowledge. The contact persons are Doug Baum (texascamelcorps@gmail.com) and Valeri Crenshaw (crenshawvaleri@hotmail.com). Registration fee for NACROA members is \$ 600 and for non-members it is \$750. Camels may attend for free. Registration/payment will be open from August 1<sup>st</sup> to 1<sup>st</sup> October 2024.

## **BEYOND DESERTS AND HIGHLANDS- held at Vienna International Centre, Vienna, Austria on 10 June 2024**

The Joint FAO/IAEA Centre organised an international on "Beyond Deserts and Highlands- Global Celebration of the International Year of Camelids 2024". It was in response to the growing global interest and the need for a collaborative platform that brings together experts, researchers, policymakers, and breeders to share knowledge, discuss challenges, and explore innovative solutions. The event aligned with the global agenda for promoting biodiversity conservation, given the unique ecological niches that camelids occupy. Recognising the potential of camelids in sustainable agriculture, ecotourism, and poverty alleviation, The event sought to foster international collaboration and partnerships that will contribute to the well-being of both camelid-dependent communities and the broader global population.

The objectives of the event were to facilitate the exchange of cutting-edge research, technological advancements, and best practices in camelid science, husbandry, and management, to provide a platform for participants to network with fellow researchers, breeders and policymakers, to share knowledge on various aspects of camelid husbandry, including genetics, nutrition, health, and socio-economic impacts and to raise awareness about the significance of camelids in diverse fields and advocate for their sustainable integration into global agricultural and economic systems. Keynote address by Dr B. Faye was in the field of camelid research and husbandry. Technical Sessions covered diverse topics such as genetics, nutrition, health, socio-economic aspects, and cultural significance of camelids. There were panel discussion on emerging challenges and opportunities in camelid management together with an exhibition.

## **CAMEL HIDES ARE SECOND ONLY TO CROCODILE HIDES**

The camel industry is expected to become one of Saudi Arabia's foremost non-oil resources if it achieves its targets in accordance with the Kingdom's Vision 2030 plan. Currently, a company called Sawani, affiliated with the Public Investment Fund, focuses on investing in the transformative camel industry, including liquid and powdered camel milk. Their products have already been exported to over 25 countries worldwide. They have utilised camel milk in the production of "Camelicious" ice cream, available in eight flavours, as well as in the cosmetics industry. In addition to Sawani, the Saudi brand "Ibil" established in 2021, has been involved in manufacturing clothing, bags, and shoes using camel wool and hides. Their aim is to become a global brand within 15 years. It is a known fact that camel hides are durable and long-lasting, second only to crocodile hides. Even their bones are used in construction practices. The camel plays a key role in the Saudi tourism sector.

(Source: <https://arab.news/222ke>, January 2024)



# CAMELOLOGY : DEFINITIONS, HISTORY AND SCIENTIFIC CHALLENGES

Bernard Faye<sup>1</sup> and Tarun Kumar Gahlot<sup>2</sup>

<sup>1</sup>Emeritus scientist, CIRAD-ES, UMR SELMET TA/C 112A, Campus International de Baillarguet, Montpellier, France

<sup>2</sup>Editor, Journal of Camel Practice and Research and Professor,  
MJF College of Veterinary and Animal Science, Chomu, Jaipur, India

## ABSTRACT

Since centuries, the camel is fascinating scientists over the world. If the first scientific investigations were focused on his remarkable adaptability to desert conditions and to his health management in the context of colonial expansion of European countries in desert areas, recent researches have mobilised several specific disciplines as archaeology, physiology, immunology, breeding and genetics, parasitology, surgery, imaging, diseases, food sciences, economy, or sociology. However, the camel can be regarded as a “full scientific object” legitimising the use of the word “camelology”. Present paper is an overview of chronological development of camel cultures, production and science.

**Key words:** Adaptation, camel, science history, scientific approach

The concept of “camel science” or “camel research” is commonly used by the scientists working on camel as “object of study”, but as researcher, they decline their speciality as parasitologist, or anatomist, or immunologist or those of other specialities. Moreover, in language other than English, the literal translation is usually not correct. In French for example, the translation of the sentence “I’m camel scientist” will mean “I am a scientist with a camel character”. Unless you use a periphrasis (“I am a scientist studying camel”), your interlocutor will not understand you properly. It is the reason why the concept of “camelology” is better to be used. In French, the word “camelologie” was reported for the first time on the website <https://camelides.cirad.fr> in 2002 as a neologism regarding camel among other words as “camelodrome” or “cameliculture”. Few times later, the word was discussed to be included in the dictionary of the French Academy. It was therefore, the humble reply was “I’m a camelologist” when people asked for our speciality.

The concept of camelology needs to be elaborated. If camelology is a discipline, we have to define the history and the field or scope of this science. The present paper is specially proposed for the international year of camelids (2024) and it aims to investigate the chronological development of camel culture and science from its primitive form to a modern but updated format.

## A brief history of the camelology in the world


### • The pioneers

Georges-Louis Leclerc, Count of Buffon, known usually under the single name of Buffon, a French aristocrat living in the XVIIIth century, was probably one of the first “naturalist” describing the camel in his book “Histoire Générale des animaux” (General history of the animals), written between 1753 and 1767. Dromedary and Camel (in fact Bactrian camel) anatomy, physiology, feeding behaviour, geographical distribution and utilisations were widely described with accuracy (Fig 1). It was the first scientific description of the large camelids since the succinct mentions of Aristote in his “history of animals” written in 343 BC (Gallimard Ed., Paris, 1994). There appears to be no specific mention of camel in the studies or publications of the most famous Arabic scholar during the middle age (except Ibn Al-‘Awwâm, the master of Moorish Agronomy-Pereira, 2024) although it was mentioned several times in the Holy Koran.

In the XIXth century, Alexandre Vallon, a French veterinarian from the imperial School of Cavalry, published a book on the natural history of camel (Vallon, 1856). But, the first scientific investigations regarding camel could be dated to the colonial period, especially when the French and English troops were invading North Africa and Sahara from Mauritania to Somaliland, especially

SEND REPRINT REQUEST TO BERNARD FAYE [email: bjfaye50@gmail.com](mailto:bjfaye50@gmail.com)





**Fig 1.** The “dromedary” and the “camel” (Bactrian) represented in the volume VI of the General history of the animals (*“les quadrupèdes”*) written by Buffon in the XVIII<sup>th</sup> century.

with the support of “Meharist companies” and other “camel corps” where military veterinarians played military and medical role.

Indeed, the first publications were mainly focused on camel diseases. Among the first books regarding camel, one could cite that of Cross (1917), a British vet serving as “camel specialist” to the Punjab Government, in India. Later, another British vet serving in India, then in Somaliland, Arnold Leese, was known for his wide experience regarding camel diseases and management, before known to be as a fascist activist. His book, “A Treatise on the Humped Camel” was published in 1927 (Leese, 1927), but previous papers were published in the Journal of Tropical Veterinary Science from 1909. The Italian military vets contributing to the colonial invasion of Libya and Somalia were not left out: Ivo Droandi who published many papers from 1920 in the Italian Colonial Institute of Agriculture (Florence) on camel surgery and diseases, published a book in 1936, after his experience in Libya. In France, the first military group using camel was created by Napoleon Bonaparte during the Egyptian campaign in 1799, but the “Meharist company” was established in 1902 in Sahara and some description regarding camel management were proposed by the vets accompanying the soldiers, as it was the case of the commandant Cauvet (Cauvet, 1925). However, the first comprehensive book regarding camel diseases was written by Gaston Curasson, a French general inspector of Veterinary Services in French colonies

(Curasson, 1947). In the same period, some books focused on generalities regarding the life of camel were published, for example, in France (Finbert, 1938).

#### • *The mysteries of the camel physiology*

After the second world war and before the independence of the colonies in the years 60, the first studies on camel were focused on the physiology of adaptation, notably the works of Knut Schmidt-Nielsen (Norwegian) and his wife Bodil (American scientist) who established the bases of the camel physiology, for example on water balance or urea excretion (Schmidt-Nielsen *et al*, 1956 and 1957). At the same time, Hilde Gauthier-Pilters started publications on camel ecology in 1958 and gathered all these observations in a book in 1981 (Gauthier-Pilters and Dagg, 1981). Physiological studies, especially regarding metabolism were continued by Emmanuel from the University of Shiraz, in Iran (Emmanuel, 1979). Renal function and physiology were later studied in details by Dr Reuven Yagil, Israel (Etzion and Yagil, 1986) and Dr N.Kataria from India (Kataria *et al*, 2002 a,b,c,d,e; 2003 and 2007). Physiology of forestomach of camelids was studied in detail by Engelhardt *et al* (1988). Physiology of racing camels was studied by Saltin and Rose (1994). More recently, “Camel Clinical Biochemistry and Haematology” was another important publication in 2018 by Bernard Faye and Mohammed Bengoumi. In Morocco, the University of Casablanca and Agro-veterinary Institute at Rabat, achieved many studies

in collaboration on water and mineral metabolism (Bengoumi *et al*, 1993; Riad *et al*, 1994). Later, the team of Casablanca University was renowned for its work on camel stress (Lemrahmed *et al*, 2019).

• ***The emergence of the camelology in the southern countries: Research institutes, specialised journals and general books***

After most of independences following the colonial time, more and more scientists from southern countries, especially from Egypt, Sudan, India, Morocco started to publish papers on camel anatomy, physiology and diseases. Some institutions began to be recognised for their research activities on camel, i.e. College of Veterinary and Animal Science, Bikaner and the National Camel Research Centre (NRCC), Bikaner, India, University of Alexandria and the Desert Research Centre (DTC) in Egypt and University of Khartoum in Sudan, Agro-veterinary Institute (IAV) in Morocco and later the Institute of Arid Lands (IRA) in Tunisia. Meanwhile, the Central Veterinary Research Laboratory (CVRL) was founded in 1985 by HH Sheikh Mohammed Bin Rashid Al-Maktoum, Vice President and Prime Minister of the UAE and it was a Government diagnostic centre that provided testing and research facilities on camels and other animals including falcons and raptors, to the UAE and neighbouring countries. CVRL is the OIE Reference laboratory for Glanders, Camel Pox, Brucellosis (*Brucella abortus*, *B. melitensis*, *B. canis*) and MERS (Middle East Respiratory Syndrome). As Scientific Director of CVRL, Dr. Wernery leads to his disease diagnosis research in camels among other species. He and the CVRL team pride themselves on being at the forefront of camel research worldwide, especially in veterinary field. In the year 2000, the World Animal Health Organisation (OIE) in Paris, decided to create a camel disease ad-hoc group gathering some camel experts in order to define the list of infectious diseases, their convenient diagnosis tools and the list of reference labs throughout the world. Dr Wernery with the authors of the present paper were among the first experts contributing to this group.

The Camel Applied Research and Development Network (CARDN) was created at Damascus (Syria) in 1991 under the umbrella of the Arab Centre for the studies of Arid Zones and Drylands (ACSAD), publishing a Camel Newsletter, then the first specific journal devoted to camel, the Journal of Camel Sciences (Wardeh, 1986) which unfortunately was stopped after few issues. In India, a book on Camels

and their Management was published in 1986 by Rathore (Rathore, 1986). The book "The Camel" authored by R. Trevor Wilson in UK (Wilson, 1984) was published as an important reference book. In 1984, too, the British Veterinary Journal also published a series of papers on Camel in Health and Disease (Higgins and Kock, 1984). In 1986, another book on camel in health and disease was authored by Higgins (1986). Then, Camel Publishing House at Bikaner (India) started publishing the Journal of Camel Practice and Research since 1994 as biannual but later in 2017, it became a triannual journal (Gahlot, 1994). In the year 2000 onwards, "The Camelid Publishers" brought out 3 important publications edited by T.K. Gahlot from India (Gahlot, 2000, 2004 and Gahlot *et al*, 2002). Compilations of the papers published in the Journal of Camel Practice and Research in the field of parasitology (Gahlot and Chhabra, 2009), gross anatomy and histology (Gahlot *et al*, 2011) and immunology (Gahlot *et al*, 2016) were proposed by Camel Publishing House. In Israel, Yagil and his team from University of Negev published many papers on physiology of adaptation and contributed also by a book on camel physiology (Yagil, 1985) to advance the knowledge of the camel. In Soviet Union also, camel sciences were implemented, especially in genetic and milk production and many books were published in Russian language (Terentyev, 1975). Few books on camel surgery were also published (Gahlot and Chouhan, 1992; Ramadan, 1994; Siddiqui and Telfah, 2010).

Scientists from the French speaking countries in Africa also started to publish the papers on camel science in the Revue d'Élevage et de Médecine Vétérinaire des pays Tropicaux (REMVT) which was integrated to CIRAD in 1996 where the coordination unit on camel breeding (UCEC in French) was created. A first comprehensive book was edited by Richard in 1984 (Hoste *et al*, 1984) and a special issue of REMVT was published in 1989 ("Le dromadaire", n°1, 1989). Then, Faye and other contributors participated to one book as support for new line of veterinary medicine proposed by SANOFI © ("Guide l'élevage du dromadaire", 1997), available also in Arabic. In Germany, in 1992, the book was published on the "One-Humped Camel In East-Africa" (Schwartz and Dioli, 1992) and in Pakistan, another book was edited (Chaudhary and Akbar, 2000). From this period, many books were published in local languages in Mongolia (Indra *et al*, 2003; Adams, 2024), China (Zhao Xin Xu, 1996), Kazakhstan (Moussaiev *et al*,

2007) and Germany (Weiss and Wernery, 2021). Recently, the French general book “Elevage des grands camélidés” (Faye *et al*, 2022) was translated in Turkish, Arabic, Spanish, Kazakh and English (Faye *et al*, 2023).

#### • The camel conferences

During the post-independence period, there were few camel conferences. These were held in the year 1970-90s in Khartoum (Sudan), Tobruk (Libya) and later in Paris (France) on the reproduction (1990) and in Nouakchott (Mauritania) on camel milk (1994) by UCEC (Saint-Martin, 1993; Bonnet, 1998), Israel (1996), Al-Ain (1998), Algeria (1989) and Kazakhstan (2000). However, one of the first important organised camel conference was at Dubai in 1992, concomitant with emergence of advanced research in the Gulf countries on the physiology of racing camel (Saltin and Rose, 1994), biotechnology of reproduction (Tibary and Anouassi, 1997) and camel diseases (Wernery and Kaaden, 1995).

In some cases, more specialised workshops were organised. For example in Morocco (1999) on the young camel and in 2000 on camel diseases (Dakkak, 2000), in Niger (2003) on camel milk in Africa organised by FAO (Lhoste, 2003), in Turkmenistan in 2004 on the role of camel in desertification combat (Faye and Esenov, 2005). Other conferences were organised in UAE (1991), India (2004 and 2007), Saudi Arabia (2007), Kenya (2010) and many others that we cannot cite all these here. The proceedings of those conferences have significantly added to the camelid literature.

In 2006, four camel scientists (B. Faye, M. Bengoumi, K. Alhadrami, A. Tibary) created the International Society of Camelid Research and Development (ISOCARD) with the objective to gather all the camelid scientists (camelologists) of the world in triennial international conferences. Nowadays, 6 conferences were organised, i.e. in 2006 (Al-Ain-UAE), 2009 (Djerba-Tunisia), 2012 (Muscat, Oman), 2015 (Almaty, Kazakhstan), 2018 (Laayoune, Morocco) and 2023 (Al-Ahsa, KSA). The Society also published an on-line journal (Journal of Camelids Sciences) available on line on the website of the society. Some of the proceedings of the ISOCARD conferences are also available on the website (Johnson *et al*, 2012; Konuspayeva, 2015; Sghiri and Kichou, 2018).

#### Some advanced camel researches

It is not possible to be exhaustive and obviously, some important topics regarding camelology are not

mentioned in detail here. The authors assume that the selected topic below are subjective.

#### • Immunology

Camel immunology has experienced important progress (Hussen and Schuberth, 2021), One of the most remarkable features of the camelid family is their unique immune system. A break-through in the camelid humoral immune system was achieved by Hamers-Casterman *et al* in 1993, who showed that the camel possesses novel class single-domain antigen binding fragments. These proteins are the naturally occurring antigen-binding domains known today as VHHs (single variable heavy-chain) or nanobodies. These nanobodies are very small and possess several bio-physical properties, that offer great advantages in various medical and biotechnological applications (Muyldermans and Lauwereys, 1999), including cancer treatment (Al-Numair *et al*, 2022), producing hyperimmune serum against snake bites and more (Tanwar *et al*, 2017). These extra ordinary camelid nanobody particularities were shown in a recent documentary by Marc Jampolsky film, DW documentary. A.K. Kataria and co-researchers from India carried out important research and studies on dromedary immunoglobulins (Kataria *et al*, 1994, 1999, 2002; Kataria, 2001; Kataria and Sharma, 1999, 2000, 2003; Kataria and Kataria, 2004).

#### Genetics and genomics

Since the years 2000, important advanced researches were performed on genetics and genomics (Burger *et al*, 2019) allowing new knowledge regarding camel biodiversity (Gaouar and Ciani, 2023), history of domestication (Almathen *et al*, 2014) or selection (Al-Abri and Faye, 2019). Scientists from Riyadh's King Abdulaziz City for Science and Technology and China's Shenzhen-based BGI (formerly the Beijing Genomics Institute) were able to decode the entire genetic makeup of the single-humped camel (*Camelus dromedarius*), thus the Arabian camel today enters a highly exclusive club of selected few mammals which have had their full genome sequenced and analysed. Genomic sequencing and analysis of eight camel-derived middle east respiratory syndrome coronavirus (MERS-CoV) isolates was done in Saudi Arabia (Al-Shomrani *et al*, 2020). Chinese scientists found origin and migration of domestic and wild Bactrian camels through genome sequencing (Jirimutu *et al*, 2012; Ming *et al*, 2020). The preservation of the camel biodiversity as an important element of the sustainable development, especially among



pastoralists is also recommended in several research papers (Kohler-Rollefson, 2022 and 2023).

#### • *Camel milk and meat studies*

Considerable progress was done in the knowledge of camel milk gross composition and fine composition of camel milk since the pioneer work of Farah (1993). A first book on milk and meat composition and processing milk was published in 2004 by Farah and Fisher. The first meta-analysis of camel milk composition was proposed by Konuspayeva *et al* (2009). The fine composition was explored by many scientists emphasising the particularities of camel milk (El-Agamy, 2017). Based on the research of Kappeler *et al*, on the camel chymosin structure in 2006, significant progress were done on the camel cheese making (Konuspayeva *et al*, 2017) and knowledge of milk microflora (Kaindi and Njage, 2020). With the emergence of the camel milk industry, especially in the Gulf countries and Central Asia, notably the establishment of Emirates Industry For Camel Milk & Products (EICMP) in the year 2006, camel milk products appeared on the market (including at international level thanks to the EU agreement) under more diversified form (Konuspayeva and Faye, 2021). The implementation of big camel dairy farms as Camelicious (EICMP) gathering thousands camel under similar environment has boosted original research on camel milk production, milk microbiology and milking management (for example, Nagy *et al*, 2013a and 2015). Intensification of the camel milk production (Nagy *et al*, 2022) by using machine milking has also boosted research on the physiology of lactation (Ayadi *et al*, 2016) and adaptation of camel to machine milking (Atigui *et al*, 2014; Nagy and Juhasz, 2016; Kaskous, 2023). The health effect linked to the regular consumption of camel milk was explored by many camel scientists with variable scientific rigour as mentioned in the recent review of Faye and Konuspayeva (2024). The impact on human diseases as autism (Adams, 2019; Gahlot and Adams, 2023), diabetes (Ashraf *et al*, 2021; Alkhurd *et al*, 2022), Crohn's disease (Rosenheck *et al*, 2012) or even some cancers (Badawy *et al*, 2021; Lal *et al*, 2023) are among the numerous recent *in vitro* and *in vivo* investigations. In addition, the interest of health effect of camel products was extended to the camel urine. Its potential anticancer, antiplatelet, gastroprotective and hepatoprotective effects were explored by different scientists in the world (Salamt *et al*, 2021). A synthetic book on the health and environmental benefits of camel products in general was published in 2020 (Alhaj *et al*, 2020).

Camel meat studies have shown recent interest due to its dietetic properties (Raiymbek *et al*, 2015 and 2018). Meat studies included research on the meat composition (Kadim *et al*, 2008), meat processing (Baba *et al*, 2021), slaughtering conditions (Moussahil *et al*, 2022) and even the health benefit of its consumption (Kadim *et al*, 2022). A synthetic book was published in 2013 (Kadim *et al*, 2013) gathering all aspects of camel meat industry, from the production to the consumption.

#### • *Biotechnology of reproduction and cloning*

Since the first conference on camel reproduction, held in Paris and cited above, many advanced researches were achieved especially in the Gulf countries (Skidmore *et al*, 2024), India (Purohit *et al*, 2023) or Egypt (El-Bahrawy *et al*, 2015). Artificial insemination and overall embryo-transfer are nowadays used routinely in intensive big farms (Nagy *et al*, 2013b). World's first cloned camel named "Injaz" was produced in year 2009 (Wani *et al*, 2010). A first report on interspecies embryo transfer which gave birth to Bactrian camel calves from dromedary camels, was published by the scientists of Iran (Niasari-Naslaji *et al*, 2009). Dozens of cloned dromedary camels have been produced from the embryos reconstructed with cells from racing champions, winners of beauty contestants, high milk yielders and elite bulls (Wani, 2021). The first cloned Bactrian camel calf was produced by interspecies SCNT using dromedary camel as a source for oocytes as well as a surrogate for carrying the pregnancy to term (Wani *et al*, 2017). Multiple cloned camels from racing, show and dairy exemplars were produced in a recent research (Olsson *et al*, 2021).

#### • *Camel health and welfare*

Health is one of the most important constraints in camel farming, both in traditional production and intensive system. Thus, it is not surprising to see that veterinary science was the most cited topic in the scientific literature devoted to camel as mentioned in the different scientometric studies assessing the number of publications, the different topics and the main journals, authors or institutions contributing to the camel sciences (Faye *et al*, 2000; Rathinasabapathy and Rajendran, 2013; Gupta *et al*, 2015; Kandeel *et al*, 2023). Several books or review papers are nowadays available for giving comprehensive information on infectious diseases (Wernery and Kaaden, 1995; Wernery *et al*, 2014), zoonotic diseases (Khalafalla, 2023) and even global health disorders affecting camels notably in traditional farming systems



(Kohler-Rollefson *et al*, 2001). The emergence of MERS-CoV in Middle East and the role of dromedary camel in the disease transmission has stimulated an important volume of research on virology and epidemiology (Ngere *et al*, 2020; Azhar *et al*, 2023). Few studies based on ecopathological approach of multifactorial diseases as calf diarrhoea were achieved (Bengoumi *et al*, 2003).

Camel welfare is an important emerging topic, formerly focused on the assessment of the stress in different conditions (dehydration, heat stroke, transport, slaughtering...) and nowadays, more and more is focussed on good management practices and globally good farming conditions for the camel (Menchetii *et al*, 2021). Recently, a book was gathering the knowledge regarding this topic including health management, environmental conditions, handling and assessment of behaviour (Padalino and Faye, 2024).

### Camel celebrations

World Camel Day is celebrated on 22nd June in many countries to raise awareness about camels and their importance to human societies and ecosystems and to celebrate these unique creatures. United Nations has declared the year 2024 as International Year of Camelids. Several organisations are celebrating it in form of meetings, workshop or conferences as LPPS (Lokhit Pashu Palak Sansthan) in India; CAMENET (Middle East Camel Network); NACROA (North America Camel Ranch Owner Association), IAEA (International Atomic Energy Agency) in Austria, many universities throughout the world (Algeria, Morocco, Tunisia, Saudi Arabia, UAE, Kuwait...) and of course, FAO.

### The camelology, a science?

In a world where disciplinary approaches are no longer enough to understand complexity, considering a livestock species such as the camel, in all its dimensions (from the cell functioning to the farming system or desert ecosystem, passing by metabolic functions and behaviour, diseases and cognition) can be considered a scientific challenge by itself. In this context, the camel is not only an animal, but an object of study that responds to 3 levels of investigation: (i) the camel as biological model, notably for medicine and adaptative studies, (ii) the camel as productive animal in desert areas or in intensive farms for providing milk, meat and other services for humanity and (iii) the camel as an element of complex desert ecosystem in relationships with changing environment, economic challenges

and human or pastoral society (Faye and Brey, 2005; Kohler-Rollefson, 2023 and Rollefson, 2004). Considering camel from such angle, the camelology faces to different challenges that can be summarised into few non-comprehensive topics as:

- To deepen the mechanisms explaining the health effect of camel milk and meat consumption
- To achieve convenient clinical trials to support the health claim of the consumption of camel products
- To implement efficient systems of selection and of feeding for improving the camel productivity (production, reproduction) in the frame of the specialisation of the farming systems
- To increase the studies for allowing sustainable development of more intensive camel systems by respecting animal welfare and health
- To investigate the multifactorial diseases by using more holistic approach and identifying the individual and environmental risk factors
- To understand and characterise the role of camel extensive systems in the desertification combat
- To evaluate the place of camel in the micro-meso- and macro-economy worldwide

And many other topics...

### Conclusion

The camel, irremediably linked to arid ecosystems, has a specific physiology and metabolism allowing its adaptation, but this desert animal is also fundamentally an element of the economy in those ecosystems and of the local culture. Moreover, it allows adding zootechnical value for people, pastoralists or settlers, living in such harsh environment. Another very important fact should be highlighted that the ship of the desert is an environmentally friendly species unlike bovines, sheep and especially goats. Reducing camel numbers on open-range land could redress desertification by allowing vegetation to recover from overgrazing. Prototype camel farming should encourage owners, to move most of their camels from open range into intensive farming. Initiating camel farms, where camels are fed on native deserts or salt-tolerant plants may help further the restoration of the ecosystem (Breulmann *et al*, 2007; Kohler-Rollefson, 2023). Like

“eremology”, the science of desert (Heuse, 1992), the camelology can consider the camel as a “full scientific object”. The camel scientists should go beyond their specific disciplines for a better understanding of the camel in his all dimensions.

## References

- Adams C. Camel Crazy: A Quest for Miracles in the Mysterious World of Camels. 2019. New World Library, 304 pages. ISBN 978-1-60868-648-3.
- Adams C. Camel Crazy, Mongolian Language Translation Book. 2024. ISBN 991952292-6, Nepko Publishing, Ulaanbaatar, Mongolia.
- Al-Abri M and Faye B. Genetic improvement in dromedary camels: challenges and opportunities. *Front. Genet.* 2019; 10(167):1-5; DOI: 10.3389/fgene.2019.00167
- Al-Haj O, Faye B and Agrawal RD (Eds). Handbook of Research on Health and Environmental Benefits of Camel Products, IGI Global, Hershey, USA, 2020, 480 p
- AlKurd R, Hanash N, Khalid N, Abdelrahim D N, Khan M A, Mahrous L, Radwan H, Naja F, Madkour M, Obaideen Abu Shihab K and Faris M. Effect of camel milk on glucose homeostasis in patients with diabetes: A systematic review and meta-analysis of randomised controlled trials. *Nutrients.* 2022; 14(6):1245, <https://doi.org/10.3390/nu14061245>
- Almathen F, Charruau P, Mwacharo J, Alnaqeeb M, Abdussamad A, Abdul R, Al-Ekna M, Faye B, Hanotte O, Burger P. Large scale geographic mitochondrial DNA analysis provide insights on the demographic and evolutionary history of the dromedary (*Camelus dromedarius*). *Proc. 34th Conf. Int. Soc. Anim. Genet. (ISAG, 2014).*, Xi'An (China), 2014.28/07-01/08/14, Abstract P4021, p. 116-117.
- Al-Numair N S, Theyab A, Alzahrani F, Shams A M, Al-Anazi IO, Oyouni AAA, Al-Amer OM, Mavromatis C, Saadeldin IM, Abdali WA and Hawsawi, YM. Camels' biological fluids contained nanobodies: promising avenue in cancer therapy. *Cancer Cell International.* 2022. 22(1), 279, <https://doi.org/10.1186/s12935-022-02696-7>
- Al-Shomrani BM, Manee MM, Alharbi SN, Altammami MA, Alshehri MA, Nassar MS, Bakhrebah MA, Al-Fageeh MB. Genomic Sequencing and Analysis of Eight Camel-Derived Middle East Respiratory Syndrome Coronavirus (MERS-CoV) Isolates in Saudi Arabia. *Viruses.* 2020; 3;12(6):611. doi: 10.3390/v12060611. PMID: 32503352; PMCID: PMC7354450.
- Allen W R. Proceedings of the First International Camel Conference: 2<sup>nd</sup>-6<sup>th</sup> February 1992 held in Dubai. R & W Publications, 1992 - Camels - 431 pages
- Ashraf A, Mudgil P, Palakkott A, Iratni R, Gan CY, Maqsood S and Ayoub MA. Molecular basis of the anti-diabetic properties of camel milk through profiling of its bioactive peptides on dipeptidyl peptidase IV (DPP-IV) and insulin receptor activity. *Journal of Dairy Science.* 2021; 104(1):61-77. <https://doi.org/10.3168/jds.2020-18627>
- Atigui M, Hammadi M, Barmat A, Farhat M, Khorchani T and Marnet PG. First description of milk flow traits in Tunisian dairy dromedary camels under an intensive farming system. *Journal of Dairy Research.* 2014; 81(2):173-182. <https://doi.org/10.1017/S0022029914000089>
- Ayadi M, Musaad A, Aljumaah R, Matar A and Faye B. Effects of manual udder stimulation on milk partitioning and flow traits during the machine milking in dairy camels. *Journal of Camel Practice and Research.* 2016; 23(1):1-5.
- Azhar EI, Velavan TP, Rungsung I, Traore T, Hui DS, McCloskey B, El-Kafrawy SA and Zumla A. Middle East respiratory syndrome coronavirus—a 10-year (2012-2022) global analysis of human and camel infections, genomic sequences, lineages and geographical origins. *International Journal of Infectious Diseases.* 2023; 131:87-94. <https://doi.org/10.1016/j.ijid.2023.03.046>
- Baba WN, Rasool N, Selvamuthukumara M and Maqsood S. A review on nutritional composition, health benefits and technological interventions for improving consumer acceptability of camel meat: an ethnic food of Middle East. *Journal of Ethnic Foods.* 2021; 8:1-13. <https://doi.org/10.1186/s42779-021-00089-1>
- Badawy AA, Othman RQA and El-Magd MA. Effect of combined therapy with camel milk-derived exosomes, tamoxifen and hesperidin on breast cancer. *Molecular & Cellular Toxicology.* 2021; 1-10. <https://doi.org/10.1007/s13273-021-00163-4>
- Bengoumi M, Riad F, Giry J, De La Farge F, Davicco MJ, Safwate A and Barlet JP. Hormonal control of water and sodium in plasma and urine of camels during dehydration and rehydration. *General and Comparative Endocrinology.* 1993; 89:378-386.
- Bengoumi M, Berrada J, Hidane K and Faye B. Diarrhées du chameau: écopathologie, physiopathologie et prévention. *Atelier Int. sur le lait de chamelle en Afrique.* (F. Lhoste, Ed), FAO-CIRAD-KARKARA, Niamey (Niger). 2003; 5-8/11/03
- Bonnet P. (Ed). Actes de l'atelier “Chameaux et dromadaires, Animaux laitiers” 24-26 Oct. 1994. Nouakchott (Mauritanie), Publ. CIRAD. 1998; (coll. Colloques), 300 pp.
- Breulmann M, Böer B, Wernery U, Wernery R, El Shaer HM, Alhadrami GA, Gallacher D, Peacock J, Ali Chaudhary S, Brown G and Norton J. The Camel from Tradition to Modern Times. UNESCO, Doha. 2007.
- Burger PA, Ciani E and Faye B. Old World camels in a modern world—a balancing act between conservation and genetic improvement. *Animal Genetics.* 2019; 50(6): 598-612.
- Cauvet G. Le chameau. Baillière et fils Publ., 1925; Paris, France
- Chaudhary ZI and Akbar SJ. The Camel and Its Diseases. A-Bayan Press Publ., 2000; Dubai, UAE
- Cross HE. The Camel and its Diseases. Bailliere, Tindall & Cox Publ., 1917; London, UK
- Curasson G. Le chameau et ses maladies. Vigot Frères Publ., 1947; Paris, France

- Dakkak A. (Ed.) Proc. Atelier régional sur « Maladies parasitaires et infectieuses du dromadaire », Actes Editions, 2000, IAV, Rabat (Morocco), 164 p.
- Droandi I. Il cammello: storia naturale, anatomica, fisiologica, zootecnica, patologia. 1936. Istituto Agricolo Coloniale Italiano, Firenze.
- El-Agamy ESI. Camel milk. In: Handbook of Milk of Non Bovine Mammals. 2017; Young W. *et al*, (Eds), Wiley Publ., 409-480, <https://doi.org/10.1002/9781119110316.ch6>
- El-Bahrawy KA, Khalifa MA and Rateb SA. Recent advances in dromedary camel reproduction: An Egyptian field experience. Emirates Journal of Food and Agriculture. 2015; 27(4):350-354, <https://doi.org/10.9755/ejfa.v27i4.19907>
- Emirates Industry for Camel Milk & Products “Camelicious”. [https://www.linkedin.com/company/camelicious/?trk=ppro\\_cprof&originalSubdomain=in](https://www.linkedin.com/company/camelicious/?trk=ppro_cprof&originalSubdomain=in)
- Emmanuel B. Comparative metabolic studies between ruminant species. In Annales de Recherches Vétérinaires. 1979; 10(2-3):480-481.
- Engelhardt W, Lechner-Doll M, Heller R, Rutagwenda Theogen and Schwartz Horst. Physiology of the forestomach in camelids with particular reference to adaptation to extreme dietary conditions: A comparative approach. Animal Research and Development. 1988; 28:56-70.
- Esam I Azhar 1, Thirumalaisamy P Velavan 2, Ikromi Rungsung 2, Tieble Traore 3, David S Hui 4, Brian McCloskey 5, Sherif A El-Kafrawy 6, Alimuiddin Zumla
- Etzion Z and Yagil R. Renal function in camels (*Camelus dromedarius*) following rapid rehydration. Physiological Zoology. 1986; 59(5):558-562.
- Farah Z. Composition and characteristics of camel milk. Journal of Dairy Research. 1993; 60(4):603-626. <https://doi.org/10.1017/S0022029900027953>
- Farah Z, Fisher A. (Eds). Milk and meat from the camel: Handbook on products and processing. VDF Hochschulverlag and ETH Zürich Publ., 2004; Switzerland, 229 pp.
- Faye B. Guide de l'élevage du dromadaire. Ed. Sanofi, Libourne, France, 1997, 126 pp.
- Faye B, Bonnet P, Charbonnier G and Marti A. Bilan des activités de recherche sur le dromadaire par analyse bibliométrique de la littérature scientifique. Cas particuliers des travaux sur le chamelon. Atelier International sur le chamelon : “le Chamelon, futur de l'élevage camélin”. Ouarzazate, 24-26 Oct. 1999, Maroc. Revue d'Elevage et de Médecine Vétérinaire des Pays Tropicaux. 2000; 53:125-131.
- Faye B and Bengoumi M. Camel Clinical Biochemistry and Haematology. 1<sup>st</sup> Edn. 2018; Springer. ISBN : 978-3-319-95560-5
- Faye B and Esenov P. (Eds), Desertification combat and food safety: the added value of camel producers”. Vol. 362 NATO Sciences Series, Life and Behavioural Sciences. IOS press Publ., Amsterdam (The Netherlands). 2005; 240 pp.
- Faye B and Brey F. Les relations entre chameaux et société: entre marginalisation et idéalisation. Revue Ethnozootechnie. 2005; n°77 -Varia, 43-50.
- Faye B, Konuspayeva G, Magnan C. L'élevage des grands camélidés. Ed. QUAE, Versailles, coll. Guide pratique. 2022; pp 204. ISBN 978-2-7592-3499-8 , <https://doi.org/10.35690/978-2-7592-3500-1>
- Faye B, Konuspayeva G and Magnan C. Large camel farming- A care-management guide from breeding to camel products. Springer Publ. Dordrecht, The Netherlands. 2023; pp 217. <https://doi.org/10.1007/978-94-024-2237-5>
- Faye B and Konuspayeva G. Camel milk composition and future market potential. CABI Reviews. 2024; 19(1), <https://doi.org/10.1079/cabireviews.2024.0021>
- Finbert EJ. La vie du chameau- Le vaisseau du désert. Albin Michel Publ., 1938; Paris, France
- Gahlot TK and Chouhan DS. Camel Surgery. 1992. Gyan Prakashan Mandir, Bikaner, India.
- Gahlot TK (Ed.). Journal of Camel Practice and Research. 1994. Vol 1, No 1. Camel Publishing House. ISSN 0971-6777 (Print); 2277-8934 (Online).
- Gahlot TK (Ed.). Selected Topics on Camelids. Edn. 2000, ISBN 81-901141-0-7. The Camelid Publishers, Bikaner 334001, INDIA.
- Gahlot Tarun Kumar and Adams C. Camel milk and other dietary treatments in autism: an overview. Journal of Camel Practice and Research. 2023; 30(2):143-147.
- Gahlot TK, Wernery U, Tibary Ahmed and Zhao XX. (Eds). Selected Bibliography on Camelids 1991-2000. Edn 2002, ISBN 81-901141-1-5. The Camelid Publishers, Bikaner 334001, INDIA.
- Gahlot TK (Ed). Selected Research on Camelid Physiology and Nutrition. 2004, ISBN 81-901141-2-3. The Camelid Publishers, Bikaner 334001. INDIA.
- Gahlot TK (Ed). Proceedings of the International Camel Conference “Recent trends in Camelids research and Future strategies for saving Camels”, College of Veterinary and Animal Sciences, Bikaner, Rajasthan, India, 16-17 February 2007.
- Gahlot TK (Ed). Proceedings of the International Camel Conference on “Sustainability of camel population and production” held at King Faisal University, Al-Hasa, Kingdom of Saudi Arabia, 17-20 February, 2013.
- Gahlot TK and Chhabra MB (Eds). Selected Research on Camelid Parasitology, Edn. 2009, ISBN 81-903140-0-9. Camelid Publishing House, Bikaner 334001. INDIA.
- Gahlot TK, Wernery U. and Mulyrdermans S. (Eds). Selected Research On Camelid Immunology. Edn. 2016. ISBN-81-903140-4-1, First Edition, Camel Publishing House, Bikaner, India.
- Gahlot TK, Saber AS, Nagpal SK and Wang Jianling (Eds). Selected Research on Gross Anatomy and Histology of Camels. 2011. ISBN 81-903140-1-7. Camelid Publishing House, 67 Gandhi Nagar West, Near Lalgargh Palace, Bikaner 334001 INDIA.
- Gaouar SBS and Ciani E. An Overview of Camel Biodiversity and Genetics. Biology and Life Sciences Forum. 2023; 22(1):15. <https://doi.org/10.3390/blsf2023022015>



- Gauthier-Pilters H and Dagg AI. The camel. Its evolution, ecology, behavior and relationship to man. Univ. Chicago Press Publ., 1981; Chicago, USA
- Gupta BM, Ahmed KM, Gupta R and Tiwari R. World camel research: a scientometric assessment, 2003–2012. *Scientometrics*, 2015; 102(1), 957-975.
- Hamers-Casterman C, Atarhouch T, Myldermans S, Robinson G, Hamers C, Songa EB, Bendahman N and Hamers R. Naturally occurring antibodies devoid of light chains. *Nature*. 1993; 363(6428):446-448.
- Heuse G. L'éremologie, science du XXI<sup>e</sup> Siècle. *Sècheresse*. 1992; 3(1):5-10
- Higgins AJ. The Camel in Health And Disease. 1986. Balliere Tindall, London. pp 168.
- Higgins AJ and Kock R. The camel in health and disease. *British Veterinary Journal*. 1984; 140(5):482-504.
- Hoste C, Peyre de Fabregues B and Richard D. Le dromadaire et son élevage. Etudes et synthèses de l'IEMVT n°12, D. Richard (Ed.), IEMVT Publ., 1984. Maisons-Alfort, France
- Hussen J and Schubert HJ. Recent advances in camel immunology. *Frontiers in Immunology*. 2021; 11:614150.
- Indra P, Maratch A and Batsouour L. [le chameaumontol]. 2003. Oulanbator, Mongolia
- Jirimutu, Wang Z, Ding G, Chen G, Sun Y, Sun Z, *et al* Genome sequences of wild and domestic Bactrian camels. *Nature Communications*. 2012; 3:1202. doi: 10.1038/ncomms2192. Erratum in: *Nat Commun*. 2013; 4. doi: 10.1038/ncomms3089. PMID: 23149746; PMCID: PMC3514880.
- Johnson EH, Maghoub O, Eljack AH, Kadim I, Bobade PA, Tageldin MH, Al-Marzooqi WS, El Tahir Ahmed Y. (Eds). Proceeding of 3<sup>rd</sup> ISOCARD conference, 29th January -1<sup>st</sup> February, 2012; Muscat (Sultanate of Oman), 431 p.
- Kadim IT, Mahgoub O, Purchas RW. A review of the growth and of the carcass and meat quality characteristics of the one-humped camel (*Camelus dromedarius*). *Meat Science*. 2008; 80:555-569, <https://doi.org/10.1016/j.meatsci.2008.02.010>
- Kadim I, Mahgoub O, Faye B and Farouk M. (Eds). *Camel Meat and Meat Products*. CAB International Publ, Oxfordshire, UK & Boston, USA. 2013; 248 p.
- Kadim IT, Al-Amri IS, Alkindi AY and HaqQM. Nutritional values and health benefits of dromedary camel meat. *Animal Frontiers*. 2022; 12(4):61-70.
- Kaindi D and Njage P. Microbial aspect of lactic acid bacteria isolated from camel milk. In: *Handbook of Research on Health and Environmental Benefits of Camel Products*. 2020; pp 54-74. <https://doi.org/10.4018/978-1-7798-1604-1.ch004>
- Kandeel M, Morsy MA, Abd El-Lateef HM, Marzok M, El-Beltagi HS, Al Khodair KM, Soliman WE, Albokhadaim I and Venugopala KN. A century of "camel research": a bibliometric analysis. *Frontiers in Veterinary Science*. 2023; 10:1157667.
- Kappeler SR, Rahbek-Nielsen H, Farah Z, Puhon Z, Hansen EB and Johansen E. Characterisation of recombinant camel chymosin reveals superior properties for the coagulation of bovine and camel milk. *Biochemical and Biophysical Research Communications*. 2006; 342(2):647-654. <https://doi.org/10.1016/j.bbrc.2006.02.014>
- Kaskous S. A quarter individual milking machine "Stimu Lactor" in a camel farm in Switzerland: According to field Study. *Journal of Camel Practice and Research*. 2023; 30(3):267-271.
- Kataria AK, Kataria N, Swarnkar CP and Sharma KN. Serum immunoglobulin status in camel (*Camelus dromedarius*). *International Journal of Animal Science*. 1994; 9:299-300.
- Kataria AK, Kataria N, Singh R, Garg SL and Agarwal VK. Serum immunoglobulin levels in camel (*Camelus dromedarius*) during dehydration and rehydration in winter season. *Journal of Immunology and Immunopathology*. 1999; 1:58-59.
- Kataria AK and Sharma KN. Dromedary immunoglobulins: Physico-chemical and biological characterisation. *Journal of Camel Practice and Research*. 1999; 6(2):279-286.
- Kataria AK and Sharma KN. Chromatographic purification of serum and colostral immunoglobulins of camel (*Camelus dromedarius*). *Journal of Camel Practice and Research*. 2000; 7(1):91-95.
- Kataria AK. Some characteristics of camel (*Camelus dromedarius*) immunoglobulins. *Journal of Immunology and Immunopathology*. 2001; 3(2):126.
- Kataria AK, Kataria N and Sharma KN. Serum protein and immunoglobulin profile in camel (*Camelus dromedarius*). *Indian Journal of Animal Health*. 2002; 41(1):5-8.
- Kataria AK and Sharma KN. Isolation and characterisation of Fab and Fc fragments from camel colostral IgG1. *Journal of Immunology and Immunopathology*. 2003; 5(1):64-66.
- Kataria AK and Kataria N. Immunoradiometric assay of serum IgE levels in dromedary camel. *Journal of Camel Practice and Research*. 2004; 11(1):11-13.
- Kataria N, Kataria AK, Agarwal VK, Garg SL and Sahani MS. Effect of long term dehydration on serum constituents in extreme climatic conditions in camel (*Camelus dromedarius*). *Indian Journal of Physiology and Pharmacology*. 2002a; 46(2):218-222.
- Kataria N, Kataria AK, Agarwal VK, Garg SL and Sahani MS. Thiosulphate clearance test for determination of glomerular filtration rate in dromedaries. *Journal of Camel Practice and Research*. 2002b; 9(1):9-15
- Kataria N, Kataria AK, Agarwal VK, Garg SL and Sahani MS. Intravenous glucose tolerance test (IVGTT) in camel during dehydration and rehydration in winter and summer. *Journal of Camel Practice and Research*. 2002c; 9(1):17-25.
- Kataria N, Kataria A K, Agarwal VK, Garg SL and Sahani MS. Osmolal and water clearances in camels during hot and cold ambience. *Journal of Camel Practice and Research*. 2002d; 9(2):87-90.
- Kataria N, Kataria AK, Agarwal VK, Garg SL and Sahani MS. Effect of hot and cold ambience on renal clearances of



- electrolytes in camels. *Journal of Camel Practice and Research*. 2002e; 9(2):91-96.
- Kataria N, Kataria AK Agarwal VK, Garg SL and Sahani MS. Effect of seasonal dehydration on creatinine clearance in Indian dromedary camels. *Journal of Camel Practice and Research*. 2003; 10(2):91-97.
- Kataria N, Kataria AK Agarwal VK, Garg SL and Sahani MS. Solute loads and transfer function of kidney in dromedary camel during dehydration and rehydration in winter and summer. *Veterinarski Arhiv*. 2007; 77(3):237-246.
- Khalafalla AI. Zoonotic diseases transmitted from the camels. *Frontiers in Veterinary Science*. 2023; 10:1244833
- Kohler-Rollefson Ilse. *Hoofprints on the Land*. 2023. ISBN: 9781645021520. Chelsea Green Publishing Co. Genre: Nature. 288 pages.
- Kohler Rollefson I, Mundy P and Mathias E. *A Field Manual of Camel diseases. A traditional health care for the dromedary*. Intermediate Technology Publ., LPP, Germany. 2001, 254 pp
- Kohler Rollefson I. *Camel Karma*. 2023. ISBN 9789354476464, 425 pages. Speaking Tiger Books LLP, 125A, Ground Floor, Shahpur Jat, near Asiad Village, New Delhi – 110049.
- Köhler-Rollefson Ilse. Camel biodiversity – and how to conserve it, *Animal Frontiers*. 2022; 12(4):17-19, <https://doi.org/10.1093/af/vfac042>
- Konuspayeva G. (Ed). *Proc. of 4th Conference of ISOCARD, "Silk Road Camel: the camelids, main stakes for sustainable development"*, June 8-12, 2015 Almaty, Kazakhstan, Special Issue of Scientific and Practical Journal *Veterinariya* # 2 (42) 2015, 483 pp.
- Konuspayeva G, Faye B and Loiseau G. The composition of camel milk : A meta-analysis of the literature data. *Journal of Food Composition and Analysis*, 2009; 22:95-101. <https://doi.org/10.1016/j.jfca.2008.09.008>
- Konuspayeva G, Camier B, Aleilawi N, Al-Shumeimyri M, Al-Hammad K, Algruin K, Alshammari F, Beaucher E and Faye B. Manufacture of dry- and brine-salted soft camel cheeses for the camel dairy industry. *International Journal of Dairy Technology*. 2017; 70(1):92-101, <https://doi.org/10.1111/1471-0307.12319>
- Konuspayeva G and Faye B. Recent advances in camel milk processing. *Animals*. 2021; 11(4):1045.
- Lal Manohar, Mahala Neelam, Mittal Aastha, Ramdas Anisha, Shrivastava Chetan and Dubey Uma S. Studies on cytotoxic activity of camel milk whey protein as a nutraceutical against hela cells. *Journal of Camel Practice and Research*. 2023; 30(2):155-164. DOI : 10.5958/2277-8934.2023.00021.8
- Leese AS. *A Treatise on the One-Humped Camel in Health and in Disease*. Haynes & Son Publ., 1927; London, UK
- Lemrhamed A, Tabite R, Fahr M, Riad F, El-Abbadi N, Tahri E., Faye B and El-Khasmi M. Evaluation of preslaughter stress responses during waiting time at lairage in dromedary camels (*Camelus dromedarius*). *Journal of Camel Practice and Research*. 2019; 26(2):149-156, <https://doi.org/10.5958/2277-8934.2019.00023.7>
- Lhoste F. (Ed.) *Comptes-Rendus de l'atelier « lait de chamelle pour l'Afrique »*, 5-8/11/2003, Niamey (Niger), FAO publ., 203 pp.
- Menchetti L, Faye B and Padalino B. New animal-based measures to assess welfare in dromedary camels. *Tropical Animal Health and Production*. 2021; 53:533, <https://doi.org/10.1007/S11250-021-02978-8>
- Ming L, Yuan L, Yi L. *et al*. Whole-genome sequencing of 128 camels across Asia reveals origin and migration of domestic Bactrian camels. *Commun Biol* 3, 1 (2020). <https://doi.org/10.1038/s42003-019-0734-6>
- Moussaiev Z, Torejanov A and Seidalyev B. [Elevage des chameaux]. *Basmaou Publ.*, 2007; Almaty, Kazakhstan
- Moussahil A, Tabite R, Lemrhamed A, Farh M, Iddar A and El Khasmi M. Impact of the duration of stress steps before slaughter and at slaughter on the physicochemical and biochemical characteristics of meat in camels. *Asian Journal of Research in Animal and Veterinary Sciences*. 2022; 10(3):1-12. <http://info.submit4journal.com/id/eprint/20>
- Muyldermans Sand Lauwereys M. Unique single-domain antigen binding fragments derived from naturally occurring camel heavy-chain antibodies. *Journal of Molecular Recognition*. 1999; 12(2):131-140. [https://doi.org/10.1002/\(SICI\)1099-1352\(199903/04\)12:2<131::AID-JMR454>3.0.CO;2-M](https://doi.org/10.1002/(SICI)1099-1352(199903/04)12:2<131::AID-JMR454>3.0.CO;2-M)
- Nagy P, Faye B, Marko O, Thomas S, Wernery U and Juhasz J. Microbiological quality and somatic cell count in bulk milk of dromedary camels (*Camelus dromedarius*): descriptive statistics, correlations and factors of variation. *Journal of Dairy Science*. 2013a; 96:5625–5640, <https://doi.org/10.3168/jds.2013-6990>
- Nagy P, Skidmore JA and Juhasz J. Use of assisted reproduction for the improvement of milk production in dairy camels (*Camelus dromedarius*). *Animal Reproduction Science*. 2013b; 136(3):205-210. <https://doi.org/10.1016/j.anireprosci.2012.10.011>
- Nagy P, Faigl V, Reiczigel J and Juhasz J. Effect of pregnancy and embryonic mortality on milk production in dromedary camels (*Camelus dromedarius*). *Journal of Dairy Science*. 2015; 98(2):975-986. <https://doi.org/10.3168/jds.2014-8546>
- Nagy P and Juhasz J. Review of present knowledge on machine milking and intensive milk production in dromedary camels and future challenges. *Tropical Animal Health and Production*. 2016; 48(5):915-926. <https://doi.org/10.1007/s11250-016-1036-3>
- Nagy PP, Skidmore JA and Juhasz J. Intensification of camel farming and milk production with special emphasis on animal health, welfare and the biotechnology of reproduction. *Animal Frontiers*. 2022; 12(4):35-45.
- Ngere I, Munyua P, Harcourt J, *et al* High MERS-CoV seropositivity associated with camel herd profile, husbandry practices and household socio-demographic characteristics in Northern Kenya. *Epidemiology and Infection*. 2020;148:e292. <https://doi.org/10.1017/S0950268820002939>
- Niasari-Naslaji A, Nikjou D, Skidmore JA, Moghiseh A, Mostafaey M, Razavi K, Moosavi-Movahedi AA.

- Interspecies embryo transfer in camelids: the birth of the first Bactrian camel calves (*Camelus bactrianus*) from dromedary camels (*Camelus dromedarius*). *Reproduction, Fertility and Development*. 2009; 21(2):333-7. doi: 10.1071/rd08140. PMID: 19210924.
- Olsson PO, Tinson AH, Al Shamsi N, Kuhad KS, Singh R, Son Y-B, Jeong Y, Jeong YW, Cai L, Sakaguchi K, Kim S, Choi EJ, Yu X and Hwang WS. Blastocyst formation, embryo transfer and breed comparison in the first reported large scale cloning of camels. *Scientific Reports*. 2021; 11:14288.
- Padalino B and Faye B. (Eds.), *Dromedary Camel Behavior and Welfare: Camel Friendly Management Practices*. Springer Publ., 2024; Cham, Switzerland
- Pereira CH. *Merveilles des dromadaires* ; Publ. Arts des 2 mondes, Aulnay/bois, France, 2024; 80 pp.
- Proceedings of the 3<sup>rd</sup> Conference of the International Society of Camelid Research and Development, Organised by Sultan Qaboos University, College of Agricultural and Marine Sciences, Department of Animal and Veterinary Sciences, held in Muscat, Sultanate of Oman from the 29<sup>th</sup> January to 1<sup>st</sup> February 2012.
- Purohit GN, Vyas S, Yadav V, Nain S, Chaudhary AK, Kumar A, Dholpuria S and Saraswat CS. Semen characteristics and artificial insemination in dromedary camels. *Small Ruminant Research*. 2023; 220:106911. <https://doi.org/10.1016/j.smallrumres.2023.106911>
- Ramadan RO. *Surgery and Radiology of the Dromedary Camel*. Al-Jawad Printing Press, 1994. ISBN 9960270246, 9789960270241. 360 pages.
- Rathore GS. *Camels and Their Management*. ICAR Publ., 1986; New Delhi, India
- Rathinasabapathy G and Rajendran L. Mapping of world-wide camel research publications: a scientometric analysis. *Journal of Library, Information and Communication Technology*. 2013; 5(1-2):35-40.
- Raiymbek G, Kadim I, Konuspayeva G, Mahgoub O, Serikbayeva A and Faye B. Discriminant amino-acid components of Bactrian (*Camelus bactrianus*) and Dromedary (*Camelus dromedarius*) meat. *Journal of Food Composition and Analysis*. 2015; 41:194-200, <https://doi.org/10.1016/j.jfca.2015.02.006>
- Raiymbek G, Kadim I, Al-Amri Issa S, Alkindi Abdulaziz Y, Faye B, Khalf SK, Kenenbay SI, Purchas RW. Concentrations of nutrients in six muscles of Bactrian (*Camelus bactrianus*) camels. *Journal of Camel Practice and Research*. 2018; 25(1):109-121.
- Riad F, Bengoumi M, Davicco MJ, Safwate A, Barlet JP. Influence of 1- $\alpha$ -hydroxycholecalciferol on calcium and phosphorus concentration in camel milk. *Journal of Dairy Research*. 1994; 61:567-571.
- Rollefson Ilse-Kohler. Proceedings of an International Conference on Saving the camel and people's livelihoods: Building a multi-stakeholder platform for the conservation of the camel in Rajasthan, held on 23-25 November 2004. Lokhit Pashu-Palak Sansthan, Sadri, Rajasthan, India.
- Rosenheck D, Ravee Y and Yagil R. Camel milk as an alternative treatment for Crohn's disease: 1264. *Official Journal of the American College of Gastroenterology*. 2012; ACG 107, S503.
- Saint-Martin G. Actes de l'atelier « Peut-on améliorer les performances de reproduction des camelins ?, Paris, 10-12/09/1990, Publ. CIRAD, 1993, Etudes et synthèses de l'ITEMVT, 438 pp.
- Salamt N, Idrus RBH, Kashim MIAM and Mokhtar MH. Anticancer, antiplatelet, gastroprotective and hepatoprotective effects of camel urine: A scoping review. *Saudi Pharmaceutical Journal*. 2021 Jul; 29(7):740-750. doi: 10.1016/j.jsps.2021.05.006. Epub 2021 Jun 6. PMID: 34400869; PMCID: PMC8347850.
- Saltin B and Rose RJ. The racing camel: physiology, metabolic functions and adaptation. *Scandinavian Physiological Society Publ.*, 1994; Stockholm, Sweden
- Schmidt-Nielsen B, Schmidt-Nielsen K, Houtp TT and Jarnum SA. Water balance of the camel. *American Journal of Physiology-Legacy Content*. 1956; 185(1):185-194.
- Schmidt-Nielsen B, Schmidt-Nielsen K, Houtp TR and Jarnum SA. Urea excretion in the camel. *American Journal of Physiology-Legacy Content*. 1957; 188(3):477-484.
- Schwartz H and Dioli M. *The One-Humped Camel in Eastern-Africa*. Publ. Verlag, 1992; Weikersheim, Germany
- Sghiri A and Kichou F. (Eds). Proceedings of the 5<sup>th</sup> Conference ISOCARD "Recent Advances in Camelids Biology, Health and Production", 12-15 Nov. 2018, Laâyoune, Morocco, IAV Hassan II, 2018, 584 pp.
- Siddiqui MI and Telfah Mohammed. *A Guide Book of Camel Surgery*. 2010. Publisher: Abu Dhabi Food Control Authority ADFCA. ISBN: 978-9948-03-773-6.
- Skidmore JA, Juhasz J and Nagy P. Welfare aspects of reproductive care and management of dromedary camels. In: *Dromedary Camel Behavior and Welfare: Camel Friendly Management Practices* (pp 189-206). Springer Publ., 2024; Cham, Switzerland.
- Tanwar PD, Ghorui SK, Kochar SK, Singh Raghvendar and Patil NV. Production and preclinical assessment of camelid immunoglobulins against Echis sochureki venom from desert of Rajasthan, India. *Toxicon*. 2017; 134:1-5. ISSN 0041-0101. <https://doi.org/10.1016/j.toxicon.2017.05.012>.
- Terentyev CM. *Verbliudovodstvo [Camel breeding]*, Kolos Publ., 1975; Moscow, USSR.
- Tibary A and Anouassi A. *Therigenology in Camelidae*. Actes Editions, 1997; Rabat, Maroc
- Vallon A. *Histoire naturelle du dromadaire*. Recueil de mémoires et observations sur l'hygiène et la médecine vétérinaire militaire. Dumaine J. Publ., Military library, 1856 ; Paris, France.
- Wani NA, Wernery U, Hassan FAH, Wernery R and Skidmore JA. Production of the First Cloned Camel by Somatic Cell Nuclear Transfer. *Biology of Reproduction*. 2010; 82:373-379. <https://doi.org/10.1095/biolreprod.109.081083>
- Wani, NA. *In vitro* embryo production (IVEP) in camelids: Present status and future perspectives. *Reproductive Biology*. 2021; 21(1):100471. <https://doi.org/10.1016/j.repbio.2020.100471>.

- Wani NA, Vettical BS and Hong SB. First cloned Bactrian camel (*Camelus bactrians*) calf produced by interspecies somatic cell nuclear transfer: A step towards preserving the critically endangered wild Bactrian camels. PLOS ONE. 2017; 12(5):e0177800. <https://doi.org/10.1371/journal.pone.0177800>
- Wardeh MF. A programme for development of camel production in the Arab countries. ACSAD/AS/69/1986, Working paper, 1986; Damascus, Syria
- Weiss Walter M and Wernery Ulrich. 111 Dinge über Kamele die man wissen muss. 2021; Emons Verlag, Germany, 240 Seiten, ISBN 978-3-7408-1206-5.
- Wernery U and Kaaden OR. Infectious Diseases of Camelids. Blackwell Wissenschafts-Verlag Publ., 1995; Berlin, Germany
- Wernery U, Kinne J and Schuster RK. Camelid Infectious Disorders. World Organisation for Animal Health (OIE) Publ., 2014; Paris, France
- Wilson RT. The Camel. Longman Publ., 1984; London, UK
- Yagil R. The Desert Camel. Comparative Physiological Adaptation. Karger publ., 1985; London, UK
- Zhao-Xin-Xu. Ecophysiology and Reproduction of the Camelidae, Lanzhou Univ. Publ., 1996; Lanzhou, China



# THE TRENDS OF CAMEL RESEARCH IN NORTH AMERICA: A BIBLIOMETRIC APPROACH

Ahmed Alsaleem<sup>1</sup> and Mahmoud Kandeel<sup>1,2</sup>

<sup>1</sup>Department of Biomedical Sciences, College of Veterinary Medicine, King Faisal University, 31982 Al-Ahsa, Saudi Arabia

<sup>2</sup>Department of Pharmacology, Faculty of Veterinary Medicine, Kafrelsheikh University, 33516 Kafrelsheikh, Egypt

## ABSTRACT

Camels are witnessing global attraction even in countries outside the natural camel habitat. This study aims to provide a comprehensive bibliometric analysis of camel research in North America, identifying key trends, influential contributors, and collaborative networks. Data was sourced from the Scopus database, yielding 786 papers on camel research affiliated with institutions in the United States and Canada. Bibliometric tools such as VOSviewer and R Studio's Bibliometrix package were used to visualise and analyse the data. The bibliometric analysis of camel research in North America highlights substantial collaborative efforts between the North American countries and international partners. The data spans 493 sources, with an annual growth rate of 1.89%. The research involves 3127 authors and a notable international co-authorship rate of 52.04%. Camel research in North America spans multiple disciplines, including Agricultural and Biological Sciences, Medicine and Veterinary Sciences, Chemistry, Genetics and Molecular Biology, Immunology and Microbiology, Environmental Science, Engineering, and Social Sciences. This interdisciplinary approach underscores the broad impact and significance of camel studies. The international partnerships have facilitated significant advancements in understanding zoonotic diseases, genetic diversity, and the nutritional benefits of camel milk, underscoring the importance of maintaining robust international research networks to enhance the quality and impact of camel studies. This bibliometric analysis highlights the growing significance of camel research in North America, with strong international collaborations driving advancements. However, gaps remain in integrating advanced technologies and exploring socio-economic impacts in non-traditional regions. Continued investment and international collaboration are essential to address these gaps and drive innovative research initiatives.

**Key words:** Bibliometric analysis, camel research, interdisciplinary collaboration, North America, publication trends

Camel research holds substantial importance due to the unique characteristics and vital roles camels play in arid and semi-arid regions (Adah *et al*, 2023). Camels are renowned for their exceptional adaptability to harsh environments, making them invaluable for sustaining livelihoods in regions where other livestock might not thrive (Sahoo, 2020). They provide essential resources such as milk, meat, and transportation, and their resilience against extreme temperatures and water scarcity positions them as key assets in the face of climate change (George, 2024). Moreover, camels have unique physiological and genetic traits that offer significant insights into medical and biological sciences, contributing to advancements in understanding disease resistance, metabolic processes, and potential biomedical applications (Hedayat-Evrigh *et al*, 2020; Ming *et al*, 2020).

Recently, we provided a bibliometric analysis of camel research, which showed increasing global attention (Kandeel *et al*, 2023). This surge in interest

can be attributed to a growing recognition of camels' ecological and economic significance, as well as their potential contributions to global health and scientific knowledge. Researchers in the United States and Canada have been particularly active in studying various aspects of camel biology, health, and management (Camel sciences and economy in the world: current situation and perspectives). The collaborative efforts between North American institutions and international partners have led to significant advancements in understanding camels' role in zoonotic diseases, genetic diversity, and the nutritional benefits of camel milk.

The primary objectives of this study are to provide a comprehensive bibliometric analysis of camel research in North America, identifying key trends, influential contributors, and collaborative networks. By analysing a robust dataset sourced from the Scopus database, this study aims to map the evolution of camel research over time, highlighting

SEND REPRINT REQUEST TO MAHMOUD KANDEEL [email: mkandeel@kfu.edu.sa](mailto:mkandeel@kfu.edu.sa)

significant periods of growth and identifying the most productive researchers and institutions. Additionally, this study seeks to elucidate the interdisciplinary nature of camel research, exploring how collaborations between North American and international researchers have contributed to the advancement of this field. Furthermore, this study intends to show the global relevance of camel research and its implications for various scientific and practical domains. By identifying gaps in the existing literature and highlighting emerging trends, this study aims to foster stronger international collaborations and drive innovative research initiatives.

## Materials and Methods

### *Data Source and search strategy*

The data was sourced from the Scopus database using a targeted search query: TITLE (camel) AND (LIMIT-TO (AFFILCOUNTRY , "United States") OR LIMIT-TO (AFFILCOUNTRY , "Canada")). This search yielded a total of 786 papers. The data extraction process involved gathering essential bibliographic details for each paper, including the title, authors, abstract, keywords, publication year, source (journal or conference), document type, and subject area.

### *Inclusion and exclusion criteria*

The inclusion criteria for the bibliometric analysis on camel research in North America were established to ensure the relevance and quality of the data collected. Studies were included if they were indexed in the Scopus database, focused on camel research, and affiliated with institutions in the United States or Canada. The search was conducted using a specific query that targeted articles with the term "camel" in the title and limited the results to those affiliated with North American countries. Additionally, only peer-reviewed articles, reviews, conference papers, letters, book chapters, notes, and short surveys were considered to provide a comprehensive overview of the scholarly contributions in this field. Exclusion criteria were applied to maintain the dataset's integrity and relevance. Duplicate records were checked to avoid redundancy, and erratum records were excluded to focus on original research contributions.

### *Data Cleaning and Preprocessing*

The data from Scopus database were exported in CSV file. The data were cleaned and pre-processed to remove duplicates and irrelevant records.

## *Bibliometric analysis*

The bibliometric analyses were conducted using two primary tools: VOSviewer and R Studio with the Bibliometrix package (Aria and Cuccurullo 2017; Arruda *et al*, 2022; Wong 2018). VOSviewer was employed for constructing and visualising bibliometric networks, such as co-authorship, and keywords networks.

### *Publication Trends Analysis*

Trends over time were examined to comprehend the expansion of research in this area, focusing on annual publication numbers, the most productive authors, and significant papers. Descriptive statistics were computed to give an overview of the dataset, covering the distribution of papers by year, document type, and subject matter.

## Results and Discussion

### *Dataset characteristics*

The dataset comprises 786 articles, each detailing various aspects of scholarly articles indexed in Scopus. The data includes 1) the authors information, such as their names, full names, and unique ID. Additionally, the dataset captures the titles of the articles, providing a quick reference to the subject matter of each entry. 2) Bibliographic details of the articles, including the year of publication, the source title (journal or conference name), volume, and issue numbers.

The provided PRISMA flow diagram presents a systematic approach to the bibliometric analysis of camel research trends in North America (Fig 1). The screening process, dated June 10th, 2024, identified a total of 786 records, 784 of which were deemed suitable for inclusion in the bibliometric analysis after confirming there were no duplicates, and 2 erratum records were removed.

### *Document Types*

The bar chart illustrates the distribution of various document types included in the bibliometric analysis of camel research in North America (Fig 2). The data shows that out of the 784 total records, the majority are research articles (634), followed by review papers (41), conference papers (36), letters (22), book chapters (20), notes (17), editorials (11), and short surveys (3). The x-axis of the bar chart represents the number of documents, while the y-axis lists the different types of articles. This representation underscores the predominance of primary research articles in the field and highlights




Fig 1. PRISMA flowchart for the bibliometric analysis on trends of camel research in North America.




Fig 2. Distribution of document types in camel research in North America.



the diverse mediums through which camel research is communicated.

Publication Trend

The trends in camel research publications in North America can be delineated into three distinct phases based on the graph provided (Fig 3). The first phase, spanning from 1859 to the mid-20th century, is characterised by a sparse and inconsistent publication output. During this period, the number of publications remained very low, often with multiple years passing without any new articles. This phase reflects the nascent stage of camel research in North America, where interest and resources dedicated to this field were minimal. The sporadic publications during this time may have been driven by isolated scientific inquiries or specific historical events that briefly captured academic interest.

The second phase, from the mid-20th century to 2008, marks a period of gradual growth and increased consistency in camel research publications. Starting around the 1960s, there is a noticeable, albeit slow, rise in the number of publications. This phase is indicative of a growing but still limited interest in camel research, possibly fuelled by expanding academic institutions and a broader recognition of the ecological and economic importance of camels. During this period, the research output began to stabilize, with a small but steady stream of publications each year, setting the stage for more significant developments in the subsequent phase.

The third phase, from 2009 to 2024, is characterised by a sharp and sustained increase in the number of camel research publications. The data showed a clear upward trend, with the number of publications accelerating rapidly, peaking in recent




Fig 3. The Trend of publications in camel research in North America.




Fig 4. Geographic distribution of camel research publications in North America and in collaboration with other countries. The intensity of the colour shading indicates the number of publications, ranging from 1 to 689.

years, particularly in 2021-2023. This explosive growth can be attributed to that camel studies have become a well-established and actively pursued area of scientific inquiry in North America.

International collaboration

The geographic distribution of camel research publications from North America, with a focus on collaborative work with other countries, reveals significant trends in academic contributions (Fig 4). The United States leads by a substantial margin, producing 689 publications. Canada, with 112 publications, also shows a strong research presence.

The collaborative nature of camel research is evident from the involvement of several countries. Egypt (94 publications) and Saudi Arabia (83

publications) are notable partners. These countries' high publication numbers highlight the importance of collaborative research efforts. The United Kingdom (51 publications) and the United Arab Emirates (46 publications) also have significant academic output, indicating strong international partnerships with North American researchers.

Other countries such as China (40 publications), France (29 publications), Germany (24 publications), and Jordan (23 publications) contributed to the global research landscape through collaborations. These numbers reflect diverse research interests and the importance of international collaborations in advancing camel research. The involvement of countries like Kenya (20 publications) and Ethiopia (18 publications) in Africa underscores the collaborative efforts to address region-specific challenges and leverage local expertise. The widespread distribution of collaborative publications across various countries demonstrates a global recognition of the importance of camel research, with contributions spanning from North America to Europe, Asia, and Africa.

### *Active institutions*

The bar chart in Fig 5 shows the number of camel research publications produced by various institutions in North America and their collaborations with other countries. The Fig was set to display affiliations with 10 or more publications. The x-axis represents the number of publications, while the y-axis lists the institutions involved. The data highlights the significant contributions of multiple institutions from Asia and Africa implying the predominance of collaboration research.

King Saud University stands out with the highest number of publications, indicating its leading role in camel research, likely due to its geographical and cultural proximity to camel populations. This prominence is followed by Columbia University, and the Centers for Disease Control and Prevention (CDC). These institutions are known for their strong research programs and resources, which likely facilitate extensive studies and collaborations.

Other notable institutions include Washington State University Pullman, the University of South Florida, and the University of Florida, all of which have made significant contributions to the field. The presence of international institutions like Cairo University, Suez Canal University, and the United Arab Emirates University highlights the collaborative nature of camel research, emphasising the global

effort to advance knowledge in this area. The diverse range of institutions, including veterinary schools, universities, research centers, and specialized institutes, underscores the interdisciplinary and collaborative approach required to address the complexities of camel research.

### *The most active funding agents*


The bar chart in Fig 6 illustrates the contributions of top funding agencies to camel research projects, highlighting those with five or more funded projects. The x-axis represents the number of funded projects, while the y-axis lists the funding agencies involved.

The National Institutes of Health (NIH) stands out as the most prominent funding agency, supporting the highest number of projects, with a total of 37. This highlights the NIH's commitment to diverse research areas, including camel studies, which may intersect with broader health and biomedical research interests. The National Science Foundation (NSF) follows closely with 32 funded projects, reflecting its role in promoting scientific research and education across various disciplines, including the biological and environmental sciences related to camels.

Other notable funding agencies include the National Institute of Allergy and Infectious Diseases (NIAID) and the National Cancer Institute (NCI), each supporting 27 and 14 projects respectively. The National Natural Science Foundation of China (NSFC) and the U.S. Department of Health and Human Services (HHS) also contribute significantly, with 13 and 12 projects funded, respectively, showing international and interdisciplinary support for camel research. Additional key funding sources include the Biotechnology and Biological Sciences Research Council (BBSRC) with 9 projects, King Abdulaziz City for Science and Technology (KACST) with 9 projects.

### *The top publishing journals*

The analysis of journals publishing camel research reveals a diverse and interdisciplinary landscape, underscoring the multifaceted nature of this field (Fig 7). The Journal of Camel Practice and Research leads with 20 publications, highlighting its specialisation and central role in disseminating camel-specific studies. This journal serves as a dedicated platform for researchers focusing on various aspects of camel biology, health, and management, reflecting its importance in the academic community dedicated to camel studies.



**Fig 5.** Active institutions of camel research publications in North America and in collaboration with other countries. The listed affiliations include only those having 10 or more publications.

Veterinary and health-related journals also play a crucial role in camel research. The Journal of the American Veterinary Medical Association and Emerging Infectious Diseases have published 11 and 10 articles, respectively, indicating a strong interest in the veterinary and zoonotic disease aspects of camel research.

The diversity of journals, such as Animal Reproduction Science, Journal of Dairy Science, and PLOS ONE, which have each published multiple camel research articles, reflects the wide-ranging applications and interests in camel studies. From reproductive science and dairy production to general scientific inquiries and public health implications,






Fig 6. Funding agencies with five or more camel research projects.





Fig 7. Journals with five or more camel research publications.

these journals show the interdisciplinary nature of camel research.

### Academic disciplines

The bar chart in Fig 8 illustrates the distribution of camel research publications across a wide range

of academic disciplines. The x-axis represents the number of publications, while the y-axis lists the academic fields. This distribution highlights the interdisciplinary nature of camel research and its relevance across multiple domains.



**Fig 8.** Distribution of camel research publications across various academic disciplines.

Agricultural and biological sciences dominate the chart with the highest number of publications, reaching 250. This prominence underscores the central role of camels in agricultural research and their biological significance, including studies on camel physiology, genetics, and their utility in farming systems.

Medicine and veterinary sciences also feature prominently, with 180 and 142 publications, respectively. The significant number of publications in these fields highlights the health and veterinary aspects of camel research, including disease prevention, treatment, and overall animal health management.

Other notable fields include chemistry, genetics and molecular biology, and immunology and


microbiology, each with substantial contributions. These disciplines focus on the molecular and genetic aspects of camel biology, exploring areas such as disease resistance, genetic diversity, and molecular mechanisms underlying camel physiology. The presence of camel research in fields like Environmental Science, Engineering, and Social Sciences further illustrates the diverse applications and broad impact of camel studies.

### ***Bibliometric Coupling Analysis of the Top Collaborations***


The authors are grouped into three clusters based on their collaborative networks (Fig 9). Cluster 1 includes Benaissa, M.H., Burger, P.A., Faye, B., Kaidi, R., and Youngs, C.R. This cluster

indicates a strong interconnected network with notable contributions from Faye, B., who has the highest number of documents (12) and significant link strength (15). Cluster 2 features Ali, M.A., Kayali, G., Shehata, M.M., and Webby, R.J., with Ali, M.A. showing the highest total link strength (14) and document contributions (7). Cluster 3 includes Alagaili, A.N., Daszak, P., Munster, V.J., And Van Doremalen, N., where Munster, V.J. leads with 10 documents and notable link strength (10), indicating significant collaborative influence.

### Co-occurrence of Keywords



data were set to a minimum of 10 times keywords co-occurrence. Of 6731 keywords 176 met this threshold.





along with demographic terms such as “child”, “adolescent”, and “adult”, points to studies focusing on specific regions and population groups.

The co-occurrence network analysis reveals 22 distinct clusters, each representing a focused thematic area in camel research. In the top 10 clusters, Cluster 1, encompassing keywords such as “adolescent”, and “child”, highlights research themes related to public health and human demographics. This cluster suggests significant interest in the impact of camel-related factors on human health, including zoonotic diseases and nutritional studies involving camel products. The presence of keywords like “united states” and “young adult” further indicates that this research might be geographically focused on regions with substantial camel populations or significant camel product consumption.

Clusters 2 through 5 primarily focus on biological and anatomical studies. Cluster 2 includes keywords like “bacteria”, “fermentation”, and “microbiology”, indicating research on microbial communities associated with camels and their fermentation processes. Cluster 3, with keywords such as “anti-bacterial agents”, “antibiotic resistance”, and “*Escherichia coli*”, underscores the importance of addressing bacterial pathogens and the effectiveness of antibiotics in camel health management. Meanwhile, Cluster 4, featuring “artiodactyla”, “mammal”, and “histology”, focuses on the anatomical and histological characteristics of camels and related species, revealing an interest in the structural and morphological analysis of these animals.

Clusters 6 through 10 highlight advanced research methodologies and genetic studies. Cluster 6, with keywords like “major clinical study”, “procedures”, and “randomised controlled trial”, emphasises the use of clinical trials and evidence-based research approaches in camel studies. Cluster 7, including “camel milk”, “milk protein”, and “milk”, focuses on the nutritional and biochemical properties of camel milk, exploring its health benefits and nutritional value. Cluster 8, featuring “gene expression”, “human cell”, and “mouse”, highlights genetic and cellular research involving camels, human, and mouse cells to understand genetic expressions and their implications. Cluster 9, with “antibodies”, “antibody”, and “single-domain antibodies”, centres on immunological studies, exploring various antibodies and their roles in immune responses related to camels. Finally, cluster 10, including “DNA sequence”, “molecular genetics”,

and “sequence analysis, DNA”, underscores the application of molecular genetics and sequencing techniques in camel research, revealing advanced genetic research trends in this field.

The diversity of clusters in the co-occurrence network analysis reflects a robust and interdisciplinary research landscape in camel studies. The 22 clusters cover a wide range of topics, from public health and microbiology to advanced genetic research and clinical studies, indicating the multifaceted nature of camel research and its relevance to various scientific and health-related fields.

### ***Bibliometric Analysis of the Top Highly Cited Publications***

The dataset comprises a variety of highly cited research articles focusing on camel-related studies, spanning several decades and involving multiple countries (Table 1). These studies highlight significant contributions in fields such as infectious diseases, veterinary sciences, and public health. The oldest article in the dataset, “Isolation and structure of an untriakontapeptide with opiate activity from camel pituitary glands” by Li and Chung, dates back to 1976 and has amassed 530 citations, indicating its lasting impact on biochemical research. This paper, published in the prestigious “Proceedings of the National Academy of Sciences of the United States of America,” underscores early explorations into bioactive peptides derived from camels.

Recent studies in the dataset, such as “Chronic Infection with Camelid Hepatitis E Virus in a Liver Transplant Recipient Who Regularly Consumes Camel Meat and Milk” by (Lee *et al*, 2016) and “Middle East Respiratory Syndrome Coronavirus Infection in Dromedary Camels in Saudi Arabia” by (Alagaili *et al*, 2014), reflect contemporary research priorities. These papers, cited 428 and 374 times respectively, discuss the zoonotic diseases and their transmission between camels and humans. The involvement of multiple countries (USA, UK, Singapore for the former, and USA, Saudi Arabia for the latter) highlights the global collaborative effort to understand and mitigate these health risks. Notably, these studies were published in high-impact journals like “Gastroenterology” and “American Society for Microbiology,” further attesting to their scientific significance.

The dataset also includes impactful public health studies, such as those examining the influence of marketing on children’s recognition of

**Table 1.** The Topmost Highly Cited Publications in Camel Research in North America and in Collaboration with other Countries.

Title	Authors	Journal	Affiliations Countries	Year	Citations	References
Isolation and structure of an untriakontapeptide with opiate activity from camel pituitary glands	Li C. H. Chung D.	Proceedings of the National Academy of Sciences of the United States of America	USA	1976	530	(Li and Chung, 1976)
Chronic Infection with Camelid Hepatitis e Virus in a Liver Transplant Recipient Who Regularly Consumes Camel Meat and Milk	Lee G. H. Tan B. H. Chi-Yuan Teo E. Lim S. G. Dan Y. Y. Wee A. Kim Aw P. P. Zhu Y. Hibberd M. L. Tan C. K. Purdy M. A. Teo C. G.	Gastroenterology	USA, UK, Singapore	2016	428	(Lee <i>et al</i> , 2016)
Middle east respiratory syndrome coronavirus infection in dromedary camels in Saudi Arabia	Alagaili A. N. Briese T. Mishra N. Kapoor V. Sameroff S. C. de Wit E. Munster V. J. Hensley L. E. Zalmout I. S. Kapoor A. Epstein J. H. Karesh W. B. Daszak P. Mohammed O. B. Ian Lipkin W.	American Society for Microbiology	USA, Saudi Arabia	2014	374	(Alagaili <i>et al</i> , 2014)
The CAMELS data set: Catchment attributes and meteorology for large-sample studies	Addor N. Newman A. J. Mizukami N. Clark M. P.	Hydrology and Earth System Sciences	USA, UK	2017	313	(Addor <i>et al</i> , 2017)
Brand Logo Recognition by Children Aged 3 to 6 Years: Mickey Mouse and Old Joe the Camel	Fischer P. M. Schwartz M. P. Richards J. W. Goldstein A. O. Rojas T. H.	JAMA: The Journal of the American Medical Association	USA	1991	296	(Fischer <i>et al</i> , 1991)
Seroepidemiology for MERS coronavirus using microneutralisation and pseudoparticle virus neutralisation assays reveal a high prevalence of antibody in dromedary camels in Egypt, june 2013	Perera, R. A. Wang, P. Gomaa, M. R. El-Shesheny, R. Kandeil, A. Bagato, O. Siu, L. Y. Shehata M. M. Kayed A. S. Moatasim Y. Li M. Poon L. L. Guan Y. Webby R. J. Ali M. A. Peiris J. S. Kayali, G.	Eurosurveillance	USA, China, Egypt, Hong Kong	2013	271	(Perera <i>et al</i> , 2013)

MERS coronaviruses in dromedary camels, Egypt	Chu D. K. W. Poon L. L. M. Gomaa M. M. Shehata M. M. Perera R. A. P. M. Zeid D. A. El Rifay A. S. Siu L. Y. Guan Y. Webby R. J. Ali M. A. Peiris M. Kayali G.	Emerging Infectious Diseases	USA, Egypt, Hong Kong	2014	240	(Chu <i>et al</i> , 2014)
RJR Nabisco's Cartoon Camel Promotes Camel Cigarettes to Children	Difranza, J. R. Richards J. W. Paulman P. M. Wolf-Gillespie N. Fletcher C. Jaffe R. D. Murray D.	JAMA: The Journal of the American Medical Association	USA	1991	234	(Difranza <i>et al</i> , 1991)
Antibacterial and antiviral activity of camel milk protective proteins	El Agamy E. S. I. Ruppanneb R.	Journal of Dairy Research	Canada, France, Egypt	1992	220	(El Agamy and Ruppanneb 1992)
Replication and shedding of MERS-CoV in upper respiratory tract of inoculated dromedary camels	Adney D. R. van Doremalen N. Brown V. R. Bushmaker T. Scott D. de Wit E. Bowen R. A. Munster V. J.	Emerging Infectious Diseases	USA	2014	214	(Adney <i>et al</i> , 2014)

cigarette brands and the seroepidemiology of MERS coronavirus in camels. For instance, Fischer *et al* (1991) “Brand Logo Recognition by Children Aged 3 to 6 Years: Mickey Mouse and Old Joe the Camel,” with 296 citations, published in “JAMA: The Journal of the American Medical Association,” investigates early brand recognition and its implications for public health. Similarly, Perera *et al* (2013) study on MERS coronavirus in Egyptian camels, cited 271 times, reveals a high prevalence of antibodies in camels, offering crucial insights into disease transmission dynamics.

Recent interest in camel research fields has surged (Kandeel *et al*, 2023), driven by the growing recognition of camels’ ecological, economic, and health-related significance (Masebo *et al*, 2023). This heightened interest is particularly evident from the sharp increase in publications in the past decade, with substantial contributions from North American institutions in collaboration with international partners. Advances in molecular genetics, vaccines and immunology, and zoonotic disease research have positioned camels as important subjects in understanding disease transmission and developing



**Fig 11.** Summary of camel research statistics in North America.



biomedical applications (Abri and Faye, 2019; Ahmed and Mahmoud, 2023; Al Ramadan *et al*, 2021). Additionally, the unique properties of camel milk and its potential health benefits have spurred research in nutrition and food sciences. The advent of AI technologies is expected to revolutionize camel health and welfare (Alsaleem *et al*, 2024).

### Summary of camel research in North America

Fig 11 presents a detailed summary of camel research statistics in North America from 1859 to 2024. It includes data spanned 493 sources and 786 documents, showing an annual growth rate of 1.89%. The research involves 3127 authors, with 130 single-authored documents, and a notable international co-authorship rate of 52.04%. Each document averages 5.35 co-authors, with a total of 1698 author’s keywords and 26,275 references. The average age of documents is 16.3 years, and they receive an average of 21.38 citations per document.

Fig 12 presents a three-field plot illustrating the associations between countries, affiliations, and keywords in camel research in North America. The left column (AU\_CO) represents various countries, including Italy, USA, Brazil, China, Saudi Arabia, and others. The middle column (AU\_UN) shows affiliations with prominent research institutions such as Inner Mongolia Agricultural University, King Abdulaziz University, Jordan University of Science and Technology, King Saud University, and many more. The right column (ID) lists the most frequently used keywords in camel research, such as “camel,” “article,” “nonhuman,” “animals,” “dromedaries,” and others. This visualisation highlights the

network of international collaborations, institutional contributions, and key research themes prevalent in the field of camel studies in North America. Wile USA is dominating in the right column, the middle column shows collaboration between USA and a lot of universities overseas.

### Strengths and limitations of this study and future directions

The strengths of the bibliometric analysis of camel research are manifold. First, the comprehensive data collection from the Scopus database ensures a robust and extensive dataset, encompassing a wide range of publications and document types. This inclusivity allows for a thorough examination of trends, key contributors, and collaborative networks. Additionally, the identification of significant periods of growth and the mapping of international collaborations highlight the global nature and interdisciplinary scope of camel research.

Despite its strengths, this bibliometric analysis of camel research has several limitations. One major limitation is the reliance on the Scopus database, which, although comprehensive, may not include all relevant publications, especially those in non-indexed regional journals or grey literature. Additionally, the analysis primarily focuses on quantitative measures such as publication counts and citation metrics, which may not fully capture the qualitative aspects of research impact and innovation.

Future direction involves fostering stronger ties between researchers in North America and those in camel-rich regions such as the Middle East and Africa. This includes tackling zoonotic diseases, which




Fig 12. Three-field plot for countries, affiliations and keywords association in camel research in North America.

have significant public health implications, as well as addressing issues related to climate change and sustainability, where camels' resilience offers valuable lessons. Furthermore, the integration of advanced technologies such as artificial intelligence and big data analytics into bibliometric studies can provide deeper insights into research trends and impact, guiding policy-making and funding decisions.

## Conclusions

This bibliometric analysis underscores the growing significance of camel research in North America, driven by the ecological and economic importance of camels. The study reveals robust collaborative efforts between North American researchers and international partners, leading to advancements in understanding zoonotic diseases, genetic diversity, and the nutritional benefits of camel milk. Despite these achievements, gaps remain in integrating advanced technologies and exploring the socio-economic impacts of camel products in non-traditional regions. Addressing these gaps through continued investment and fostering international collaborations is essential to drive innovative research and enhance the overall impact of camel studies.

## Ethics approval and consent to participate

Not applicable

## Consent for publication

Not applicable

## Availability of data and materials

Not applicable

## Competing interests

There is no conflict of interests

## Funding

This work was supported by the Deanship of Scientific Research, Vice Presidency for Graduate Studies and Scientific Research, King Faisal University, Saudi Arabia (Project# KFU241218).

## Authors' contributions

"Conceptualisation, M.K.; methodology, M.K.; software, M.K.; validation, M.K.; formal analysis, M.K.; investigation, A.A. and M.K.; resources, M.K.; data curation, A.A. and M.K.; writing—original draft preparation, M.K.; writing—review and editing, A.A. and M.K.; funding acquisition, A.A. and M.K. All authors have read and agreed to the published version of the manuscript.

## Acknowledgements

This work was supported by the Deanship of Scientific Research, Vice Presidency for Graduate Studies and Scientific Research, King Faisal University, Saudi Arabia (Project# KFU241218).

## References

- Abri MAA and Faye B. Genetic improvement in dromedary camels: challenges and opportunities. *Frontiers in Genetics*. 2019; 10:428917.
- Adah AS, Ayo JO and Adah DA. Unique physiological and behavioural adaptive features of the one-humped camel (*Camelus dromedarius*) to arid environments. *Journal of Applied Veterinary Sciences*. 2023; 8:57-64.
- Addor N, Newman AJ, Mizukami N and Clark MP. The CAMELS data set: Catchment attributes and meteorology for large-sample studies. *Hydrology and Earth System Sciences*. 2017; 21:5293-5313.
- Adney DR, van Doremalen N, Brown VR, Bushmaker T, Scott D, de Wit E, Bowen RA and Munster VJ. Replication and shedding of MERS-CoV in upper respiratory tract of inoculated dromedary camels. *Emerging Infectious Diseases*. 2014; 20:1999-2005.
- Ahmed A and Mahmoud K. MVA-based vaccine efficacies on the immune response and seroprevalence of MERS-CoV: A systematic review. *Journal of Camel Practice and Research*. 2023; 30:9-17.
- Al Ramadan SY, Salem KTA-M, Alshubaith IH, Al-Ali AM, Abohelaika S, Moqbel MS and Alluwaimi AM. Selected aspects of camel immune system and immune responses. *Open Journal of Veterinary Medicine*. 2021; 11:177-211.
- Alagaili AN, Briesse T, Mishra N, Kapoor V, Sameroff SC, de Wit E, Munster VJ, Hensley LE, Zalmout IS, Kapoor A, Epstein JH, Karesh WB, Daszak P, Mohammed OB and Ian Lipkin W. Middle east respiratory syndrome coronavirus infection in dromedary camels in Saudi Arabia. *mBio*. 2014; 5.
- Alsalem A, Shousha S, Marzok M, Afzal S, Alameen A, Albokhadaim I and Kandeel M. Transformative technologies for the future of camel welfare: artificial intelligence for improved diagnostics, therapeutics and health outcomes. *Journal of Camel Practice and Research*. 2024; 31:1-9.
- Aria M and Cuccurullo C. bibliometrix: An R-tool for comprehensive science mapping analysis. *Journal of Informetrics*. 2017; 11:959-975.
- Arruda H, Silva ER, Lessa M, Proenca D, Jr. and Bartholo R. VOSviewer and Bibliometrix. *Journal of the Medical Library Association*. 2022; 110:392-395.
- Camel Sciences and Economy in the World: Current Situation and Perspectives.
- Chu DKW, Poon LLM, Gomaa MM, Shehata MM, Perera RAPM, Zeid DA, El Rifay AS, Siu LY, Guan Y, Webby RJ, Ali MA, Peiris M and Kayali G. MERS coronaviruses in dromedary camels, Egypt. *Emerging Infectious Diseases*. 2014; 20:1049-1053.

- Difranza JR, Richards JW, Paulman PM, Wolf-Gillespie N, Fletcher C, Jaffe RD and Murray D. RJR Nabisco's Cartoon Camel Promotes Camel Cigarettes to Children. *JAMA: The Journal of the American Medical Association*. 1991; 266:3149-3153.
- El Agamy ESI and Ruppanneb R. Antibacterial and antiviral activity of camel milk protective proteins. *Journal of Dairy Research*. 1992; 59:169-175.
- Fischer PM, Schwartz MP, Richards JW, Goldstein AO and Rojas TH. Brand Logo Recognition by Children Aged 3 to 6 Years: Mickey Mouse and Old Joe the Camel. *JAMA: The Journal of the American Medical Association*. 1991; 266:3145-3148.
- George AS. Camel milk production as an adaptation to climate change induced drought in East Africa. *Partners Universal Multidisciplinary Research Journal*. 2024; 1:109-126.
- Hedayat-Evrigh N, Khalkhali-Evrigh R and Bakhtiarzadeh MR. Genome- Wide identification and analysis of variants in domestic and wild bactrian camels using whole- genome sequencing data. *International Journal of Genomics*. 2020; 2020:2430846.
- Kandeel M, Morsy MA, El-lateef A, Hany M, Marzok M, El-Beltagi H, Alkhodair K, Albokhadaim I and Venugopala KN. A Century of" Camel Research": A Bibliometric Analysis. *Frontiers in Veterinary Science*. 2023; 10:700.
- Lee GH, Tan BH, Chi-Yuan Teo E, Lim SG, Dan YY, Wee A, Kim Aw PP, Zhu Y, Hibberd ML, Tan CK, Purdy MA and Teo CG. Chronic infection with camelid hepatitis e virus in a liver transplant recipient who regularly consumes camel meat and milk. *Gastroenterology*. 2016; 150:355-357.e353.
- Li CH and Chung D. Isolation and structure of an untriakontapeptide with opiate activity from camel pituitary glands. *Proceedings of the National Academy of Sciences of the United States of America*. 1976; 73:1145-1148.
- Masebo NT, Zappaterra M, Felici M, Benedetti B and Padalino B. Dromedary camel's welfare: literature from 1980 to 2023 with a text mining and topic analysis approach. *Frontiers in Veterinary Science*. 2023; 10.
- Ming L, Yuan L, Yi L, Ding G, Hasi S, Chen G, Jambl T, Hedayat-Evrigh N, Batmunkh M and Badmaevna GK. Whole-genome sequencing of 128 camels across Asia reveals origin and migration of domestic Bactrian camels. *Communications Biology*. 2020; 3:1.
- Perera RA, Wang P, Gomaa MR, El-Shesheny R, Kandeil A, Bagato O, Siu LY, Shehata MM, Kayed AS, Moatasim Y, Li M, Poon LL, Guan Y, Webby RJ, Ali MA, Peiris JS and Kayali G. Seroepidemiology for MERS coronavirus using microneutralisation and pseudoparticle virus neutralisation assays reveal a high prevalence of antibody in dromedary camels in Egypt, June 2013. *Eurosurveillance*. 2013; 18.
- Sahoo A. Camel: a fast-declining animal species but can strive with its unique climate resilience and 'desert to medicine' application". *EC Veterinary Science*. 2020; 5:43-57.
- Wong D. VOSviewer. *Technical Services Quarterly*. 2018; 35:219-220.



## JOURNAL OF CAMEL PRACTICE AND RESEARCH BEING THE MOST PROLIFIC SOURCE OF RESEARCH ON CAMEL MILK AND URINE

Research on camel milk and urine (CMUR) has gained significant attention because of its nutritional and medicinal applications. The aim of this study was to conduct a bibliometric analysis of the CMUR from 1947 to 2023. Following PRISMA rules, the Scopus database was used to extract 1338 English-language studies, which were analysed using Scopus Analytics, Bibliometrix, and VOSviewer. The findings indicated that 7.78% of studies focused on camel urine within the CMUR domain. The analysis showed a steady increase in the number of publications over 77 years ( $R\text{-squared} = 0.8308$ ). The CMUR encompasses a wide range of research disciplines, with particular emphasis on the agricultural and medical fields. Saudi Arabia is the top producer of CMUR. In terms of the most prolific sources of publications in the CMUR, the Journal of Camel Practice and Research takes the lead, with 80 papers. This was followed by the Journal of Dairy Science with 51 papers, the Emirates Journal of Food and Agriculture with 39 papers, the International Dairy Journal with 39 papers and Food Chemistry with 36 papers.

King Saud University contributed 103 documents to the CMUR, making it the most prolific institution globally. The co-word analysis identified seven prominent research themes within the CMUR, which can be further categorised into four distinct clusters. Notably, 'Probiotics', 'Anti-inflammatory', and 'Diabetes' emerged as trending subjects. The analysis of the CMUR yielded significant insights into the present research trends, identified gaps in the existing knowledge, and provided valuable recommendations for future investigations.

(Source: Abdelwahab, S. I., Taha, M. M. E., Mariod, A. A., Mohamed, H. Y., Farasani, A. M., & Jerah, A. (2024). Unleashing the potential: Camel Milk and Urine Research insights for performance, collaboration, structure and future trends (1947–2023). *Cogent Food & Agriculture*, 10(1). <https://doi.org/10.1080/23311932.2024.2311435>)

## GENETIC STRUCTURE OF CAMELS UNDER FOCUS

Thirteen countries participated in a conference from April 7 to 9, 2024 in Riyadh which looked at changing the genetic structure of camels to improve milk and meat yields. The three-day conference focussed on ways to improve the performance of camels based mainly on genetic tools. International experts on camels were invited to present new techniques for implementing a program for camels. In December 2011, a national workshop on camel biodiversity was held in the Saudi capital with the help of two international experts. On that occasion, it was concluded that there was a need for further studies on camels. The conference was held as part of the technical cooperation programme between the Ministry of Agriculture and the FAO.

## 3<sup>rd</sup> ANNUAL INTERNATIONAL CAMEL PARADE IN PARIS

The event in Paris was organised by the French Federation for the Development of Camelids in France and Europe, under the umbrella of the International Camel Organisation, and was sponsored by the Saudi Ministry of Culture and the Kingdom's Camel Club. This was the third year in which the event took place. The event was first held in January 2019 and repeated in 2022. The participants in the parade of camels, llamas, alpacas and other members of the camelid family of creatures participated from more than 50 representatives of camel-related organisations from more than 30 countries, along with camel breeders, government officials, and others with an interest in the animals, and entertainers from various branches of the performing arts.

# SOME ASPECTS OF CAMEL MILK AND ITS SAFETY FOR HUMAN CONSUMPTION-AN OVERVIEW

Shehadeh Kaskous<sup>1</sup> and Michael W. Pfaffl<sup>2</sup>

<sup>1</sup>Department of Research and Development, Siliconform, Schelmengriesstrasse 1, 86842 Türkheim, Germany

<sup>2</sup>Department of Animal Physiology and Immunology, School of Life Sciences, Technical University of Munich, 85354 Freising, Liesel-Beckmann-Straße 1, Germany

## ABSTRACT

Camel milk is in high demand in many countries around the world. There appears to be increasing awareness of the use of camel milk in human diet. Therefore, the aim of this study was to determine the importance of camel milk and its safety for human consumption. Camel milk contains numerous antimicrobial components and protective factors, making it a unique milk compared to the milk of other lactating animal species. Camel milk production varies between 1000 and 12000 kg during the lactation period of 8 to 18 months, with large differences between geographical regions, feeding and husbandry conditions. The composition of camel milk is 2.5-5.5% fat, 2.2-4.5% protein, 2.5-5.5% lactose, 0.7-1.0% ash and 8.0-15% dry matter. To increase milk production for each camel and improve the quality and safety of raw camel milk, the use of a suitable milking machine such as StimuLactor is necessary. In addition, correct husbandry, feeding and hygiene measures must be considered. In conclusion, regarding the importance of camel milk and the associated related health benefits of its bioactive ingredients, improvements in milking hygiene, milk storage and transport conditions are required to ensure the quality of camel milk meets consumer needs.

**Key words:** Camel, human milk, milking machine, milk yield

Camel milk is one of the most important nutritional sources for the population in many arid and semi-arid areas of the world, as it contains almost all the vital nutrients needed in dry climate. The camel population in the world is approximately 35 million, about 95% of which are dromedaries (FAO, 2022). Camel milk yield varies between 1000 and 12000 kg during the lactation period of 8 to 18 months (Boujenane, 2020; Swelum *et al*, 2020). However, the amount of milk per lactation depends on many factors such as breed (Type), animal health, lactation stage, lactation number, living conditions and season (Yamina *et al*, 2013; Chamekh *et al*, 2020; Swelum *et al*, 2021; Boudalia *et al*, 2023). Camel milk was named as the most valuable product (Davati *et al*, 2015) and it is known as “white gold of the desert” (Wernery, 2006). However, camel milk contains numerous minor components that have special bioactive properties (Kaskous and Pfaffl, 2017; Swelum *et al*, 2021; Oselu *et al*, 2022). These are present in significant concentrations and are extremely important and beneficial to human nutrition and health (Kaskous, 2016). Therefore, camel milk is usually consumed as a fresh or naturally fermented product and is therefore not pasteurised (Mehari *et al*, 2007; Matofari *et al*, 2013; Abera *et al*, 2016; Mwangi *et al*, 2016; Serda *et al*, 2018).

But camel milk is an excellent food for the growth of microorganisms (Zangerl, 2007; Matofari *et al*, 2013) and non-heat-treated milk and raw milk products are the main cause of diseases caused by pathogens (De Buyser *et al*, 2001; Smits *et al*, 2023). It is noteworthy that the milk from a healthy udder is usually “sterile” (Johnson *et al*, 2015) because the camel udder is protected by a variety of defence mechanisms such as innate or specific immunity as well as physiological peculiarities and is only contaminated with germs when it passes through the teat canal (Zangerl, 2007). In addition, the germs that get into the milk come from the udder-and teat-surface, the stall, the air, the water, the feed, the milker, and the milking equipment. Contamination of raw camel milk can be very high if udder health is not checked, plastic containers are used during milking and storage, hygiene measures are not carried out during milking, and water is not available to clean the teat and udder before milking (Mulwa *et al*, 2011; Ismaili *et al*, 2019; Atigui *et al*, 2023). During milk transportation, especially when the route is long, and poor and the milk is not refrigerated, the microbial content of the milk multiplies rapidly (Mulwa *et al*, 2011; Ismaili *et al*, 2019). Due to the many contaminated factors, it has been shown that the poor hygiene status in the

SEND REPRINT REQUEST TO SHEHADEH KASKOUS [email: skaskous@siliconform.com](mailto:skaskous@siliconform.com)

production and handling of camel milk as well as the lack of low-cost post-harvest technologies mean that camel milk has lost its potential to improve the living conditions of the camel farmer (Matofari *et al*, 2007; Ismaili *et al*, 2019; Hassen *et al*, 2022) and this limitation leads to losses in quality and quantity after harvest, especially physical-chemical and microbiological deterioration of the milk (Odongo *et al*, 2016; Ismaili *et al*, 2019; Atigui *et al*, 2023). The production of hygienically perfect, ready-to-process milk places high demands on the camel farmer. This means optimal animal husbandry, animal care, good feeding, ideal milking machines and cooling of the milk during storage and transport to deliver consistently high-quality milk to the consumer. To achieve this goal, the transition from traditional production systems to intensive production systems must be accelerated. In the following some aspects of camel milk and its safety for human consumption are presented and discussed.

### Composition of camel's milk

The composition of camel milk is different from those of other lactating animals' species particularly due to its protein composition, milk fat structure and mineral and vitamin content (El-Hatmi *et al*, 2015; Abduku and Eshetu, 2024), but it contains all the important nutrients (Farah, 2011; Kanca, 2017; Boujenane, 2020; Swelum, 2021; Lajnaf *et al*, 2023). Camel milk contains the following components: 2.2-4.5% protein, 2.5-5.5% fat, 2.5-5.5 lactose, 0.7-0.95% ash and 8.0-15% total dry matter (El-Hatmi *et al*, 2006; Konuspayeva *et al*, 2009; Kaskous *et al*, 2012; Ali *et al*, 2019; Ismaili *et al*, 2019; Roy *et al*, 2020; Seifu 2022; Kaskous, 2023). However, the composition varies greatly and depends on various factors, such as the breed (type), individual, husbandry and feeding condition, age, milking interval, stage of lactation, number of lactations, season, management, and udder health (Wernery, 2007; Shuiep *et al*, 2008; El-Hatmi *et al*, 2015; Jrad *et al*, 2015; Patel *et al*, 2016; Ismaili *et al*, 2019; Swelum *et al*, 2021; Behrouz *et al*, 2022; Kraimia *et al*, 2024). In general, camel milk has lower amounts of fat, protein, and lactose compared to bovine milk (Smits *et al*, 2011; Konuspayeva *et al*, 2011; Arab *et al*, 2014; Alhaj *et al*, 2022).

### Camel milk Proteins

Camel's milk is a rich source of proteins with potential anti-microbial and protective activity (Kaskous and Pfaffl, 2017; Mohamed *et al*, 2020; Lajnaf *et al*, 2023). It was observed that camel milk has 21 different amino acids compared to 18 amino

acids in bovines. Such amino acids are the synthesis basis of all proteins in camel milk (Fig 1). Practical results have shown that the average protein content of camel milk under European conditions was 3.34% in the Netherlands (Smits *et al*, 2011), 2.28% in Germany (Kaskous, 2019) and 2.39% in Switzerland (Kaskous, 2023), while the average protein content in camel milk outside Europe was 3.1% (Konuspayeva *et al*, 2009; Al haj and Al Kanhal, 2010). In general, the protein content of camel milk ranges between 2.2 and 4.5% (Konuspayeva *et al*, 2009; Yadav *et al*, 2015; Kula and Dechasa, 2016; Patel *et al*, 2016; Swelum *et al*, 2021; Seifu, 2022). This difference is due to environmental conditions and the availability of food and water for the lactating camels (Yamina *et al*, 2013; Ismaili *et al*, 2019). Camel milk proteins appear as: Casein, whey proteins, fat globule membrane proteins. However, the average casein content and whey protein content of camel milk vary between 1.9 and 2.3% and 0.7 and 1.0%, respectively (Farah, 2011). Caseins make up the highest proportion of the protein fraction in camel milk (Hamed *et al*, 2012; Swelum *et al*, 2021). This represents 65-70% of total proteins compared to an average of 83% in bovine milk (Khaskheli *et al*, 2005; Frister, 2007; Hamed *et al*, 2012). Like cow's milk, casein in camel milk consists of four fractions. These are  $\alpha$ s1,  $\alpha$ s2,  $\beta$  and k-casein (Al haj and Al kanhal, 2010; Abbas *et al*, 2013). However, camel milk contains more  $\beta$ -casein (75%) than  $\alpha$ -casein (21%) of total casein, compared to cows' milk, which contains approximately the same content of  $\beta$ -casein and  $\alpha$ -casein (36 and 38%) (Devendra *et al*, 2016; Kappeler *et al*, 2003; Yirda *et al*, 2020). Interestingly, camel milk contains very little k-casein (3.5%) of total casein, compared to the k-casein content (13%) in cow's milk (Devendra *et al*, 2016; Mbye *et al*, 2022). It was observed that whey proteins in camel milk constituted about 30-35% of total protein (Khaskheli *et al*, 2005; Jilo and Tegegne, 2016; Dugassa, 2021; Vincenzetti *et al*, 2022). Camel milk as human milk does not contain  $\beta$ -Lactoglobulins (El-Hatmi *et al*, 2007; Elagamy *et al*, 2009; Kaskous and Pfaffl, 2017; Rahmeh *et al*, 2019). Therefore,  $\alpha$ -lactalbumin is the main whey protein in camel milk (Wernery, 2007; Redington *et al*, 2016; Lajnaf *et al*, 2023). Camel milk contains other important components of whey proteins such as serum albumin with a variation of 2.9 and 13.8 g/l (El-Hatmi *et al*, 2006) and lactoferrin with 227-229 mg/l (Konuspayeva *et al*, 2007; Kaskous *et al*, 2012). In addition, several reports have shown that camel milk has higher immunoglobulins (0.72 g/l) than cow's milk (0.47 g/l) (Kaskous and Pfaffl, 2017).



## Camel milk fat

The fat content of camel milk varies between 2.5 and 5.5% depending on the nutritional status, stage of lactation, breed and the season of the year (Mal *et al*, 2006; Yamina *et al*, 2013; Devendra *et al*, 2016; Kaskous, 2019; 2023; Abduku and Eshetu, 2024). The fat globules in camel milk are the smallest among all ruminants ranging from 0.1 to 18 micrometers in diameter (El-Zeini, 2006), and these do not naturally aggregate due to the absence of agglutinin (Khalesi *et al*, 2017). 96% of the fat in camel milk is triacylglycerol (Nikkhah, 2011; Dugassa, 2021). Compared to cow's and buffalos' milk, camel milk fat contains lower concentrations of short-chain fatty acids and higher concentration of long-chain fatty acids (C<sub>14</sub>-C<sub>18</sub>) (Abu-Lehia, 1989; Konuspayeva *et al*, 2008; Abbas *et al*, 2013; Sara *et al*, 2022). The saturated fatty acid (SFA) in camel milk varied between 50% (Bactrian camel milk) and 60% (dromedary camel milk) of total fatty acids (Rahmeh *et al*, 2019) and about 65% of total fatty acids according to Wang *et al* (2011) and Oselu *et al* (2022). However, the monounsaturated fatty acids (MUFA) in dromedary camel milk were 56-80% of total fatty acids (Medhammar *et al*, 2012). In addition, the proportion of unsaturated fatty acids in camel milk was between 35 and 50% of the total fatty acids (Izadi *et al*, 2019). It is noteworthy that the proportion of unsaturated fatty acids in the fatty acid pattern of camel milk in the intensive production system was 43.1% (Stahl *et al*, 2006). In addition, the fat content in camel milk was found to be highly dependent on the stage of lactation, with the highest concentration occurring in the first three months of lactation (34.67±1.30 g/l), compared to the concentration in the middle (27.40±5.69 g/l) and the end of lactation (29.12±2.83 g/l) (Kraimia *et al*, 2024).

## Camel milk lactose

The lactose content in camel milk is not stable, although it is involved in the osmotic pressure in the udder. It is noteworthy that a different mechanism exists in camels, as regular feeding and water are not available in many regions of the world. Therefore, different concentrations of lactose have been found in camel milk. Milk lactose content was observed to be 4.3 ±0.2% higher in camels deprived of water for four days compared to 4.1±0.2% for 16 days (Bekele *et al*, 2011). These results illustrate that the variation in lactose content in camel milk is mainly related to water intake and the type of feed consumed (Al haj and Al kanhal, 2010; Kula and Dechasa, 2016). In general, the lactose content in camel milk

varied between 2.5% and 5.5% (Khan and Iqbal, 2001; Konuspayeva *et al*, 2009; Devendra *et al*, 2016; Faraz, 2020; Dugassa, 2021; Alhaj *et al*, 2022; Karaman *et al*, 2022). In addition, lactose content in camel milk was observed to be high in the first months of lactation (42.5 g/l), followed by a significant ( $P<0.05$ ) decrease at the end of lactation (38.35 g/l) (Kraimia *et al*, 2024). At this point, it must be said that the significant change in lactose concentration in camel milk is due to the health of the udder. It is known that in clinical or subclinical mastitis, the lactose content of the milk decreases and new synthesis is reduced (Schulz, 2003). New research has shown that the average lactose content in the milk of a healthy camel udder was around 4.09±0.03% after using new "StimuLactor" milking machines on a practical farm in Switzerland (Kaskous, 2023).

## Camel milk vitamins and minerals

Studies on the vitamin content of camel milk have been carried out for a long time (Sawaya *et al*, 1984; Farah *et al*, 1992). Stahl *et al* (2006) found that fresh dromedary milk contains less vitamin A, E, B<sub>1</sub> and  $\beta$ -carotene than cow's milk. An interesting aspect: The vitamin C content in camel milk (34.16 mg/l) was 2 to 3 times higher than in cows' milk (Farah *et al*, 1992) and 6 times higher than in human milk (Gizachew *et al*, 2014). However, the vitamin C content of camel milk was between 24 and 52 mg/kg and was therefore, 2-4 times higher than that of cow's milk (15±6.3 mg/l) (Mehaia, 1994; Stahl, 2005; Kamal and Karoui, 2017). The high vitamin C content in camel milk makes nutritional sense because fruits and vegetables are rarely part of the human diet in dry areas and intake of vitamin C *via* camel milk can help prevent vitamin deficiencies in humans. It has also been found that the lower carotene content makes the colour of camel milk whiter compared to cow's milk (Stahl *et al*, 2006; Devendra *et al*, 2016). In addition, studies using Bactrian camel milk showed that camel milk was rich in vitamin D and riboflavin and consumption of two cups daily provided 160% of the recommended dietary intake of vitamin D (5 µg/day) and riboflavin (0.5 mg/day) (Wijesinha-Bettoni and Burlingame, 2013).

Furthermore, research has shown that camel milk is rich in minerals such as calcium, phosphorus, sodium, potassium, chloride, iodine, and magnesium (Shamsia, 2009; Gizachew *et al*, 2014; Aljumaah *et al*, 2012; Alhadrami and Faye, 2016). The average contents of Ca, K, Mg and Na in camel milk were 1.47±0.38, 0.98±0.24, 0.07± 0.01, and 0.65±0.11 g/l,

respectively (Jrad *et al*, 2015). However, the average ash content in camel milk varied between 7.5 and 8.8 g/l (Jrad *et al*, 2015; Abdel Galil *et al*, 2016). It is noteworthy that the content of iron, zinc and copper in camel milk was higher than in cows’ milk (Singh *et al*, 2006). It should be noted that many factors such as breed (type), diet, water consumption and analytical methods can affect the mineral content of camel milk (Mehaia *et al*, 1995; Haddadin *et al*, 2008). A new study has shown that the season can have an impact on the minerals in camel milk (Kraimia *et al*, 2024) (Table 1).

### Safe raw camel milk for human consumption

In many countries, particularly in Africa and the Middle East, the consumption of raw camel milk has increased and is often considered as a “health food” with positive effects on digestion, the immune system and for treatment of various diseases (Gonfa *et al*, 2001; Seifu, 2007; Zaitlin *et al*, 2013). According to an online survey of 852 consumers conducted in the United Arab Emirates (UAE), about 58.4% consumed unpasteurised camel milk (Cheikh Ismail *et al*, 2022). The reason for consuming unpasteurised camel milk was that this milk strengthens the immune system and has higher nutritional content (Smits *et al*, 2023). Therefore, raw camel milk was used to treat, alleviate, or prevent health conditions such as diabetes, autism, cancer, dementia, allergies, and parasites (Kaskous, 2016). In Ethiopia, most of camel milk is consumed in the raw state without

any heat treatments (Eyassu, 2007; Mehari *et al*, 2007; Roess *et al*, 2023). In the Arabian Peninsula, consumption of unpasteurised camel milk is also common (Omrani *et al*, 2015). Furthermore, fresh and fermented camel milk has been also used in India, Russia, and Sudan for human consumption as well as for treatment of a series of diseases (Kumar *et al*, 2016). Unpasteurised sour camel milk and fresh cow’s milk form the core of the Somali diet (Seifu, 2007; Sadler and Catley, 2009; Carruth, 2014). On the other hand, some countries like UAE, Australia and USA warned that camel raw milk was not generally recognised as safe or effective for the therapeutic uses. However, consuming raw or unpasteurised dairy products poses several known risks, especially if refrigeration is not ensured (Zaitlin *et al*, 2013; Carruth *et al*, 2017). The Food and Drug Administration (FDA) in USA warned that if the camel farmer was going to market their product as a “drug”, they needed to get federal approval, which would require the farm to provide scientific data demonstrating the safety and effectiveness of their product. Moreover, FDA warned that a consumption of raw camel milk is a health risk, because it is associated with foodborne illness caused by pathogens including *Campylobacter*, *Escherichia coli*, *Listeria*, *Brucella*, *Staphylococcus* and *Salmonella* (Yamina *et al*, 2013; Swinburne, 2017; Wernery *et al*, 2017; Dadar *et al*, 2019). Based on the observation of Spargue *et al* (2012) that brucellosis a bacterial disease caused by various infectious *Brucella*

**Table 1.** Some factors that affect the concentration of minerals in camel milk, according to Kraimia *et al* (2024), with some changes.

Factors		Minerals in camel milk					
		Na (g/l)	K (g/l)	Ca (g/l)	Mg (g/l)	I (g/l)	P (g/l)
Day	Morning	0.39	2.44	1.90	0.07	0.13	0.70
	Evening	0.45	2.39	1.73	0.07	0.13	0.83
	P<0.05			*			**
Stage of lactation	Beginning	0.43	2.63	1.51	0.07	0.15	0.72
	Middle	0.44	2.35	1.67	0.07	0.14	0.65
	End	0.41	2.37	1.75	0.07	0.12	0.69
	P<0.05					*	
Lactation number	1	0.49	2.28	1.67	0.07	0.13	0.63
	2-6	0.46	2.19	1.48	0.07	0.12	0.78
	>6	0.44	2.40	1.66	0.08	0.11	0.66
	P<0.05						
Season	Winter	0.46	2.26	1.48	0.07	0.12	0.66
	Spring	0.39	2.44	1.90	0.07	0.13	0.70
	Summer	0.45	2.38	1.55	0.08	0.15	0.68
	Fall	0.52	2.28	1.61	0.08	0.13	0.77
	P<0.05	***		***	***	**	

\*: P<0.05; \*\*: P<0.01; \*\*\*: P<0.001

species and is transmitted through both close contact with camels and consumption of raw camel milk. However, camel brucellosis has been diagnosed in all camel-rearing countries except Australia and mainly depends on the management system (Wernery, 2014). Zimmermann (2016) reported that one of the primary risks of camel's milk is consumed in unpasteurised form. Reviewed by Roess *et al* (2023) found that consumption of days-old unrefrigerated raw camel milk was significantly associated with gastrointestinal symptoms. The Saint Louis Institute for conservation medicine studied the consumption of camel milk in northern Kenya, where around 10% of people drink unpasteurised camel milk, exposing themselves to a few animal-based pathogens. The study found a higher prevalence of pathogenic bacteria in raw camel milk than in sheep and cattle milk. Furthermore, Musinga *et al* (2008) found that contaminations of raw camel milk in Kenya can occur along the chain from producers to final consumers and the consumption of raw camel milk should be of major concern to public health. Research in Saudi Arabia and Morocco found that raw camel milk samples were contaminated due to poor handling practices and hygiene conditions (El-Ziney, 2007; Alaoui Ismaili *et al*, 2019). Matofari *et al* (2013) found in Kenya that salmonella enteric occurrence along the camel milk chain had an incidence of 13% with the highest being in the farm environment. The sources of this pathogen may constitute the risk factors that are associated with its prevalence in the environment. However, milk from individuals without clinical or subclinical mastitis may also contain mastitis-causing pathogens, and other factors, such as soil, water, and pastoralists (Elmoslemany *et al*, 2010). Farm management practices and seasonality have also been shown to have a significant impact on the contamination of raw camel milk (Shuiep *et al*, 2008; Smits *et al*, 2023). The control of raw camel milk is an important issue for human consumption (Alebie *et al*, 2021). In fact, pasteurisation is a good preservation method to produce high-quality milk.

### Raw camel milk and udder inflammation

Camel udder can get clinical or subclinical mastitis, like other dairy animals (Alebie *et al*, 2021; Rahmeh *et al*, 2022). A high percentage of subclinical mastitis in camels is reported by several authors (Obeid *et al*, 1996; Almaw and Molla, 2000; Wanjohi *et al*, 2013; Niasari-Naslaji *et al*, 2016) and the values varied between 15 and 70% (Bhatt *et al*, 2004; Abera *et al*, 2010; Seifu and Tafesse, 2010; Alamin *et al*, 2013).

In general, 46% of the global camel population suffers from mastitis (Aqib *et al*, 2022). It was shown that mastitis pathogens of the dromedary are the same as cultured from the mammary gland of bovines and these are *Streptococcus agalactiae*, *Staphylococcus aureus*, *Coagulase-negative Staphylococcus*, *Streptococcus bovis*, *Streptococcus uberis*, *Streptococcus dysgalactiae* (Wernery *et al*, 2008). In the traditional husbandry systems, poor management, and unhygienic milking lead to mastitis in camels (Obeid *et al*, 1996; Almaw and Molla, 2000). The results from Golestan province in Iran have shown that out of 243 camel milk samples from individual quarters (95 milking camels), 18.1% were subclinical mastitis and somatic cell count values beyond  $306 \times 10^3$  cells/ml could be considered as subclinical mastitis in camel (Niasari-Naslaji *et al*, 2016). Bekele and Molla (2001) reported that, out of 152 camels in Afar Region, north-eastern Ethiopia examined, 19 (12.5%) were diagnosed as clinical mastitis cases based on clinical signs and bacteriological examinations. The main mastitis pathogens isolated were *Staphylococcus aureus*, *coagulate negative staphylococci*, *Streptococcus agalactiae*, *S. dysgalactiae*, and other species of *streptococci*, *Pasteurella haemolytica* and *E. coli*. Similar results have been shown by Wanjohi *et al* (2013) that subclinical mastitis was prevalent in dromedary camels of two districts of north-eastern province of Kenya, and that Gram-positive cocci (*Staphylococcus* and *Streptococcus*) were the dominant mastitis pathogens isolated. Other isolated bacteria were found as *Klebsiella/Enterobacter*, *Escherichia coli* and *Bacillus*. In addition, *Staphylococcus aureus*, *Streptococcus agalactiae*, *Escherichia coli* and *Corynebacterium bovis* were the main pathogens of camel mastitis (Aqib *et al*, 2022). Rahmeh *et al* (2022) found that the genera *Staphylococcus*, *Streptococcus*, *Schlegella*, unclassified *Enterobacteriaceae*, *Lactococcus*, *Jeotgalicoccus* and *Klebsiella* were abundant in mastitis milk compared to healthy samples. Abdel Gadir Atif *et al* (2006) have performed comparison of California mastitis test (CMT), somatic cell counts (SCC) and bacteriological examinations for detection of camel mastitis in Ethiopia. A total of 956 quarter milk samples from 253 camels were detected. 59.7% quarter milk samples had microorganisms. A positive correlation was found between CMT scores and bacteriological classes ( $P < 0.001$ ) and SCC ( $p < 0.001$ ). Detection of subclinical mastitis in dromedary camels using somatic cell counts, California mastitis test and udder pathogen was also done in Saudi Arabia (Saleh and Faye, 2011). A total of 120 quarter milk samples from 30 clinically healthy dromedary camels were cultured. SCC varied from 9000 to 2 000.000



cells/ml with an average of 125000. Intramammary infections were present in most of examined quarter milk samples. The current review by Aqib *et al* (2022) showed the forms of camel mastitis. Clinical mastitis is characterised by hardening and swelling of the udder, pain on palpation, and visible changes in the colour and texture of the milk. Subclinical mastitis is inflammation without obvious signs (El Tigani-Asil *et al*, 2020) that can be detected by CMT, SCC and microbiological examination. It is noteworthy that an increase in SCC above the physiological value not only indicates a problem with udder health, but also leads to reduced milk yield, a change in milk composition, an impairment of milk processing and a change in the bioactive ingredients of camel milk (Kaskous *et al*, 2021)

Based on that, the disease has been reported in numerous camel-producing countries in Africa and Asi (Aqib *et al*, 2022). The following table presents the results of some works on contaminated udder quarters with microorganisms (Table 2).

In Sudan, raw camel milk samples were collected and the isolated aerobic bacteria (115 isolates) were identified as Gram-negative (85.26%),

while 14.73% of samples were Gram-positive. The authors emphasised that raw camel milk is a source of many bacteria which may lead to health hazards for men (Elhaj *et al*, 2014). In the southern province of Jordan, raw camel milk samples were collected from 90 dromedary camels. About 21% of the camels revealed clinical signs of mastitis. The most predominant bacteria isolates were *Staphylococcus aureus*, *Streptococcus* spp. *Micrococcus* spp. and *Corynebacterium* spp. (Hawari and Hassawi, 2008). The microbial quality of raw camel milk in United Arab Emirates was investigated, 50 samples were analysed for: Aerobic plates count, total coliform, total *Staphylococcus aureus*, total yeast, and Mold. The results indicated that the mean value of aerobic plate count was  $1.8 \times 10^5$  cfu/ml, the mean value of total coliform was  $6.8 \times 10^{-1}$ , the mean value of *Staphylococcus aureus* was  $1.2 \times 10^3$ , and the yeast mean value was  $4.1 \times 10^{-1}$  cfu/ml (Omer and Eltinay, 2008). Furthermore, it should also be noted that the calf may be a source of prevalence of microbes in camel milk. So, in many countries, the calf will have their mother suckle to induce the milk ejection reflex. But the calf may compromise the udder hygiene since

**Table 2.** Raw camel milk with subclinical and clinical mastitis in some countries.

Country	Number of milk samples (quarter or udder)	Positive samples of subclinical or clinical mastitis (%)	References
Egypt	90 (udder)	87.78	Asfour and Anwer (2015)
Ethiopia	956 (quarter)	59.7%	Abdel Gadir Atif <i>et al</i> (2006)
Ethiopia	205 (quarter)	37.6%	Abdurahman (2006)
Ethiopia	145 (udder)	29%	Abera <i>et al</i> (2010)
Ethiopia	47 (udder)	76.60%	Abera <i>et al</i> (2016)
Ethiopia	543 (quarter)	63%	Bekele and Molla (2001)
Ethiopia	34 (udder)	5.88 %	Hadush <i>et al</i> (2008)
Ethiopia	374 (quarter)	8.9%	Alebie <i>et al</i> (2021)
Iran	243 (95)	18.1%	Niasari-Naslaji <i>et al</i> (2016)
Jordan	90 (udder)	21%	Hawari and Hassawi (2008)
Kenya	86 (quarter)	81.4%	Guliye <i>et al</i> (2002)
Kenya	107 (udder)	66%	Matofari <i>et al</i> (2013)
Kenya	66 (udder)	Most of examined quarter milk samples	Odongo <i>et al</i> (2016)
Kenya	380 (quarter)	44.5%	Toroitich <i>et al</i> (2017)
Kenya	207 (udder)	23%	Younan <i>et al</i> (2001)
Kenya	384 (udder)	61.2%	Wanjohi <i>et al</i> (2013)
Kuwait	25 (udder)	36%	Rahmeh <i>et al</i> 2022
Saudi Arabia	120 (quarter)	Most of examined quarter milk samples	Saleh and Faye (2011)
Saudi Arabia	740 (quarter)	33 % of tested quarters had subclinical mastitis based on CMT	Al Jumaah <i>et al</i> (2012)
Sudan	160 (udder)	71.9%	Elhaj <i>et al</i> (2014)
Sudan	391 (quarter)	43.5%	Abdurahman <i>et al</i> (1995)

after suckling no cleaning of the udder before milking is done (Noor *et al*, 2013). Several lines of evidence suggest that the risk factors for camel mastitis were severe tick infestation, teat injuries, poor milking hygiene and physiological disorders (Ahmad *et al*, 2012; Aqib *et al*, 2022). Finally, the negative effects of mastitis on camel owners include a reduction in milk production, deterioration in milk quality, a decrease in milk price due to high SCC, loss of milk due to antibiotic treatment, and increased animal care costs (Hertl *et al*, 2010; Tuteja *et al*, 2013).

**Using a milking machine to obtain safe raw camel milk for consumption.**

Usually, camels are milked by hand in most countries of the world in traditional farming systems (Bekele *et al*, 2002; Alhadrami and Faye, 2016; Atigui *et al*, 2023). The introduction of machine milking makes only slow progress and is limited to intensive dairy camel farms in a few countries (Kaskous and Fadlelmoula, 2014). To improve the quality and the safety of raw camel milk, machine milking must be used instead of hand milking (Hammadi *et al*, 2010; Nagy and Juhasz, 2016; Kaskous, 2018a, 2019). Saleh *et al* (2013) showed a clear difference between two milking methods in terms of udder health and milk quality. Therefore, microbiological contamination was higher in farms with hand milking than in farms with machine milking (Table 3).

**Table 3.** Bacteriological finding of camel milk samples in two farms with different milking methods (Saleh *et al*, 2013).

Parameters	Farm with machine Milking	Farm with hand milking
Number of Camels	14	14
Duration of the investigation	6 months after calving	6 months after calving
Total samples	84 (100%)	84 (100%)
Uninfected samples	65 (77.4%)	53 (63.1)
Coagulase-negative Staphylococci	15 (17.8%)	22 (26.2%)
<i>Staphylococcus aureus</i>	-	3 (3.6%)
Micrococcus	4 (4.8%)	6 (7.1%)

However, it must be noted that many milking machines used do not succeed in completely emptying the udder. The amount of residual milk after machine milking was high (up to 30% or even more) (Kaskous, 2018b). Review by Ayadi *et al* (2013) clearly showed that low fat levels were observed in milk during mechanical milking,

indicating incomplete milk letdown. In addition, it has also been found that improper use of the milking machine, especially improper use of liners, can damage the camel’s udder, lead to oedema, and promote the colonisation of *Staphylococcus aureus* during the time of machine milking (Juhasz and Nagy, 2008). Results from Tunisia have shown, that the use of machine milking in the field was associated with higher milk yields, but also resulted in increased microbial contamination compared to hand milking (Atigui *et al*, 2023). Kaskous (2023) has shown reverse results that no pathogenic bacteria were detected in the milk produced after using the milking machine (StimuLactor). Improper use of the milking machine is known to have negative effects on udder health as the udder cannot be completely emptied (Kaskous and Pfaffl, 2023). As a result, the remaining milk after milking can serve as a substrate for pathogens and increase the risk of mastitis (Bruckmaier and Wellnitz, 2008). Therefore, completely emptying the udder during milking promotes milk synthesis and secretion and ensures that the udder remains healthy.


As the market is looking for a suitable milking machine for camels, a special modern milking machine called “StimuLactor” was developed by Siliconform, Germany in 2018, which has been used in practice since then (Fig 2). This milking machine for camels was necessary to enable rapid and complete milk extraction during milking and to maintain udder health. A field study showed that the udder remained healthy and completely emptied after using the “StimuLactor” milking machine (Kaskous, 2019), since this milking machine has been adapted to the anatomical, morphological, and physiological characteristics of the camel’s udder.

Training on the milking machine is an important point because by getting the camels used to machine milking, their stress behaviour is reduced, their milk production is increased and the milking time is normalised (Brahmi *et al*, 2024).


Finally, maintaining healthy udders and teats during milking is a key component of an effective milking machine to achieve good and higher quality milk production by preventing mastitis and maintaining animal welfare. In this way, we can help provide consumers with good and safe camel milk.

**Factors affecting safe raw camel milk after harvest**

Raw camel milk is a natural food that can be contaminated with microbiota in the chain from the milking to the consumer as the milk is very good



**Fig 1.** Representative electropherogram of bovine and dromedary camel milk samples determined by capillary electrophoresis according to Mohamed *et al* (2020).



**Fig 2.** A: A StimuLactor milking machine during milking in camel farm, B+C: Attach milk teat cups.

suitable liquid for these microbiotas (Mohammed *et al*, 2016; Kaskous, 2018a). Therefore, a great deal of research has been done to determine the prevalence of microbial content in the raw camel milk after harvesting from the healthy udder (Wanjohi *et al*, 2013; Matofari *et al*, 2013; Odongo *et al*, 2016; Serda *et al*, 2018; Ismaili *et al*, 2019). Autochthonous microbiota from the exterior of the camel's udder and teats can contribute to the contamination as well as microbiota that are derived from the environment in which the camel is housed and milked (Bachmann, 1992; Bekele and Molla, 2001; Hawari and Hassawi, 2008; Omer and Eltinay, 2008; Wanjohi *et al*, 2013; Atigui *et al*, 2023). However, most important it appears to be the contribution of microbiota from teats soiled with manure, mud, and feed. Teats and udders of camels inevitably become contaminated while they are lying or when allowed in dirty lots. The influence of dirty camels on total bacteria counts

depends on the extent of soiling of the teat surface and the udder prep procedures employed. Such microbiota could be reduced if hygiene measures were implemented during milking. However, many milkers or camel owners did not adhere to the hygiene measures during milking (Hassen *et al*, 2022). This means that the milk can become contaminated after it leaves the streak canal. Therefore, due to the traditional nature of camel milk production and the lack of appropriate hygiene measures, there is a high risk of milk contamination with microbiota in the milk production sector in many developing countries (Getachew, 2003). The factors that increase microbiota contamination are mainly due to the high ambient temperatures coupled with a lack of on-farm refrigeration, long distances to markets and a lack of transportation options (Husein *et al*, 2016). Matofari *et al* (2013) reported that camel milk is less contaminated at farm because it has not undergone



many handlers. The only contamination at this stage may come from the infected udder mostly caused by the cocci group. Abera *et al* (2016) reported that the two dominant factors of the quality of camel raw milk after harvesting are the condition of keeping the product and the time before delivery to the consumer. In any case, a high number of microbiotas in the collected milk samples is an indication of unhygienic milk production conditions (Abdurahman, 2006; Kamal *et al*, 2010). It is noteworthy that many farmers store their raw camel milk in plastic canisters after harvest (Hassen *et al*, 2022). An interesting aspect was that 66% of raw camel milk samples at the farm (Production area) had a microbial load less than  $10^5$  cfu/ml, compared to 54% at the collection point (bulk tank) and marketing where the microbial load was above  $10^6$  cfu/ml (Matofari *et al*, 2013). In addition, bicycles, donkeys, and existing vehicles were used to transport raw camel milk from production areas in Kenya to collection or market centres 10 to 20 km away. The ambient temperature in the production areas and on the transport, route was around 39 °C. The raw camel milk reaches the collection points in 2 to 3 hours and the large markets in the cities in 6 to 8 hours (Matofari *et al*, 2013). Through these transportation processes, the raw camel milk could maintain millions of microbiotas, and when this raw camel milk is consumed, the health situation will be severely affected. In this context, Husein *et al* (2016) reported that consuming raw camel milk from production areas was less dangerous than consuming raw camel milk from the market. Furthermore, Abera *et al* (2016) found that about 85.7% of raw camel milk samples in Somali Regional State of Ethiopia were bacterially contaminated and the total bacterial count (TBC) of the contaminated raw camel milk samples was on average  $4.75 \pm 0.17$  log cfu/ml. These bacteria multiply rapidly from the udder to the market. These results indicate that there was a lack of hygiene in the production and milking areas, during transport, and during sale of raw camel milk. A new study in Nigeria has shown that milk samples were taken from five healthy dromedaries and the microbiota *Escherichia coli*, *Salmonella* in the first camel and *Staphylococcus aureus* in the remaining camels were found (Dogondaji *et al*, 2023). The authors emphasise that the consumption of raw camel milk should therefore, be done with caution. Kaskous (2019) found that by adhering to hygienic measures, good, safe raw camel milk with normal composition was obtained. Therefore, hygiene measures must also be observed after the harvest in order to maintain the quality standard of raw camel milk.

## Conclusion

- Regarding the importance of camel milk and the associated related health benefits of its bioactive ingredients, improvements in milking hygiene, milk storage and transport conditions are required to ensure the quality of camel milk meets consumer needs.
- Camels must be kept healthy and raw milk must be refrigerated immediately after milking and during storage and transport.
- The used milking system should adapt to the morphological, anatomical, and physiological characteristics of the udder and teat of the lactating camels and it should achieve a physiologically ideal milking process that meets high animal welfare standards and increases milk production with a high-quality standard.
- To obtain safe camel milk for human consumption, it is recommended to consume pasteurised camel milk.

## References

- Abbas S, Ashraf H, Nazir DA and Sarfraz DL. Physiochemical analysis, and composition of camel milk. International Research Journal. 2013; 2:83-98.
- Abdel Gader AGM and Alhaider AA. The unique medicinal properties of camel products: A review of the scientific evidence. Journal of Taibah University Medical Sciences. 2016; 11(2):98-103.
- Abdel Gadir Atif E, Hildebrandt G, Kleer JN, Molla B, Kyule MN and Baumann MP. Comparison of California Mastitis Test (CMT), somatic cell counts (SCC) and bacteriological examinations for detection of camel (*Camelus dromedarius*) mastitis in Ethiopia. Berliner und Münchener Tierärztliche Wochenschrift. 2006; 119(1-2):45-49.
- Abduku H and Eshetu M. Physico-chemical properties and processing characteristics of camel milk as compared with other dairy species: A review. Asian Journal of Dairy and Food Research. 2024; 43(1):1-7.
- Abdurahman OA, Agab H, Abbas B and Aström G. Relations between udder infection and somatic cells in camel (*Camelus dromedarius*) milk. Acta Veterinaria Scandinavica. 1995; 36(4):423-431.
- Abdurahman OA. Udder health and milk quality among camels in the Error Valley of eastern Ethiopia. Livestock Research for Rural Development. 2006; 18: Article 110.
- Abera M, Abdi O, Abunna F and Megersa B. Udder health problems and major bacterial causes of camel mastitis in Jijiga, eastern Ethiopia: implication for impacting food security. Tropical Animal Health and Production. 2010; 42:341-347.
- Abera T, Legesse Y, Mummed B and Urga B. Bacteriological quality of raw camel milk along the market value chain in Fafen zone, Ethiopian Somali regional state. BMC Research Notes. 2016; 9:285-290.

- Abu-Lehia IH. Physical and chemical characteristics of camel milk fat and its fraction. *Food Chemistry*. 1989; 34(4):261-267.
- Ahmad S, Yaqoob M, Bilal MQ, Khan MK, Muhammad G, Yang LG and Tariq M. Factors affecting yield and composition of camel milk kept under desert conditions of central Punjab, Pakistan. *Tropical Animal Health and Production*. 2012; 44:1403-1410.
- Al haj A and Al Kanhal HA. Compositional, technological, and nutritional aspects of dromedary camel milk. *International Dairy Journal*. 2010; 20(12):811-821.
- Alamin MA, Alqurashi AM, Elsheikh AS and Yasin TE. Mastitis incidence and bacterial causative agents isolated from lactating she-camel (*Camelus dromedarius*). *IOSR Journal of Agriculture Veterinary Science*. 2013; 2:7-10.
- Alaoui Ismaili M, Saidi B, Zahar M, Hamama A and Ezzaier R. Composition and microbial quality of raw camel milk produced in Morocco. *Journal of the Saudi Society of Agricultural Sciences*. 2019; 18:17-21.
- Alebie A, Molla A, Adugna W, Tesfaye A and Ejo M. Prevalence, isolation, identification, and risk factors of major bacterial cause of camel subclinical mastitis. *BioMed Research International*. 2021; 5522331.
- Alhadrami GA and Faye B. Animals that produce dairy foods: Camel. Reference Module in Food Science. Elsevier, 1<sup>st</sup> Ed. 2016; pp 1-12.
- Alhaj OA, Lajnaf R, Jrad Z, Alshuniaber MA, Jahrami HA and Serag El-Din MF. Comparison of ethanol stability and chemical composition of camel milk from five samples. *Animals*. 2022; 12(5):1-11.
- Ali W, Akyol E, Ceyhan A, Dilawar S, Firdous A, Qasim MZ and Ahmad MM. Milk production and composition in camel and its beneficial uses: A review. *Turkish Journal of Agriculture - Food Science and Technology*. 2019; 7(12):2142-2147.
- Aljumaah RS, Almutairi FF, Ismail E, Alshaikh MA, Sami A and Ayadi M. Effects of production system, breed, parity, and stage of lactation on milk composition of dromedary camels in Saudi Arabia. *Journal of Animal and Veterinary Advances*. 2012; 11:141-147.
- Almaw G and Molla B. Prevalence and aetiology of mastitis in camels (*Camelus dromedarius*) in Iraq. *Journal of Camel Practice and Research*. 2000; 7:97-100.
- Aqib AI, Muzammil I, Naseer MA, Shoaib M, Bakht P, Zaheer T, Khan YR, Khan RL, Usman M, Shafeeq M, Tanveer Q, Hussain HI, Saleem A and Prince K. Pathological insights into camel mastitis. *Acta Tropica*. 2022; 231: 106415.
- Arab HH, Salama SA, Eid AH, Omar HA, Arafa E-SA and Maghrabi IA. Camel's milk ameliorates TNBS-induced colitis in rats via downregulation of inflammatory cytokines and oxidative stress. *Food and Chemical Toxicology*. 2014; 69:294-302.
- Asfour HAE and Anwer AM. Some bacteriological and immunological studies on camel's milk. *Alexandria Journal of Veterinary Sciences*. 2015; 47:38-46.
- Atigui M, Fguiri I, Arroum S, Brahmi M, Ghzaïel B and Hammadi M. Effect of milking routines and hygiene practices and evolution along the market value chain on raw camel milk quality in Tunisia. *Italian Journal of Animal Science*. 2023; 22:337-346.
- Ayadi M, Aljumaah RS, Musaad A, Samara EM, Abdelrahman MM, Alshaikh MA, Saleh SK and Faye B. Relationship between udder morphology traits, alveolar and cisternal milk compartments and machine milking performance of dairy camels (*Camelus dromedarius*). *Spanish Journal of Agriculture Research*. 2013; 11(3):790-797.
- Bachmann T. Quality control at reception. In: *Handbook on Milk Collection in Warm Developing Countries*. IDF, Brussels. 1992; pp 1-12.
- Behrouz S, Saadat S, Memarzia A, Sarir H, Folkerts G and Boskabady MH. The antioxidant, anti-inflammatory and immunomodulatory effects of camel milk. *Front. Immunology*. 2022; 13:855342.
- Bekele T and Molla B. Mastitis in lactating camels (*Camelus dromedarius*) in Afar Region, north-eastern Ethiopia. *Berliner und Münchener Tierärztliche Wochenschrift*. 2001; 114(5-6):169-172.
- Bekele T, Lunddeheim N and Dahlborn K. Milk production and feeding behaviour in the camel (*Camelus dromedarius*) during 4 watering regimens. *Journal of Dairy Science*. 2011; 94(3):1310-1317.
- Bekele T, Zeleke M and Baars RMT. Milk production performance of the one humped camel (*Camelus dromedarius*) under pastoral management in semi-arid eastern Ethiopia. *Livestock Production Science*. 2002; 76:37-44.
- Bhatt L, Chahar A, Tuteja FC and Verma D. Prevalence aetiology and antibiogram of subclinical mastitis isolates from camel. *Veterinary Practitioner*. 2004; 5:61-65.
- Boudalia S, Gueroui Y, Zebba R, Arbia T, Chiheb AE, Benada M, Hadri Z, Youcefi A and Bousbia A. Camel livestock in the Algerian Sahara under the context of climate change: Milk properties and livestock production practices. *Journal of Agriculture and Food Research*. 2023; 11:100528.
- Boujenane I. Review of milk let-down in camels and proposition of a milk recording method. *Tropical Animal Health and Production*. 2020; 52(6):2845-2853.
- Brahmi M, Atigui M, Hammadi I, Seddik M-M, Ben Salem W, Marnet P-G and Hammadi M. Effect of camel's previous experience on behavioural response and milk production during training to machine milking. *Applied Animal Behaviour Science*. 2024; 270:106118.
- Bruckmaier R and Wellnitz O. Induction of milk ejection and milk removal in different production systems. *Journal of Animal Science*. 2008; 86:15-20.
- Carruth L, Roess AA, Mekonnen YT, Hosh FM and Salman MD. Antimicrobial resistance and food safety in Africa. *The Lancet Infectious Diseases*. 2017; 7(6):575-576.
- Carruth L. Camel milk, amoxicillin, and a prayer: medical pluralism and medical humanitarian aid in the Somali region of Ethiopia. *Social Science and Medicine*. 2014; 120:405-412.

- Chamekh L, Khorchani T, Dbara M, Hammadi M and Yahyaoui MH. Factors affecting milk yield and composition of Tunisian camels (*Camelus dromedarius*) over complete lactation. *Tropical Animal Health and Production*. 2020; 52:3187-3194.
- Cekh Ismail L, Osaili TM, Mohamad MN, Zakaria H, Ali A, Tarek A, Ashfaq A, Al Abdouli MA, Saleh ST, Daour RA, Alrajaby R, Stojanovska L and Al Dhaheiri AS. Camel milk consumption patterns and perceptions in the UAE: A cross-sectional study. *Journal of Nutritional Science*. 2022; 21(11):e59
- Dadar M, Shahali Y and Whatmore AM. Human brucellosis caused by raw dairy products: a review on the occurrence, major risk factors and prevention. *International Journal of Food Microbiology*. 2019; 292:39-47.
- Davati N, Yazdi FT, Zibae S, Shahidi F and Edalatian MR. Study of lactic acid bacteria community from raw milk of Iranian one humped camel and evaluation of their probiotic properties. *Jundishapur Journal of Microbiology*. 2015; 8(5):1-6.
- De Buyser ML, Dufour B, Maire M and Lafarge V. Implication of milk and milk products in food-borne disease in France and different industrialised countries. *International Journal of Food Microbiology*. 2001; 67:1-17.
- Devendra K, Verma KA, Chatli MK, Singh R, Kumar P, Mehta N and Malav OP. Camels milk: Alternative milk for human consumption and its health benefits. *Nutrition and Food Science*. 2016; 46:217-227.
- Dogondaji RA, Lawal M, Wasagu RSU, Yakubu A and Umar S. Microbial assessment of raw milk of some female camels (*Camelus dromedarius*) in Sokoto, Nigeria. *PSM Microbiology*. 2023; 8(1):1-8.
- Dugassa D. Quality and therapeutic aspect of camel milk: A review. *Journal of Food Processing and Technology*. 2021; 12(8):Nr. 902.
- El Tigani-Asil ETA, Abdelwahab GE, Veedu JTVP, Khalafalla AI, Mohamed ZSA, Ishag HZA, Shah AAM, Alhosani MAA and Muhairi SSM. Gangrenous mastitis in dromedary camels in UAE caused by *Streptococcus agalactiae*. *BMC Veterinary Research*. 2020; 16:174-178.
- Elagamy EI, Nawar M, Shamsia SM, Awad S and Haenlein GF. Are camel milk proteins convenient to the nutrition of cow milk allergic children? *Small Ruminant Research*. 2009; 82:1-6.
- Elhaj AE, Freigoun, Somaya AB and Mohamed TT. Aerobic bacteria and fungi associated with raw camel's milk. *Online Journal of Animal and Feed Research*. 2014; 4(1):15-17.
- El-Hatmi H, Girardet J-M, Gaillard J-L, Yahyaoui MH and Attia H. Characterisation of whey proteins of camel (*Camelus dromedarius*) milk and colostrum. *Small Ruminant Research*. 2007; 70:267-271.
- El-Hatmi H, Jrad Z, Salhi I, Aguiabi A, Nadri A and Khorchani T. Comparison of composition and whey protein fractions of human, camel, donkey, goat, and cow's milk. *Mljekarstvo/Dairy*. 2015; 65:159-167.
- El-Hatmi H, Levieux A and Levieux D. Camel (*Camelus dromedarius*) immunoglobulin G,  $\alpha$ -lactalbumin, serum albumin and lactoferrin in colostrum and milk during the early post-partum period. *Journal of Dairy Research*. 2006; 73(3):288-293.
- Elmoslemany AM, Keefe GP, Dohoo IR, Wichtel JJ, Stryhn H and Dingwell RT. The association between bulk tank milk analysis for raw milk quality and on-farm management practices. *Preventive Veterinary Medicine*. 2010; 95:32-40.
- El-Zeini HM. Microstructure, rheological and geometrical properties of fat globules of milk from different animal species. *Polish Journal of Food and Nutrition Sciences*. 2006; 56(2):147-154.
- El-Ziney MG. Microbiological quality and safety assessment of camel milk (*Camelus dromedarius*) in Saudi Arabia (Qassim region). *Applied Ecology and Environmental Research*. 2007; 5:115-122.
- Eyassu S. Handling, preservation and utilization of camel milk and camel milk products in Shinile and Jigjiga Zones, Eastern Ethiopia. *Livestock Research for Rural Development*. 2007; 19(6): Article 86.
- FAO stat. World Food and Agriculture-Statistical Yearbook. Rome, Italy, 2022; pp 184-188.
- Farah Z, Rettenmaier R and Atkins D. Vitamin content of camel milk. *International Journal for Vitamin and Nutrition Research*. 1992; 62(1): 30-33.
- Farah Z. Camel milk. In: Fuquay JW, Fox PF, McSweeney PLH, *Encyclopaedia of Dairy Sciences*. 2<sup>nd</sup> Ed., London, UK: Academic Press, 2011; pp 512-517.
- Faraz A. Composition of camel milk: A Blessing for Health. *Annals of Public Health and Epidemiology*. 2020; 1(2): APHE.MS.ID.000509.
- Frister H. Zusammensetzung der Milch, In: Krömker, V. *Kurzes Lehrbuch Milchkunde und Milchhygiene*. Parey, MVS Medizinverlag Stuttgart GmbH. 2007; pp 80-101.
- Getachew F. Milk and dairy products, post-harvest losses and food safety in sub-Saharan Africa and the near east. *Assessments Reports in the dairy sub sector in Ethiopia*. Action programme for the prevention of food losses. FAO, Rom. 2003; pp 1-60.
- Gizachew A, Teha J, Birhanu T and Nekemete E. Review on medicinal and nutritional values of camel milk. *Natural Science*. 2014; 12(12):35-41.
- Gonfa A, Foster HA and Holzapfel WH. Field survey and literature review on traditional fermented milk products of Ethiopia. *International Journal of Food Microbiology*. 2001; 68:173-186.
- Guliye AY, Van Creveld C and Yagil R. Detection of subclinical mastitis in dromedary camels (*Camelus dromedarius*) using somatic cell counts and the N-acety-beta-D-glucosaminidase test. *Tropical Animal Health and Production*. 2002; 34(2):95-104.
- Haddadin MS, Gammoh SI and Robinson RK. Seasonal variation in the chemical composition of camel milk in Jordan. *Journal of Dairy Research*. 2008; 75(1):8-12.
- Hadush B, Kebede E and Kidanu H. Assessment of bacteriological quality of raw camels' milk in Ab-Ala,



- north eastern Ethiopia. Livestock Research for Rural Development. 2008; 20(9):Article 151.
- Hamed H, Trujillo A-J, Juan B, Guamis B, Elfeki A and Gargouri A. Interrelationships between somatic cell count, lactation stage and lactation number and their influence on plasmin activity and protein fraction distribution in dromedary (*Camelus dromedarius*) and cow milk. Small Ruminant Research. 2012; 105:300-307.
- Hammadi M, Atigue M, Ayadi M, Barmat A, Belgacem A, Khaldi G and Khorchani T. Training and short time effects of machine milking on milk yield and milk composition in Tunisian Maghrebi camels (*Camelus dromedarius*). Journal of Camel Practice and Research. 2010; 17(1):1-7.
- Hassen M, Amentie T, Abdimahad K, Maalin A and Mahamed A. Hygienic production, and post-harvest handling practices of raw camel milk in Degahbour District of Jarar Zone, Somali Regional State, Ethiopia. Open Journal of Animal Sciences. 2022; 12:303-316.
- Hawari AD and Hassawi DS. Mastitis in one humped she-camels (*Camelus dromedarius*) in Jordan. Journal of Biological Science. 2008; 8:958-961.
- Hertl JA, Grohn Yrjo, Leach JdG, Bar D, Bennett GJ, Gonzalez RN, Rauch BJ, Welcome FL, Tauer LW and Schukken YH. Effects of clinical mastitis caused by gram-positive and gram-negative bacteria and other organisms on the probability of conception in New York State Holstein dairy cows. Journal of Dairy Science. 2010; 93:1551-1560.
- Husein M, Selamawit H, Ashenafi G, Fikre Z and Ashenafi F. Assessment on safety status of camel raw milk marketed in samara-logia town of afar National regional state, northeast Ethiopia. Food Science and Quality Management. 2016; 49:80-88.
- Ismaili MA, Saidi B, Zahar M, Hamama A and Ezzaier R. Composition, and microbial quality of raw camel milk produced in Morocco. Journal of the Saudi Society of Agricultural Science. 2019; 18:17-21.
- Izadi A, Khedmat L and Mojtahedi SY. Nutritional and therapeutic perspectives of camel's milk and its protein hydrolysates: a review on versatile bio functional properties. Journal of Functional Foods. 2019; 60(2):103441.
- Jilo K and Tegegne D. Chemical composition and medicinal values of camel milk. International Journal of Research in BioSciences. 2016; 4(4):13-25.
- Johnson B, Joseph M, Jose Sh, Jose S, Kinne J and Wernery U. The microflora of teat canals and udder cisterns in nonlactating dromedaries. Journal of Camel Practice and Research. 2015; 22(1):55-59.
- Jrad Z, Oulahal N, Adt I, Khorchani T, Degraeve P and El-Hatmi H. Camel colostrum: Nutritional composition and improvement of the antimicrobial activity after enzymatic hydrolysis. Emirates Journal of Food and Agriculture. 2015; 27(4):384-389.
- Juhasz J and Nagy P. Challenges in the development of a large-scale milking system for dromedary camels. In: Nagy P, Huszenicza G, and Juhasz J (Eds.) WBC/ ICAR Satellite Meeting on Camelid Reproduction, Budapest, Hungary. 2008; pp 1-4
- Kamal M and Karoui R. Monitoring of mild heat treatment of camel milk by front-face fluorescence spectroscopy. LWT-Food Science and Technology. 2017; 79:586-593.
- Kamal ZL, Kamal ZA and Heydar NA. Total bacteria, coliforms and the *Staphylococcus aureus* bacteria count of raw milk (from farms to the processing factory), and pasteurised milk in Khuzestan province. The 4th Congress on Animal Science, Tehran, Iran. 2010; pp 21-18.
- Kanca H. Milk production and composition in ruminants under heat stress. In: Watson RR, Collier RJ, Preedy VR, Nutrients in dairy and their implications for health and disease. London, UK, Elsevier, 2017; pp 97-109.
- Kappeler S, Farah Z and Puhan Z. 5-Flanking regions of camel milk genes are highly similar to homologue regions of other species and can be divided into two distinct groups. Journal of Dairy Science. 2003; 86:498-508.
- Karaman A, Akgül FY, Ogut S, Canbay HS and Alvarez V. Gross composition of raw camel's milk produced in Turkey. Food Sci. Technol. Campinas. 2022; 42:e59820.
- Kaskous S and Fadlelmoula AA. The challenge of machine milking in dromedary camel. Scientific Journal of Review. 2014; 3(2):1004-1017.
- Kaskous S and Pfaffl M. Bioactive properties of minor camel milk ingredients- an Overview. Journal of Camel Practice and Research. 2017; 24(1):15-26
- Kaskous S and Pfaffl MW. Milking machine settings and liner design are important to improve milking efficiency and lactating animal welfare-Technical note. Agri Engineering. 2023; 5:1314-1326.
- Kaskous S, Alasaad A, Nouh A, Mohamed U, Sauerwein H and Bruckmaier RM. The concentration of lactoferrin and other milk constituents in Syrian Shami camel during different lactation seasons. Journal of Agriculture Sciences, Damascus University. 2012; 28(2):273-287.
- Kaskous S, Al-Momani AQ, Al-Yacoub AN and Al-Najjar KA. Physiological perspective of milk somatic cell counts in lactating camels. Journal of Camel Practice and Research. 2021; 28(3):319-325.
- Kaskous S. A quarter individual milking machine "Stimu Lactor" in a camel farm in Switzerland: According to field Study. Journal of Camel Practice and Research. 2023; 30(3):267-271.
- Kaskous S. Camel milk composition, udder health and effect of different storage times and temperatures on raw milk quality using camel milking machine "StimuLactor." Agriculture and Food Sciences Research. 2019; 6(2):172-181.
- Kaskous S. Importance of camel milk for human health. Emirates Journal of Food and Agriculture. 2016; 28(3):158-163.
- Kaskous S. Physiology of lactation and machine milking in dromedary she-camel. Emirates Journal of Food and Agriculture. 2018b; 30(4):295-303.
- Kaskous S. Sources of contamination of raw camel milk with microorganisms. Conference "Tropentag" Global food security and food safety: The Role of Universities. Ghent, Belgium, September 17-19, 2018a, pp 251.

- Khalesi M, Salami M, Moslehishad M, Winterburn J and Moosave-Movahedi AA. Biomolecular content of camel milk: A traditional superfood towards future healthcare industry. *Trends in Food Science and Technology*. 2017; 62:49-58.
- Khan BB and Iqbal A. Production and composition of camel milk-review. *Pakistan Journal of Agricultural Sciences*. 2001; 38(3-4):64-68.
- Khaskheli M, Arain MA, Chaudhry S, Soomro AH and Qureshi TA. Physioco- chemical quality of camel milk. *Journal of Agriculture and Social Science*. 2005; 2:164-166.
- Konuspayeva G, Faye B and Loiseau G. The composition of camel milk: A meta-analysis of the literature data. *Journal of Food Composition and Analysis*. 2009; 22:95-101.
- Konuspayeva G, Faye B, Loiseau G and Levieux D. Lactoferrin and Immunoglobulin contents in camels' milk (*Camelus bactrianus*, *Camelus dromedarius*, and hybrids) from Kazakhstan. *Journal of Dairy Science*. 2007; 90:38-46.
- Konuspayeva G, Faye B, Pauw ED and Focant J-F. Level and trends of PCDD/Fs and PCBs in camel milk (*Camelus bactrianus* and *Camelus dromedarius*) from Kazakhstan. *Chemosphere*. 2011; 85:351-360.
- Konuspayeva G, Lemarie E, Faye B, Loiseau G and Montet D. Fatty acid and cholesterol composition of camels (*Camelus bactrianus*, *Camelus dromedarius* and hybrids) milk in Kazakhstan. *Dairy Science and Technology*. 2008; 88(3):327-340.
- Kraimia M, Adamou A, Boudjenah-Haroun S and Benaissa MH. Factors influencing the physicochemical and mineral composition of camel milk in eastern Algeria. *Journal of Camel Practice and Research*. 2024; 31(1):17-24.
- Kula JT and Dechasa T. Chemical composition and medicinal values of camel milk. *International Journal of Research in Biosciences*. 2016; 4:13-25.
- Kumar D, Verma AK, Chatli MK, Singh R, Kumar P, Mehta N and Malav OP. Camel milk: alternative milk for human consumption and its health benefits. *Nutrition and Food Science*. 2016; 46(2):217-227.
- Lajnaf R, Attia H and Ayadi MA. Technological properties, and biological activities of camel  $\alpha$ -lactalbumin-A review. *International Dairy Journal*. 2023; 139:105563.
- Mal G, Dande S and Sahani MS. Milk production potential and keeping quality of camel milk. *Journal of Camel Practice and Research*. 2006; 13(2):175-178.
- Matofari JW, Shalo PL, Younan M, Nanua JN, Adonogo A and Qabale Misiko BN Analysis of microbial quality and safety of camel (*Camelus dromedarius*) milk chain and implications in Kenya. *Journal of Agricultural Extension and Rural Development*. 2013; 5(3):50-54.
- Matofari JW, Shitandi A, Shalo PL, Nanua NJ and Younan M. A Survey of *Salmonella* enteric contamination of camel milk in Kenya. *African Journal of Microbiological Resources*. 2007; 1(4):46-50.
- Mbye M, Ayyash M, Abu-Jdayil B and Kamal-Eldin A. The texture of camel milk cheese: effect of milk composition, coagulants, and processing conditions. *Front. Nutr*. 2022; 9:868320.
- Medhammar E, Wijesinha-Bettoni R, Stadlmayr B, Nilsson E, Charrondiere UR and Burlingame B. Composition of milk from minor dairy animals and buffalo breeds: A biodiversity perspective. *Journal of the Science of Food and Agriculture*. 2012; 92:445-474.
- Mehaia MA, Hablas MA, Abdel-Rahman KM and El-Mougy SA. Milk composition of Majaheim Wadah and Hamra camels in Saudi Arabia. *Food Chemistry*. 1995; 52(2):115-122.
- Mehaia MA. Vitamin C and riboflavin content in camel's milk: effects of heat treatments. *Food Chemistry*. 1994; 50(2):153-155.
- Mehari Y, Mekuriaw Z and Gebru G. Potentials of camel production in Babilie and Kebribeyah woredas of the Jijiga Zone, Somali Region, Ethiopia. *Livestock Research for Rural Development*. 2007; 19(4):Article 58.
- Mohamed H, Johansson M, Lundh A, Nagy P and Kamal-Eldin A. Short communication: Caseins and  $\alpha$ -lactalbumin content of camel milk (*Camelus dromedarius*) determined by capillary electrophoresis. *Journal of Dairy Science* 2020; 103:11094-11099.
- Mohammed H, Hailu S, Geberegiorgis A, Zeru F and Feyisa A. Assessment of safety status of camel raw milk marketed in Samara-Logia Town of Afar national regional state, Northeast Ethiopia. *Food Science and Quality Management*. 2016; 49:80-88.
- Mulwa WD, Schelling E, Wangoh J, Imungi KJ, Farah Z and Meile L. Microbiological quality of raw camel milk across the Kenyan market chain. *Global Science Book*. 2011; 5(1):79-83.
- Musinga M, Kimenye D and Kivilonzi P. The camel milk industry in Kenya: Results of a study commissioned by SNV to explore the potential of camel milk from Isiolo District to access formal markets. *Netherlands Development Organization/Resource Mobilisation Centre, Kenya*. 2008; 100:43-48.
- Mwangi LW, Matofari JW, Muliro PS and Bebe BO. Hygienic assessment of spontaneously fermented raw camel milk (suusa) along the informal value chain in Kenya. *International Journal of Food Contamination*. 2016; 3:18-26.
- Nagy P and Juhasz J. Review of present knowledge on machine milking and intensive milk production in dromedary camels and future challenges. *Tropical Animal Health and Production*. 2016; 48(5):915-926.
- Niasari-Naslaji A, Pezeshk H, Atakpour AB, Ghaffari S, Nickchi P, Safi S, Shirazi-Beheshtiha SH, Arabha H, Samiei R, Amjadi M, Haji Moradlou AA, Narimani I and Moosavi-Movahedi AA Estimation of somatic cell count, as gold standard to detect subclinical mastitis, in dromedary camel. *Journal of Camel Practice and Research*. 2016; 23(1):175-178.
- Nikkhah A. Equidae, camel and yak milks as functional foods: A review. *Journal of Nutrition and Food Sciences*. 2011; 1:116.
- Noor IM, Guliye AY, Tariq M and Bebe BO. Assessment of camel and camel milk marketing practices in an emerging periurban production system in Isiolo county,

- Kenya. Pastoralism Research Policy and Practice. 2013; 3:28-35.
- Obeid AI, Bagadi HO and Mukhtar MM. Mastitis in *Camelus dromedarius* and the somatic cell content of camel's milk. Research in Veterinary Science. 1996; 61(1):55-58.
- Odongo NO, Lamuka PO, Matofari JW and Abong GO. Risk factors associated with the post-harvest loss of milk along camel milk value chain in Isiolo County, Kenya. African Journal of Agricultural Research 2016; 11(8):674-682.
- Omer RH and Eltinay AH. Microbial quality of camel's raw milk in central and southern regions of United Arab Emirates. Emirates Journal of Food and Agriculture. 2008; 20(1):76-83.
- Omrani AS, Al-Tawfiq JA and Memish ZA. Middle east respiratory syndrome coronavirus (MERS-CoV): animal to human interaction. Pathogens and Global Health. 2015; 109(8):354-362.
- Oselu S, Ebere R and Arimi JM. Camels, camel milk, and camel milk product situation in Kenya in relation to the world. Journal of Food Science. 2022; 8:1237423.
- Park YW and Haenlein GFW. Milk from other minor species (Reindeer, Caribou, Musk Ox, Llama, Alpaca, Moose, Elk and others), chapter 30. In: Milk and Dairy Products in Human Nutrition. Wiley-Blackwell Publishers, Oxford, UK. 2013; pp 644-658.
- Patel AS, Patel SJ, Patel NR and Chaudhary GV. Importance of camel milk-An alternative dairy food. Journal of Livestock Science. 2016; 7:19-25.
- Rahmeh R, Akbar A, Alomirah H, Kishk M, Al-Ateeqi A, Shajan A, Alonaizi T and Esposito A. Assessment of mastitis in camel milk using high-throughput sequencing. PLOS One. 2022; 717(12):e0278456.
- Rahmeh R, Alomirah H, Akbar A and Sidhu J. Composition and properties of camel milk, In: Javed K. Milk production, processing and marketing. IntechOpen. 2019; 82592, pp 1-17.
- Redington JM, Breydo L, Almeshdar HA, Redwan EM and Uversky VN.  $\alpha$ -Lactalbumin: of camels and cows. Protein and Peptide Letters. 2016; 23(12):1072-1080.
- Roess AA, Hosh FM, Morton LC, Bestul N, Davis J and Carruth L. Associations between unpasteurised camel and other milk consumption, livestock ownership, and self-reported febrile and gastrointestinal symptoms among semi-pastoralists and pastoralists in the Somali region of Ethiopia. Epidemiology and Infection 2023; 151:e44, 1-7.
- Roy D, Ye A, Moughan PJ and Singh H. Composition, structure, and digestive dynamics of milk from different species-A review. Frontiers in Nutrition. 2020; 7:577759.
- Sadler K and Catley A. Milk matters: The role and value of milk in the diets of somali pastoralist children in Liben and Shinile, Ethiopia. Feinstein International Centre, Tufts University and Save the Children, Addis Ababa. Report. 2009; pp 1-28
- Saleh SK and Faye B. Detection of subclinical mastitis in dromedary camels (*Camelus dromedarius*) using somatic cell counts, California mastitis test and udder pathogen. Emirates Journal of Food and Agriculture. 2011; 23(1):48-58.
- Saleh SK, Al-Ramadhan G and Faye B. Monitoring of monthly SCC in she-camel in relation to milking practice, udder status and microbiological contamination of milk. Emirates Journal of Food and Agriculture. 2013; 25 (5):403-408.
- Sara B, Elhassan MBM and Zubeir IEME. Evaluation of pasteurisation and sterilisation process on camel milk quality. Global Journal of Science Frontier Research 2022; 22(2):1-11.
- Sawaya WN, Khalil JK, Al-Shalhat A and Al-Mohammad H. Chemical composition and nutritional quality of camel milk. Journal of Food Science. 1984; 49:744-747.
- Schulz J. Bau der Milchdrüse und Physiologie der Laktation. In: Fahr R-D and von Lengergen G. Milcherzeugung. Deutscher Fachverlag. 2003; pp 45-79.
- Seifu E and Tafesse B. Prevalence and aetiology of mastitis in traditionally managed camels (*Camelus dromedarius*) in selected pastoral areas in eastern Ethiopia. Ethiopian Veterinary Journal. 2010; 14:103-113.
- Seifu E. Handling, preservation and utilisation of camel milk and camel milk products in Shinile and Jijiga zones, eastern Ethiopia. Livestock Research for Rural Development 2007; 19(6):Article 86.
- Seifu E. Recent advances on camel milk: Nutritional and health benefits and processing implications-A review. AIMS Agriculture and Food. 2022; 7(4):777-804.
- Serda B, Bekele A and Abebe D. Prevalence and contamination level of *Staphylococcus aureus* in raw camel milk and associated risk factors in Jijiga District, eastern Ethiopia. Journal of Veterinary Science and Technology. 2018; 9(1):501-505.
- Shamsia S. Nutritional and therapeutic properties of camel and human milks. International Journal of Genetics and Molecular Biology. 2009; 1(4):52-58.
- Shuiep ES, El Zubeir IEM, El Owani OAO and Musa HH. Influence of season and management on composition of raw camel milk (*Camelus dromedarius*) milk in Khartoum state, Sudan. Tropical and Subtropical Agroecosystems. 2008; 8(1):101-106.
- Singh R, Ghorui S and Sahani M. Camel milk: properties and processing potential. In: Sahani MS, The Indian camel. NRCC, Bikaner. 2006; pp 59-73.
- Smits M, Huppertz T, Alting A and Kiers J. Composition, constituents, and properties of Dutch camel milk. Journal of Camel Practice and Research. 2011; 18(1):1-6.
- Smits M, Joosten H, Faye B and Burger PA. The flourishing camel milk market and concerns about animal welfare and legislation. Animals (Basel). 2023; 13(47):1-16.
- Spargue LD, Al-Dahouk S and Neubauer H. A review on camel brucellosis: a zoonosis sustained by ignorance and indifference. Pathogens and Global Health. 2012; 106:144-149.
- Stahl T, Sallmann H-P, Duehlmeier R and Wernery U. Selected vitamins and fatty acid patterns in dromedary milk and colostrum. Journal of Camel Practice and Research. 2006; 13(1):53-57.



- Stahl T. Vitamingehalte und Fettsäuremuster in Kamelmilch. PhD thesis, Tierärztliche Hochschule Hannover, Hannover 2005; pp 1-130.
- Swelum AA, El-Saadony MT, Abdelnour SA, Ba-Awadh H, Abd El-Halck ME and Sheiha AM. Relationship between concentration of macro and trace elements in serum and follicular, oviductal, and uterine fluids of the dromedary camel (*Camelus dromedarius*). Tropical Animal Health and Production. 2020; 52:1315-1324.
- Swelum AA, El-Saadony MT, Abdo M, Ombarak RA, Hussein EOS, Suliman G, Alhimaidei AR, Ammari AA, Ba-Awadh H, Taha AE, El-Tarabily KA and Abd El-Halck ME. Nutritional, antimicrobial and medicinal properties of camel's milk: A review. Saudi Journal of Biological Sciences. 2021; 28(5):3126-3136.
- Swinburne M. Raw milk: Camels, the FDA, and State deregulation. In: The Network for Public Health Law-Eastern Region. Report. 2017; February 13 ([www.networkforphl.org](http://www.networkforphl.org)).
- Toroitich KC, Gitau GK, Kitala PM and Gitao GC. The prevalence and causes of mastitis in lactating traditionally managed one-humped camels (*Camelus dromedarius*) in West Pokot County, Kenya. Livestock Research for Rural Development. 2017; 29(4):Article 62.
- Tuteja F, Patil N, Narnaware S, Nagarajan G and Dahiya S. Primarily human pathogenic fungi causing dermatophytosis in camel. Journal of Camel Practice and Research. 2013; 20(2):151-155.
- Vincenzetti S, Cammertoni N, Rapaccetti R, Santini G, Klimanova Y, Zhang J and Polidori P. Nutraceutical and functional properties of camelid's milk. Beverages. 2022; 8(1):1-12.
- Wang SY, Liang JP, Shao WJ and Wen H. Mineral, vitamin, and fatty acid contents in the camel milk of dromedaries in the Anxigansu China. Journal of Camel Practice and Research. 2011; 18(2):273-276.
- Wanjohi M, Gitao CG and Bebora L. Subclinical mastitis affecting hygienic quality of marketed camel milk from Northeastern Province, Kenya. Microbiology Research International. 2013; 1(1):6-15.
- Wernery U, Johnson B and Jose SH. The most important dromedary mastitis organisms. Journal of Camel Practice and Research. 2008; 15(2):159-161.
- Wernery U, Soellner NK, Joseph S, Varghese P, Johnson B and Kinne J. Passive immunisation against brucella melitensis in dromedaries. Journal of Camel Practice and Research. 2017; 24(3):235-237.
- Wernery U. Camel milk, the white gold of the desert. Journal of Camel Practice and Research. 2006; 13:15-26.
- Wernery U. Camelid brucellosis: a review. Rev. Science. Tech. 2014; 33(3):839-857.
- Wernery U. Camel milk-new observations. Proceedings of the International Camel Conference, "Recent trends in camelids research and future strategies for saving camels". Rajasthan, India, 16-17 February 2007; 200-204.
- Wijesinha-Bettoni R and Burlingame B. Milk and dairy product composition. In: Milk and Dairy Products in Human Nutrition. FAO Rome, Wiley-Blackwell. 2013; pp 41-102.
- Yadav AK, Kumar R, Priyadarshini L and Singh J. Composition and medicinal properties of camel milk: a review. Asian Journal of Dairy and Food Research 2015; 34(2): 83-91.
- Yamina M, Wassila C, Kenza Z, Amina Z, Noureddine S, Eddine HJ and Mebrouk K. Physico-chemical and microbiological analysis of Algerian raw camels' milk and identification of predominating thermophilic lactic acid bacteria. Journal of Food Science and Engineering. 2013; 3:55-63.
- Yirda A, Eshetu M and Babege K. Current status of camel dairy processing and technologies: a review. Open Journal of Animal Sciences. 2020; 10:362-377.
- Younan M, Ali Z, Bornstein S and Müller W. Application of the California Mastitis Test in intramammary *Streptococcus agalactiae* and *Staphylococcus aureus* infections of camels (*Camelus dromedarius*) in Kenya. Preventive Veterinary Medicine. 2001; 51:307-316.
- Zaitlin P, Dwyer J and Gleason GR. Mistaken beliefs and the facts about milk and dairy foods. Nutrition Today. 2013; 48:135-143.
- Zangerl P. Mikrobiologie der Produkte. In: Krömker V. Kurzes Lehrbuch Milchkunde und Milchhygiene, Parey. 2007; pp 156-179.
- Zimmermann KA. Camel milk: Nutrition Facts, Risks and Benefits. Published February 3, 2016 ([www.livescience.com/53579-camel-milk](http://www.livescience.com/53579-camel-milk)).

## SAUDI EXHIBITION IN GENEVA SHOWCASES CAMEL CONVERSION PRODUCTS' SIGNIFICANCE IN FOOD SECURITY

The Saudi Exhibition at the International Year of Camelids (IYC) 2024 was inaugurated in Geneva, Switzerland from June 25 to 28. Dr. Ali Al-Shaikhi, undersecretary for fisheries and livestock wealth at the Ministry of Environment, Water, and Agriculture, inaugurated the event. The event was held in partnership with the State of Bolivia at the United Nations Palace in Geneva. The event was attended by United Nations Human Rights Council Vice President Mayra Mariela Macdonal Alvarez, UN Food and Agriculture Organisation (FAO) Director General Qu Dongyu, Saudi Ambassador to Switzerland Dr. Adel Merdad, and several ministers, ambassadors, and senior officials representing about 100 UN member states.

At least 13 government and private entities participated in the Saudi exhibition, showcasing the most important locally manufactured camel conversion products and their significance in achieving food security. The exhibition also included leather, medical, and cosmetic products derived from camels. It sought to embody the great cultural value of camels in Saudi society. The Kingdom's participation in the event stems from its presidency of the International Year of Camels 2024 in partnership with the Group of Latin American and the Caribbean (GRULAC), represented by Bolivia. Saudi Arabia has taken important steps to develop the camel sector, including raising public awareness about the importance of camels and their role in providing food and livelihood for their owners and breeders, encouraging increased productivity, combating diseases and epidemics that afflict them, vaccinating them periodically, and improving the marketing of camel products to raise the income of breeders and thus maximise and raise their living standards. Saudi Arabia emphasised the significance of camels in promoting economic development, contributing to food security, and achieving sustainable development goals for livestock, all in line with the adoption of several initiatives to support the camel sector, which represents a cultural heritage and a long-standing tradition.

(Source: Staff Writer, Saudi Gazette, June 26, 2024 RELATED TOPICS FOOD SECURITY ENVIRONMENT SAUDI ARABIA SUSTAINABILITY)

## CAMEL CRAZY: A QUEST FOR MIRACLES IN THE MYSTERIOUS WORLD OF CAMELS: MONGOLIAN EDITION



The award-winning book CAMEL CRAZY: A Quest for Miracles in the Mysterious World of Camels, by Christina Adams, gained wide global acclaim in the original English version. It is now available in the Mongolian language from NEPKO Publishing of Ulaanbaatar, Mongolia. The new edition is updated with photos of Mongolian camel herders, the camel dairy farm of Solid Partners Group, and a new table on the nutritional content of Bactrian camel milk, as well as information on the endangered wild camel species found in Mongolia.. A "page turning odyssey of a mother on a mission-from Bedouin camps in the Middle East to Amish farms in Pennsylvania to camel-herder villages in India--to obtain camel milk, which dramatically helps her son's autism symptoms. Adams' work uncovers studies of camel milk for possible treatment of autism, allergies, diabetes, and immune dysfunction, as well as ancient traditions of healing. But the most fascinating aspect of Adams' discoveries is the gentle-eyed,

mischievous camels themselves." Contact Bayarmagni Ganbold of NEPKO Publishing for copies at phone 976-8810 1714 or email gbayarmagnaid@gmail.com. Website: www.nepoko.mn.

# MORPHOLOGY AND FUNCTION OF THE SPLEEN IN CAMEL: A COMPARATIVE REVIEW

A.M. Marwa-Babiker<sup>1,2</sup> and Nawal M. Elkhair<sup>3,4</sup>

<sup>1</sup>Department of Anatomy, <sup>3</sup>Department of Biomedical Sciences, College of Veterinary Medicine, King Faisal University, P.O. Box 400, Al-Ahsa 31982, Saudi Arabia

<sup>2</sup>Department of Anatomy, College of Veterinary Medicine, University of Bahri, Khartoum-North, Sudan, P.O. Box 012, Khartoum North 12217 Sudan

<sup>4</sup>Department of Physiology, Faculty of Veterinary Medicine, University of Khartoum, 13314, Shambat, Sudan

## ABSTRACT

The morphology of the spleen varies among different mammalian species. In camels, it is C-shaped, while in dogs, it is elongated, dumbbell shaped, and ventrally oversized. Ruminants have a flat and oblong spleen, while in horses it is comma shaped. The spleen is located in the left upper quadrant of the caudal peritoneal cavity, behind the diaphragm, and in close proximity to the colon, kidney, lower ribs, stomach, and pancreas in camels and other mammalian species, respectively. The camel's spleen is enclosed within a thick capsule. This capsule is divided into an inner layer composed of smooth muscle cells and an outer layer primarily made of connective tissue. The two layers of the capsule are smooth muscle and dense irregular connective tissue. Species differences exist in both the overall thickness and relative quantity of smooth muscle. Collagen and elastic fibres make up trabeculae, while smooth muscle cells extend into the parenchyma from the capsule and the hilus. The spleen in camels plays a crucial role in the storage and regulation of blood supply. The central artery, which emerged from the periarterial lymphatic sheath, was branched to four penicilli and then extended as sheathed arterioles, which were found near or surrounded by blood sinusoids of the red pulp. The white pulp is surrounded by a broad marginal zone that lacks marginal sinusoids but has sheathed arteries. The marginal zone macrophages, a key player in the spleen's immune response, were characterised by its large size and intimate connections between its cell processes and the nearby marginal zone B cells. Scanning electron microscopy of the camel foetus's spleen demonstrated the primary structure, including mesothelium, numerous erythrocytes, medium-sized and small lymphocytes, reticular cells, and reticular connective tissue fibres. Differential features of spleen of other animals are also discussed.

**Key words:** Anatomy, camel, function, histology, spleen, ultrastructure

The spleen is an intriguing organ that performs various functions such as haematopoiesis, immunological response, blood filtration, and blood storage, and these functions vary among different animal species. As the animal develops, there are exciting changes in the presence and significance of each function, with a reduction in haematopoietic activities and an increase in the specialisation of other tasks over time (Udroiu and Sgura, 2017).

The spleen, an intraperitoneal lymphoid organ, is situated on the left side of the abdomen, below the diaphragm (Dyce *et al*, 1987). During gestation, the spleen plays a significant role in haematopoiesis, (Boes and Durham, 2017). Post-birth, the spleen carries out a range of vital physiological functions, including haematopoiesis and lymphopoiesis (lymphocyte proliferation and maturation), blood filtration via recycling of senescent and damaged erythrocytes, erythrocyte and thrombocyte storage,

iron metabolism, and immune surveillance (Cheung and Nadakavukaren, 1983; Mebius and Kraal, 2005; Cesta, 2006; Eurell and Frappier, 2006; Moura *et al*, 2008; Khalil *et al*, 2009; Bello *et al*, 2019; Gnanadevi *et al*, 2019; Lewis *et al*, 2019; Xu *et al*, 2020; Abdellatif, 2021).

The spleen's structure reflects its complexity. It is primarily composed of red and white pulp (RP and WP), with a distinct marginal zone between them (Bronte and Pittet, 2013). The WP, despite occupying less than a quarter of the spleen's tissue, is its principal immunologic zone. The RP makes up the majority of the tissue and serves a different immunological function than the WP. The absence of afferent lymphatic arteries in the spleen contributes to its unique structure, as all cells and antigens enter the spleen via the blood (Lewis *et al*, 2019).

This shift from scattered lymphocytes to the segregation of the WP indicates a rise in the complexity of immunological characteristics and the

SEND REPRINT REQUEST TO A.M. MARWA-BABIKER [email: mbabakr@kfu.edu.sa](mailto:mbabakr@kfu.edu.sa), [marwa.babiker22@gmail.com](mailto:marwa.babiker22@gmail.com)



organisation of T and B lymphocyte regions, leading to the formation of germinal centres. Additionally, different features of the RP have evolved in mammalian species, enhancing the specialisation of the filtration function of the pulp sinuses and the storage function (Udroiu and Sgura, 2017).

In adult animals, lymphopoiesis is the spleen's main haematopoietic activity, while erythropoiesis occurs during foetal life. However, splenic erythropoiesis persists in newborn horses and ruminants for several weeks postpartum (Boes and Durham, 2017). Erythrocytes and thrombocytes are typically stored in the RP and splenic cords. As animals age, the elasticity of the erythrocyte cell membrane decreases, leading to the detection and removal of old and damaged erythrocytes by the splenic cords' reticuloendothelial system and macrophages in the red pulp's venous sinuses. A reticular fibre network of macrophages and reticular cells across the reticuloendothelial system helps the spleen filter blood more effectively, resulting in numerous macrophages containing engulfed erythrocyte fragments and hemosiderin pigment in the RP (Eurell and Frappier, 2006).

### Gross anatomy of the spleen in camel

The morphology of the spleen varies among different mammalian species (Fig 1). In camels, it is C-shaped, while in dogs, it is elongated, dumbbell-shaped, and ventrally oversized. Ruminants have a flat and oblong spleen, while in horses it is comma-shaped. According to Smuts and Bezuidenhout (1987) and Dyce *et al* (1987), the spleen is located in the left upper quadrant of the caudal peritoneal cavity, behind the diaphragm, and in close proximity to the colon, kidney, lower ribs, stomach, and pancreas in camels and other mammalian species, respectively. The spleen is typically not noticeable on physical inspection, but in thin adults, children, and adolescents may be detected (Nguyen and Zhang, 2020).

Accessory spleens, also known as supernumerary spleens, splenules, or splenunculi, are benign and asymptomatic. They are estimated to be present in 10% to 30% of the population, with an individual potentially having one to six accessory splenic buds. Accessory spleens are typically only a few millimetres long, but can occasionally grow to a few centimetres in length, resulting in variation in size (Nguyen and Zhang, 2020; Yildiz *et al*, 2013; Mohammadi *et al*, 2016).

Khalel (2010) conducted a comprehensive study on the gross anatomy of the spleen in Awasi


sheep, unearthing unique findings that significantly advanced our understanding in the field. The spleen, for instance, was found to possess a distinctive triangular shape with rounded corners, weighing approximately  $69\pm6.7$  g and measuring an average length and width of  $9.93\pm0.3$  and  $6.48\pm0.2$  cm, respectively. It reached its maximum thickness near the hilus ( $2.48\pm0.1$  cm). The spleen also exhibited two distinct surfaces: the visceral, which was concave and had a hilus, and the parietal, which appeared to be convex. In sheep, the spleen had two ends: the ventral end, which was narrower and thinner than the base end, and the base end, which was broad and thick. These detailed observations, along with other intriguing findings such as the quadrangular shape of goat spleens (Suri *et al*, 2017; Gnanadevi *et al*, 2019) have significantly enriched our understanding of the gross anatomy of the spleen in various animal species.

The spleen, a vital organ in camels, was characterised by its dark brown colour and unique C-shape with blunt, rough surfaces and edges. It was thick at the hilus and the midline and thin at the edges (Maina *et al*, 2014). According to Nawal and Maher (2018), the spleen of a camel had four segments, two hilus, and two splenic arteries and veins. The same authors stated that the splenic blood supply had only one arterial segmentation with no anastomosis. In buffalo calves, sheep, and goats, only one tiny hilus was found, and anastomosis and venous segmentation between the dorsal and intermediate segments were absent. According to Marwa-babiker *et al* (2023), the spleen has C-shaped, rough surfaces and serrated edges in camel fetuses. It is located on the caudolateral side of the abdominal cavity, medially to the ribs and caudally to the stomach. These findings underscore the practical importance of understanding the gross anatomy of the spleen in different animal species, making a significant contribution to the field of veterinary medicine and anatomy.


The spleen of the Asian elephant was dark reddish, long and narrow. The caudal-visceral surface was concave and displayed a longitudinal hilus towards its cranial border; the cranial-parietal surface was convex with serrations on the caudal and cranial splenic borders (Rajani *et al*, 2021).

### Histology of mammals' spleen

Zidan *et al* (2000) found that the camel's spleen is enclosed within a thick capsule. This capsule is divided into an inner layer composed of smooth muscle cells and an outer layer primarily made of connective tissue. The same authors showed that




**Fig 1.** Photograph showing the spleen of different domestic mammalian species, S: stomach, L: gastro splenic ligament.




**Fig 2.** Photomicrograph showing general histological view of the mammalian (rat) spleen: stroma of the spleen consists of connective tissue capsule surround the spleen (C), which sends the connective tissue trabeculae (T). The parenchyma consists of white pulp (WP) and red pulp (RP).

the vascular and avascular trabeculae extending from the capsule to the parenchyma, subcapsular, and per trabecular blood sinuses around primary and vascular trabeculae are unique structures of the camel spleen. These unique structures are not found



**Fig 3.** a and b: Photomicrograph showing the spleen; C: connective tissue capsule, T: connective tissue trabeculae, GC: germinal centre of the splenic corpuscle (dotted circle). MZ: marginal zone, which found between the white pulp and the red pulp (RP). Central arteriole (arrow) and splenic sinuses (arrows head).



**Fig 4.** a and b: Scanning electron micrograph of spleen of camel foetus showing mesothelium (M) erythrocytes (white arrows), lymphocytes (red arrows) reticular cells (arrowhead) and reticular connective tissue fibres (blue arrows).



in other mammalian species, indicating a distinct evolutionary path for the camel spleen. In other mammalian species, the spleen is surrounded by a capsule of thick connective tissue invested by the peritoneum (Eurell and Frappier, 2006) (Figs 2 and 3).

The two layers of the capsule are smooth muscle and dense irregular connective tissue. Species differences exist in both the overall thickness and relative quantity of smooth muscle (Eurell and Frappier, 2006 and Marwa-babiker *et al*, 2023). Collagen and elastic fibres make up trabeculae, while smooth muscle cells extend into the parenchyma from the capsule and the hilus.

The spleen, a vital component of the body's immune system, is a complex organ. It is composed of RP, responsible for storing red blood cells, and WP, rich in lymphocytes and active in immune responses (Eurell and Frappier, 2006; Bloom and Fawcett, 1986). The RP is made up of cell cords and sinusoids, while WP is lymphoid in nature, with a marginal zone containing pores where cells can leak (Van Krieken, 1997). The WP houses two types of blood vessels: one with muscled walls, and the other with thinner walls sheathed by reticular cells. Within the reticular cells, a network of lymphoblast, small and medium lymphocytes, and dendritic macrophages is formed. This intricate network of cells and vessels is crucial to the body's defense system, underscoring the importance of studying the unique camel spleen.

According to Zidan *et al* (2000), the spleen in camels plays a crucial role in the storage and regulation of blood supply. The central artery, which emerged from the periarterial lymphatic sheath, was branched to four penicilli and then extended as sheathed arterioles, which were found near or surrounded by blood sinusoids of the RP. The WP is surrounded by a broad marginal zone that lacks marginal sinuses but has sheathed arteries. The RP was divided into cords by secondary trabeculae and contained numerous sizes of venous sinusoids. The observation demonstrated that the spleen of the camel was a sinusal type that can store blood, as well as the thick muscular capsule and trabeculae pump according to the body's need. Closed and open circulations facilitated the blood supply of the spleen. A unique venous return was drained from the venous sinusoids of the RP to the peritrabecular sinuses and subcapsular sinuses to the splenic vein. These histological structures showed no significant differences related to age.

During the first trimester of pregnancy, the spleen's capsule in camels was made of a thin connective tissue, indicating a relatively low blood

storage capacity. However, as the pregnancy progressed into the second and third trimesters, the spleen underwent significant changes. The capsule transformed into a thick, dense, irregular connective tissue with descending trabeculae of collagen fibres and bundles of smooth muscle fibres, suggesting an increased blood storage capacity. The parenchyma, which consisted of a variety of cell types in the first trimester, differentiated into white and red pulps in the second and third trimesters. The RP contained megakaryocytes in all three trimesters, indicating a continuous production of blood cells throughout the pregnancy (Marwa-Babiker *et al*, 2022).

The marginal zone macrophages, a key player in the spleen's immune response, were characterised by its large size and intimate connections between its cell processes and the nearby marginal zone B cells. It also boasts a highly developed phagocytic potential (Dijkstra *et al*, 1985). In mice, Humphrey and Grennan (1981) noted that the marginal zone macrophages were larger than the other macrophages in the spleen and had a distinct morphology. These unique characteristics of the marginal zone macrophages contribute to our understanding of the spleen's immune system and its role in maintaining the camel's health.

In a meticulous study of rats, the authors observed that the spleen, before differentiated into the periarteriolar lymphoid sheath in the interior layer of the marginal zone (MZ), migrating lymphocytes tended to gather in some places near marginal-zone macrophages (Dijkstra, 1982; Brelinska and Pilgrim, 1983). This careful observation underscores the importance of understanding the spleen's structure in different animal species.

In rats (Figs 2 and 3), in line with Suri *et al* (2017) and Gnanadevi *et al* (2019) in goats, has revealed that the spleen was enclosed in a thick capsule of irregular dense connective tissue, and red and white pulps. The WP, a crucial component, is composed of periarterial lymphatic sheath and splenic nodule, which are scattered throughout the RP. The RP, another key element, consists of splenic cords and sinusoids; the endothelial cells with prominent nuclei that protruded into the sinusoidal lumen bordered the sinusoids (Gnanadevi *et al*, 2019 and Suri *et al*, 2017).

Rahman *et al* (2016) studied the histomorphometry of spleen in human, goat, buffalo, rabbit and rat and found significant implications. Identical capsular thickness measurements were observed in goats and buffaloes, whereas the thin and thinnest capsule was measured in rat and rabbit, respectively.



In sheep, the capsule was made of collagen, elastic, reticular, and smooth muscle fibres (Thanvi *et al*, 2020). The red and white pulps, the main components of the splenic parenchyma, play a crucial role in the spleen's function. The WP, consisting of lymphoreticular tissue, lymphatic nodule and periarterial lymphatic sheath, is less abundant. The RP, made up of pulp arterioles, splenic sinusoids, splenic cords, and sheathed and terminal capillaries, is also less abundant and poorly developed (Thanvi *et al*, 2020).

A novel finding conducted by Rajani *et al* (2021), described that the spleen of the Asian elephant is surrounded by a thick capsule of dense irregular connective tissue. Most of the splenic parenchyma, a unique feature, consisted of RP, while the remaining 20.25% was WP, composed of the periarterial lymphatic sheath and splenic nodules (Rajani *et al*, 2021). Many investigators have observed numerous megakaryocytes in the RP of adult and foetal dromedary camel (Zidan *et al*, 2000; Marwa-Babiker *et al*, 2022), whereas it was observed in both the spleen and bone marrow of pregnant rats (Marien and McFadden, 1970). The findings are significant in understanding the spleen's structure in different animal species.

### Ultrastructure of spleen

Scanning electron microscopy of the camel foetus's spleen demonstrated the primary structure, including mesothelium, numerous erythrocytes, medium-sized and small lymphocytes, reticular cells, and reticular connective tissue fibres (Fig 4). During the first trimester of pregnancy, a comparatively limited number of reticular cells were observed in the parietal sheath, marginal zones, and cordal gaps. Dendritic macrophages and lymphoblasts were organised into a network by the reticular cells throughout the second and third trimesters. Marwa-Babiker *et al* (2023) investigated the spleen of a camel foetus and found that it was lined with mesothelial cells and divided by reticular connective tissue fibres using scanning electron microscopy.

Weiss (1974) studied the scanning electron microscopy of normal rat spleen and demonstrated that the reticulum and vascular sinuses were made of macrophages, erythrocytes, thrombocytes, and other migratory elements. Large, bulky, irregular reticular cells with broad processes were found in the periarterial lymphatic sheath, the marginal zone and cordal spaces. Flattened reticular cells at the periphery of the periarterial lymphatic sheath formed cylinder shapes attached to the central artery and associated with unusually heavy extracellular fibres.

Vascular sinuses were suspended in the reticulum by attachments of cordal reticular cells to the adventitial surface. Adventitial cells of the sinus were branched into the cords. Endothelial cells were typically laid side by side without gaps, except as migratory cells which passed through the wall. The erythrocytes were observed in passage across the sinus wall. Sinuses and cords were often swollen, irregular, and bore blebs. Macrophages displayed rich surface folds and processes, whereas the thrombocytes were abundant and adherent to the endothelium of sinuses and the surface of reticular cells.

Polak *et al* (2009) demonstrated that the splenic cords originated from a three-dimensional network of fibroblastic reticular cells situated among branched sinuses and artery terminals in human spleen. A unique variety of rod-shaped endothelial cells that were oriented perpendicular to the longitudinal axis of the sinuses lined their interior walls. The fibroblastic reticular cells were transformed into fixed phagocytes with no phagocytosis properties. The capsule and trabeculae were composed of random smooth muscle fibres.

The reticular cells in the spleen were three-dimensional stellate shapes with smooth surfaces that extended slender processes. Although the reticular cells they exposed in some places, the reticular fibres were typically covered by reticular cell processes (Fujita and Ushiki, 1992).

Electron microscopy of albino rabbit's spleen revealed that the white pulp was arranged in clumps or nests separated by a loose-irregular meshwork composed of basement membrane-like material, collagen, and the cellular prolongations of fibroblasts and stellate cells, lymphocytes, and numerous reticular cells. The plasma cells, macrophages, erythrocytes, and thrombocytes were situated at the periphery of the white pulp and adjacent to the marginal zone. The lymphocytes of the white pulp varied in size and was difficult to differentiate from the polymorphic reticular cells (Burke and Simon, 1970).

According to Kashimura and Fujita (1987), the central arteries in human were continuous and without branching into follicular arteries in the white pulp, while developing into penicillar arteries in the red pulp. Although some penicillar arteries retreated in the marginal zone, others travelled through it, penetrating the white pulp and developing into follicular arteries.

In humans and rats, the spleen was examined using scanning electron microscopy (Sasou *et al*, 1986). Both species had the marginal zone and white and red

pulps; however, there were considerable differences in the artery termination in the marginal zone. In rats, the follicular arteries formed the marginal sinus, a circulatory net that terminated at the edge of the white pulp. Numerous vascular termini of the follicular and sheathing arteries were dispersed throughout the human's marginal zone. Despite these differences, both in humans and rats, the central artery was found to be surrounded by flat reticular cells, a reassuring similarity. In the red pulp of rats, the vascular termini were funnel- or tubular-shaped. The sheath of human arteries had a circumferential lamellar structure consisting of flat reticular fibres (Sasou *et al*, 1986).

## Conclusion

The spleen, a vital organ in various mammals, exhibits significant variations in shape. For instance, it was bigger ventrally and bell-shaped in dogs, comma-shaped in horses, C-shaped in camels, and elongated in ruminants. The celiac plexus provides precise innervation and blood supply via the splenic artery and vein. This intricate network ensures the organ's optimal functioning. The spleen is a soft, reddish-purple organ with two surfaces, visceral and diaphragmatic, and its limits are divided into three areas: cranial, caudal, and intermediate. The splenic hilum, located on the caudomedial side of the gastric impression, contains nerves, splenic vessels, splenorenal, and gastrosplenic ligaments and is enclosed in an elastic and fibrous tissue capsule that extends into the parenchyma as trabeculae. The parenchyma, supported by an exemplary network of reticular fibres, comprises two types of tissue: white pulp and red pulp, separated by marginal zones.

## Acknowledgements

The authors would like to acknowledge the 1st level Veterinary Medicine female students 2022-2023; Department of Anatomy, College of Veterinary Medicine, King Faisal University for providing gross anatomy image used in this review paper.

## Conflict of Interest

The authors declare that they have no conflict of interest.

## Data Availability Statement

The data presented in this study are available on request from the corresponding author.

## References

Abdellatif AM. Structure of the Eurasian moorhen spleen: A comprehensive study using gross anatomy, light, and

transmission electron microscopy. *Microscopy Research and Technique*. 2021; 84(8):1696-1709. <https://doi.org/10.1002/jemt.23728>.

- Bello A, Onu JE, Onyeansi BI, Sonfada ML, Umaru MA, Danmaigoro A. and Hena SA. Unique anatomy of the spleen of one humped camel (*Camelus dromedarius*): A review. *Open Access Journal of Biomedical Engineering and Biosciences*. 2019; 3(3):294-298.
- Bloom W and Fawcett DW. *A Textbook of Histology*. 11th edition, WB Saunders, Philadelphia and London. 1986.
- Boes KM and Durham AC. *Bone Marrow, Blood Cells, and the Lymphoid/Lymphatic System. Pathologic Basis of Veterinary Disease*. 2017; 724-804.e2. doi: 10.1016/B978-0-323-35775-3.00013-8.
- Brelinska R and Pilgrim C. Macrophages and interdigitating cells; their relationship to migrating lymphocytes in the white pulp of rat spleen. *Cell Tissue and Research*. 1983; 233:671-688.
- Bronte V and Pittet MJ. The spleen in local and systemic regulation of immunity. *Immunity*. 2013; 14; 39(5):806-18. doi: 10.1016/j.immuni.2013.10.010.
- Burke JS and Simon GT. *Electron Microscopy of the Spleen 1. Anatomy and Microcirculation*. *American Journal of Pathology*. 1970; pp 127-154. <https://pubmed.ncbi.nlm.nih.gov/5414015>.
- Cesta MF. Normal Structure, Function, and Histology of the Spleen. *Toxicologic Pathology*. 2006; 34:455-465. <https://doi.org/10.1080/01926230600867743>.
- Cheung HT and Nadakavukaren MJ. Age-dependent changes in the cellularity and ultrastructure of the spleen of Fischer F344 rats. *Mechanisms of Ageing and Development*. 1983; 22:23-33. [https://doi.org/10.1016/0047-6374\(83\)90004-0](https://doi.org/10.1016/0047-6374(83)90004-0).
- Dijkstra CD. Characterisation of non-lymphoid cells in rat spleen, with special reference to strongly I a positive branched cells in the T-cell areas. *J Reticuloendothelial Soc*. 1982; 32:167-178. PMID: 6296385.
- Dijkstra CD, van Vliet E, Dopp EA, van der Lelij AA. and Kraal G. Marginal zone macrophages identified by a monoclonal antibody: characterisation of immuno- and enzyme-histochemical properties and functional capacities. *Immunology*. 1985; 55:23-30. <http://www.ncbi.nlm.nih.gov/pmc/articles/pmc1453576>.
- Dyce KM, Sack WO And Wensing CJG. *Textbook of Veterinary Anatomy*. W.B. Saunders Company, Philadelphia, London, Toronto. 1987.
- Eurell JA and Frappier BL. *Dellmann's Text Book of Veterinary Histology*. Blackwell publishing. 6<sup>th</sup> edition. 2006; pp 147-150.
- Fujita T and Ushiki T. Scanning electron microscopic observations of the immunodefensive systems with special reference to the surface morphology of the non-lymphoid cells. *Archives of Histology and Cytology*. 1992; 55:105-13. [https://doi.org/10.1679/aohc.55.suppl\\_105](https://doi.org/10.1679/aohc.55.suppl_105).
- Gnanadevi R, Senthilkumar S, Kannan TA. and Ramesh G. Comparative Histoarchitectural study of splenic components in sheep and goat. *International Journal*

- of Current Microbiology and Applied Sciences. 2019; 8(5):1387-1394. <https://doi.org/10.20546/ijcmas.2019.805.158>.
- Humphrey JH and Grennan D. Different macrophage populations distinguished by means of fluorescent polysaccharides. Recognition and properties of marginal zone macrophages. The European Journal of Immunology. 1981; 11:221-228. <https://doi.org/10.1002/eji.1830110311>.
- Kashimura M and Fujita T. Scanning electron microscopy study of human spleen: relationship between the microcirculation and functions. Scanning Electron Microscope. 1987; 1(2):841-851. PMID: 3616578.
- Khalel EM. Anatomical and histological study of the spleen in Iraqi sheep (Awasi sheep). Basrah Journal of Veterinary Research. 2010; 10(2):163-171.
- Khalil M, Sultana SZ, Rahman M, Mannan S, Ahmed S, Ara ZG. and Chowdhury AI. Study of prenatal and postnatal development of spleen of *Gallus domesticus* (Deshi chicken). Mymensingh Medical Journal. 2009; 18:169-174. PMID: 19623142.
- Lewis SM, Williams A. and Eisenbarth SC. Structure-function of the immune system in the spleen. Science Immunology 1. 2019; 4(33):10.1126.
- Maina MM, Usende IL, Igwenagu E, Onyiche TE, Yusuf ZM. and Ntung NO. Gross, Histological and Histomorphometric Studies on the Spleen of One Humped Camel (*Camelus dromedarius*) Found in the Semi-Arid Region of North Eastern Nigeria. Journal of Veterinary Advances. 2014; 4(10):703-711. DOI:10.5455/jva.20141025045543.
- Marien GJ and McFadden KD. Splenic megakaryocytes and circulating platelets in pregnant rats. American Journal of Anatomy. 1970; 128(2):225-33. <https://doi.org/10.1002/aja.1001280207>.
- Marwa-Babiker AM, AlKhodair KM, Abdelhay MA, Elseory AMA, Zolain HB, Al-Ramadan SY. and Althnaian TA. Morphology of the spleen in the camel's foetus (*Camelus dromedarius*): gross anatomy, histology and scanning electron microscopic studies. Journal of Applied Veterinary Sciences. (2023); 8(4):54-61. DOI: 10.21608/JAVS.2023.219602.1250.
- Marwa-Babiker AM, Althnaian TA. and Ali AM. Histological study of prenatal development of the spleen in camel (*Camelus dromedarius*). Journal of Camel Practice and Research. 2022; 29(2):251-255. <http://dx.doi.org/10.5958/2277-8934.2022.00035.2>.
- Mebius RE and Kraal G. Structure and function of the spleen. Review. Nature Reviews Immunology. 2005; 5(8): 606-16.
- Mohammadi S, Hedjazi A, Sajjadian M, Ghrobi N, Moghadam MD. and Mohammadi M. Accessory spleen in the splenic hilum: a cadaveric study with clinical significance. Medical Archives. 2016; 70(5):389-391. <https://doi.org/10.5455%2Fmedarh.2016.70.389-391>.
- Moura R, gua-Doce A, Weinmann P, Graça L, Fonseca JE. B cells: from the bench to the clinical practice. Acta Reumatológica Portuguesa. 2008; 33:137-154. <https://www.researchgate.net/publication/5247192>.
- Nawal AN and Maher MA. Gross Anatomical, Radiographic and Ultra-structural identification of splenic vasculature in some ruminants (Camel, Buffalo Calf, Sheep and Goat). International Journal of Advanced Research in Biological Sciences. 2018; 5(2):44-65. <http://dx.doi.org/10.22192/ijarbs.2018.05.02.006>.
- Nguyen L and Zhang L. Anatomy and Histology of the Normal Spleen and Liver. 2020. <https://api.semanticscholar.org/CorpusID:219019314>.
- Polák Š, Gálfiová P. and Varga I. Ultrastructure of human spleen in transmission and scanning electron microscope. Biologia. 2009; 64/2:402-408.
- Rahman N, Tandon R, Ghaus F, Moinuddin A, Akram W. and Faruqi NA. Comparative anatomy of spleen: histomorphometric study in human, goat, buffalo, rabbit and rat. Academia Anatomica International. 2016; 2(1):28-32. <https://doi.org/10.21276/aaanat.2016.2.1.6>.
- Rajani CV, Indu VR, Patki HS, Surjith KP. and Pradeep M. Morphology of spleen in Asian elephant (*Elephas maximus indicus*). Indian Journal of Veterinary Anatomy. 2021; 33 (1):29-31.
- Sasou S, Satodate R, Masuda T. and Takayama K. Scanning electron microscopic features of spleen in the rat and human: a comparative study. Scanning Electron Microscope. 1986; (Pt 3):1063-9. <https://pubmed.ncbi.nlm.nih.gov/3798009>.
- Smuts MMS and Bezuidenhout AJ. Anatomy of the Dromedary. Clarendon Press, Oxford. 1987.
- Suri S, Sasan JS, Kamal Sarma KS. and Dibyendu Chakraborty D. Comparative gross and histomorphological studies on the spleen of sheep and goat of Jammu region of India. Exploratory Animal and Medical Research. 2017; 7(2):179-183.
- Thanvi PK, Joshi S and Singh D. Histomorphological studies on spleen of sheep (*Ovis aries*). Veterinary Practitioner. 2020; 21(1):48-53.
- Udroiu I and Sgura A. The phylogeny of the spleen. The Quarterly Review of Biology. 2017; 92(4):411-443. <http://dx.doi.org/10.1086/695327>.
- Van Krieken JHJM. Histology of the spleen: structure and cellular distribution in the normal spleen. Current Diagnostic Pathology. 1997; 4(2):100-105. [https://doi.org/10.1016/S0968-6053\(05\)80089-4](https://doi.org/10.1016/S0968-6053(05)80089-4).
- Weiss L. A scanning electron microscopic study of the spleen. Blood. 1974; 43(5):665-691.
- Xu M, Li W, Yang S, Sun X, Tarique I. Yang P. and Chen Q. Morphological characterisation of postembryonic development of blood-spleen barrier in duck. Poultry Science. 2020; 99:3823-3830. <https://doi.org/10.1016/j.psj.2020.05.012>.
- Yildiz AE, Ariyurek MO. and Karcaaltincaba M. Splenic anomalies of shape, size, and location: pictorial essay. Scientific World Journal. 2013; 321810. doi: 10.1155/2013/321810
- Zidan M, Kassem A. and Pabst R. Megakaryocytes and platelets in the spleen of the dromedary camel (*Camelus dromedarius*). Anatomia, Histologia, Embryologia. 2000; 29(4):221-4. <https://doi.org/10.1046/j.1439-0264.2000.00268.x>.



## INTERNATIONAL WORKSHOP ON CAMELID PASTORALISM



The International Workshop on Camelid Pastoralism, held in Sadri, Rajasthan February 2024, marked the inauguration of the International Year of Camelids in 2024. Hosted by Lokhit Pashu-Palak Sansthan (LPPS), the League for Pastoral Peoples (LPP), and the Godwar camel pastoralist milk producers, the event gathered stakeholders from across the globe. Participants hailed from regions like Rajasthan, Gujarat, Mongolia, Kenya, Iran, UK, Germany, Peru, Pakistan, UAE, and

USA, representing a diverse array of perspectives and experiences in camelid pastoralism.

The workshop highlighted the indispensable role of camelids in ensuring food security and biodiversity protection, particularly in arid and semi-arid regions worldwide. Participants emphasised the importance of mobile management systems, recognising camelids' resilience and environmental benefits, including their soft-footed grazing habits. Traditional knowledge held by camelid pastoralists, accumulated over centuries, was acknowledged as invaluable in sustaining camelid herds and producing a wide range of camelid-derived products.

Women emerged as leaders in camelid development, driving innovation and marketing efforts for camelid-based products. Their contributions not only enrich their own communities but also support camelid pastoralists globally. However, camelid pastoralists face numerous challenges, including land dispossession, policy neglect, and encroachment by extractive industries and urbanisation.

Rejecting the extractive model inherited from colonial legacies, participants advocated for a fossil-fuel-free approach to camelid development. They called upon governments to recognise camelids' territories of life, respect customary rights, and provide legal, institutional, and financial support. Access to veterinary services and incentivizing youth engagement in mobile camelid pastoralism were deemed essential for ensuring the continuation of traditional practices.

To preserve camelid herding as a cultural heritage and ensure the well-being of pastoralists and their herds, participants stressed the need for fair and ethical value chains. Community-based initiatives that prioritise the pastoralists' benefits and remunerate them fairly for their products were identified as development priorities.

As the International Year of Camelids unfolds, stakeholders are urged to centre their efforts on understanding and valuing the knowledge and work of camelid-herding communities. Collaboration and solidarity among global camelid pastoralists are essential to promote a development model that benefits camelids, people, and the planet, setting the stage for a successful International Year of Rangelands and Pastoralists in 2026.

(Courtesy: Ilse Kohler Röllefson and Hanwant Singh, LPPS, Sadri, India)

# HISTOLOGICAL STRUCTURE OF THE PROSTATE GLAND DURING ANTE AND POSTNATAL PERIOD OF ONTOGENESIS IN DROMEDARY CAMEL IN ALGERIA

Hafiza Aidoudi<sup>1</sup>, Djallal Eddine Rahmoun<sup>2</sup> and Tarek Khenenou<sup>1</sup>

<sup>1</sup>Laboratory of Sciences and Technics of the Livings, <sup>2</sup>Laboratory of Animal Production, Biotechnologies and Health (PABIOS) Institute of Agriculture and Veterinary Sciences, University of Souk Ahras, Taoura 41009, Algeria

## ABSTRACT

The current investigation was aimed to assess morpho-functional (histological) variations in the prostate of clinically healthy dromedaries during both ante- and postnatal phases of ontogeny. Thirty prostates from two distinct age cohorts (ante- and postnatal) were procured from the EL OUAD slaughterhouse (located in the eastern region of ALGERIA) for microscopic examination, with a focus on delineating the microscopic architecture of the prostate in this species. A range of staining methods, including Haematoxylin and Eosin, as well as silver nitrate impregnation, were employed to elucidate the normal histological framework and developmental trajectory of the prostate across different age groups. Subsequent statistical analysis was conducted to interpret the acquired data with significance set at  $P < 0.05$ . Our findings revealed discernible patterns of structural and functional differentiation, specialisation and integration within the prostate gland parenchyma. Notable milestones in structural markers were identified, during the seventh month of the intrauterine phase and the eleventh and twelfth months of the foetal period. Epithelial cell secretory activity commenced during the animal's 36th month, while the formation of lobules and thickening of the capsule layer were observed during the terminal phase of gestation and the postnatal period, respectively. Concurrently, interlobular connective tissue expanded alongside epithelial cell formation and an increase in terminal capillary count, peaking during the pubertal phase of the animals. In summary, our results underscore the association between ante- and postnatal phases of ontogenesis and the initiation of puberty, shedding light on the intricate anatomical and histological dynamics of the dromedary prostate.

**Key words:** Anatomy, dromedary, histology, ontogenesis, prostate

The camel prostate, notably substantial and easily identifiable, comprises of two distinct elements: a compact and a diffuse segment and divides into three zones: central, peripheral and transition (Mahmud *et al*, 2016). Together, these segments form an L-shaped structure positioned dorsally to the pelvic urethra. Shaaeldin and Tingari (2019) found that the camel prostate is oval-shaped, encircling the proximal part of the urethra at the bladder neck, with ducts opening around the entire urethral circumference. Dorsally, the prostate splits into two lobes separated by a septum.

Ultrastructure of the prostate of camel has been studied previously (Ali *et al*, 1976; Soliman *et al*, 2010), illustrating it as a complex of branched tubule-alveolar glands whose ducts open into the prostatic urethra. Moreover, it is enveloped by a fibro-elastic capsule rich in smooth muscle cells, with septa from

this capsule effectively segmenting the gland into lobes, albeit indistinct in adult male camels, a finding echoed in research on gazelles by Mohammed *et al* (2007).

While existing literature elucidates the anatomy, topography (Degen and Lee, 1982) and histological structure of the camel's prostate (Luo *et al*, 2016), there is a paucity of studies examining its prenatal and postnatal developmental phases. Building upon this gap, Biancardi *et al* (2017) emphasised the importance of morphological developmental analysis, linking it to systems theory and organ tissue regional concepts. Consequently, the aim of this study was to conduct a comprehensive investigation into the histological changes in the prostate gland of camel from prenatal to postnatal stages, elucidating critical developmental periods and their relationship with the onset of puberty.

SEND REPRINT REQUEST TO HAFIZA A [email: h.aidoudi@univ-soukahras.dz](mailto:h.aidoudi@univ-soukahras.dz)

## Materials and Methods

### Ethical approval

The sampling and data collection in this study was performed in accordance to the ethical recommendations of the Algerian government.

The present research was conducted on camels from a slaughterhouse in El-Oued -a province in the southeast of Algeria- and spanned from July 2021 to May 2023. Thirty prostates were obtained from clinically healthy and unvaccinated male foetuses aged between 07 to 13 months, neonatally. They were taken from animals aged 12, 24, 36, 48 and 60 months postnatally, with foetal age verified through femur length measurement. The methodological framework adhered to in this study followed the protocols outlined by Kadim *et al* (2008) and Saber *et al* (2020).

The investigations took place at the Laboratoire des Sciences et Techniques du Vivant, Institut d'Agriculture et des Sciences Vétérinaires, Université de Souk Ahras, Algeria. Specimens selected to elucidate morpho-functional changes in prostates concerning age were preserved using a 10% neutral buffered formalin solution. Subsequently, some fixed organ fragments were embedded in paraffin to facilitate the creation of thin sections measuring 3-5 microns thick.

For histological examination of the prostate's structure, staining with haematoxylin and eosin, the protocols of Cardiff *et al* (2014) and Abrahamsson *et al* (1989) were employed. Additionally, frozen sections of prostate tissue underwent staining with silver nitrate impregnation to highlight proteins prominently expressed on silicone.

Stained histological sections were examined and morphometry and microphotography were conducted using a MOTIC DM-52 light microscope, mono, digital (eyepiece x4, objective x10/0.25, x40/0.65).

Statistical analysis was carried out using standard "statist SF" software, with data analysis performed using SPSS Version 26.0.2019 (SPSS, Armonk, NY: IBM Corp). Data were expressed as means  $\pm$  SD (standard deviation) and subjected to statistical analysis using Duncan's test, with statistical significance set at  $P < 0.05$ .

### Results

Histological analysis of neonatal camel prostates revealed a thin prostatic capsule that was mainly composed of collagen and which is surrounded by isolated cells and isolated bundles of internal smooth muscle. Each tubuloalveolar gland was encapsulated in fibrous connective tissue that contained delicate

bundles of smooth muscle fibres. This glandular organisation takes the form of lobes subdivided into lobules by thin collagenous partitions, with scattered smooth muscle cells in this arrangement (Fig 1).

Examination of the thickness of the prostatic capsule revealed a significant variation ( $P < 0.01$ ) with the age of the subjects studied. At the 7th month of gestation, capsule thickness reached  $54.32 \pm 0.67 \mu\text{m}$ , increasing significantly to  $76.32 \pm 1.96 \mu\text{m}$  at the end of gestation. During the postnatal period in the spring, the thickness of the prostatic capsule in newborns measured  $85.1 \pm 1.04 \mu\text{m}$  ( $P < 0.05$ ). At the age of 60 months, this measurement reached a value of  $98.6 \pm 1.27 \mu\text{m}$ .

Concerning the secretory ducts, the application of silver nitrate to the histological sections revealed three distinct colours, indicating different cellular structures. At low magnification, the presence of multiple lumens suggests the presence of grouped empty spaces, with two forms of these lumens observed in foetal prostates.

The intra-lobular ducts were lined with tall columnar secretory cells. The interlobular ducts and the main excretory ducts were lined with stratified transitional epithelium visible in histological sections in neonatal prostates. Light brown staining indicated the presence of reticular tissue which was concentrated around urethral areas rich in this type of tissue.

Moreover, the intensity of the proliferative process in the connective tissue of the prostate varied, creating a distinctive architecture of the acini, leading to active proliferation of the mucous glands, a feature only observed in animals aged two years and over, while the acinar cells of the intercalary glands showed less active proliferation, the proliferative activity of the acinar cells of the main glands was lower during the prenatal period. These results justify the rejection of the term "prostatic activity" (Fig 2).

At higher magnification, the glandular epithelium appeared with a white lumen and a pale eosinophilic layer bordering the prostate glands. Papillary projections, sometimes with a connective or vascular axis, bordered the prostate glands and were seen in this lumen. Although this feature was absent in the prostates of camel foetuses, it was visible in those of newborns, particularly the older ones (Fig 3).

The development of true acini in the glandular epithelium is exceptionally rare, beginning only from the age of 24 months according to the researcher's observations.



The secretory endings revealed that large fibromuscular septa originate in the muscle bundles, distinctly delimiting the lobules of the gland, particularly in neonates. The branches of these septa, also fibromuscular, give rise to intra-lobular collecting ducts, lined with a two-layered pseudostratified epithelium, at the heart of each lobule. Numerous small ducts branch into tubular outgrowths which terminate in tubular endings.

In fact, it was noted that myoepithelial cells were absent and these septa, which were seen as detached filaments in the lobules, form a delicate intra-lobular stroma surrounding the alveoli that was constituting the secretory endings. These endings, aligned in the active secretory gland, were characterised by simple cylindrical cells with compressed nuclei. Each alveolus was surrounded by a capillary network within a sparse framework of loose connective fibres. There were variations in the stages of secretion. While some alveoli have glandular cells with dark pink cytoplasm, others had less dilated cytoplasm due to secretion.

The body of the prostate remains inactive in late-gestation foetuses with groups of undilated intra-lobular ducts without tubular secretory endings. It was also revealed from the sections of the pelvic part of the urethra that the transitional epithelium was similar to that of the body.

As far as the glandular epithelium is concerned, from the 7<sup>th</sup> month of gestation, the prostate takes the form of solid cell buds on the urethral epithelium. By the 8<sup>th</sup> month, some tubules are ducted, but most remain without a lumen. At this stage, the prostate is lined with a morphologically identical simple epithelium, frequently expressed by large pyramidal cells with flattened nuclei at the base. These characteristics persist in the prostate of the 9-month-old embryo.

The excretory ducts were initially lined with a two-layer epithelium, replaced by a transitional epithelium near the point of entry into the urethra. These changes occurred at a stage when the appearance of the prostate approaches that of a normal adult camel. The number of glandular tubules increases, accompanied with an increase in intertubular connective tissue.

Examination of the glandular epithelium's height of neonatal prostate glands reveals significant fluctuations throughout the postnatal period, with a minimum height at 12 and 60 months. Between 24 and 48 months, the height of the glandular epithelium was relatively uniform in the central, peripheral and transitional zones. From 36 months, the epithelium in

the central zone became significantly higher than that in the other two zones. At 48 months, the thickness of the lobule mucosa in postnatal study animals differed slightly by about 2 to 3  $\mu\text{m}$ . The epithelial height of all three layers of the prostate in 48-month-old animals was greater, suggesting a decrease in epitheliocyte height. In the 60-month-old animals, the surface area of the lobule was significantly increased, reaching around 1.5 times its initial size.


Concerning the lumen and number of glands, lobular formations of different sizes were observed in the prostatic part of 12-month-old foetuses. In newborns aged 60 months and over, a significant increase in prostate volume was observed, accompanied with a dilatation of the terminal sections of the prostate. The septa became thin and narrow. As such, the smooth muscle fibres became thinner, leading to a visible reduction in the thickness of the interlobular septa and a slight increase in the height of the epithelium of the terminal sections of the prostate was seen.

Observation of the interlobular connective tissue in foetal prostates showed that it appeared around the 9<sup>th</sup> month of gestation, with an initial thickness of  $26.61 \pm 3.21 \mu\text{m}$ , increasing to  $45.41 \pm 3.15 \mu\text{m}$  at the end of gestation. The height of the epithelium of the secretory endings in the foetal prostates reached  $4.57 \pm 2.45 \mu\text{m}$  at the 9<sup>th</sup> month of gestation and increased to  $16.58 \pm 2.67 \mu\text{m}$  at the end of gestation.


The diameter of the secretory ducts reached  $13.57 \pm 2.07 \mu\text{m}$  at the 9<sup>th</sup> month of gestation and doubled to reach  $16.58 \pm 2.67 \mu\text{m}$  at the end of gestation. It doubled to reach  $26.12 \pm 3.47 \mu\text{m}$  at the end of gestation, with the thickness of the epithelium increasing by  $4.68 \pm 3.47 \mu\text{m}$  to  $12.35 \pm 3.78 \mu\text{m}$ . The diameter of the illuminated acini was  $21.65 \pm 3.87 \mu\text{m}$  at the 9<sup>th</sup> month of gestation and increased to  $38.25 \pm 2.74 \mu\text{m}$  at the end of gestation. The diameter of the non-illuminated acini was  $14.57 \pm 2.36 \mu\text{m}$  at the 9<sup>th</sup> month of gestation and increased to  $32.47 \pm 2.74 \mu\text{m}$  at the end of gestation. It was also noted that the lumen of the glands decreases significantly from 12 to 36 months and remained relatively constant up to 60 months. This trend, correlated with previous data in the tables on glandular content at different ages, suggests an increase in the number of glands over time.

Examination of the number of glands in the prostates of camels at different stages of development revealed their presence after birth, with a slight increase at two months, followed by a significant decrease towards the third year of life, to reach minimum values (Table 1).






**Fig 1.** Histological section of the dromedary prostate gland in the postnatal period, **A:** 12<sup>th</sup> month after birth, **B:** 60<sup>th</sup> month after birth, stained with Hematoxylin and Eosin; 1-acinar cells; 2- connective tissue; 3-intraluminal secretions; 4-stroma; 5-duct. X 100.



**Fig 2.** Histological section of the dromedary prostate gland during the antenatal period, **A:** 7<sup>th</sup> month of gestation, **B:** 11<sup>th</sup> month of gestation, Hematoxylin and Eosin staining; 1-acinar cells; 2-connective tissue; 3-intraluminal secretions; 4-stroma; 5-duct. X100.



**Fig 3.** Histological section of the dromedary prostate gland during the antenatal period, **A:** 7<sup>th</sup> month of gestation, **B:** 11<sup>th</sup> month of gestation, silver nitrate staining; 1-acinar cells; 2-connective tissue; 3-intraluminal secretions; 4-stroma; 5-duct. X 100.



Discussion

The histological description of the developing prostate during the prenatal and postnatal period in animals has been studied extensively. The present research focuses on the analysis of prostate development in the dromedary. As an organ, the prostate is surrounded by a fibromuscular capsule. Previous studies on boars (Sakairi *et al*, 2003) have reported that the gland is covered externally by striated urethral muscles and lined internally by an intermittent layer of smooth muscle. In contrast, studies on bovines (Abou-Elmagd and Wrobel, 1989) described the capsule surrounding the gland as a dense, irregular connective tissue containing numerous smooth muscle cells and striated muscles. Our research findings are consistent with those of Gibson *et al* (2013), who observed that the prostate surrounds the urethra at its origin from the urinary bladder. This finding may be related to the need to produce a large amount of seminal fluid to ensure reproduction under dry conditions. The smooth muscle cells present in the capsule of the prostate could help to control the flow of seminal fluid towards the urethra.

Previous research on sheep by Lopes *et al* (2011) showed that the interlobular ducts were characterised by a low cylindrical epithelium lining. In contrast, the gland ducts in the current study were found opened into the urethral canal, with the excretory ducts exhibiting a single cylindrical epithelium lining. In addition, we observed a transition in the epithelial composition of the prostate, where single columnar epithelium becomes transitional epithelium in the terminal segments of the ducts. This may be related to the difference in environmental conditions, prostate epithelial cells may be more sensitive to desiccation in a desert climate. In this case, the presence of transitional epithelium in the terminal segments of the prostate ducts could help to protect the epithelial cells from desiccation.

Conversely, Hafez and Hafez (2001) reported the presence of predominantly serous glands with some mucosal acini in the disseminated pars of camels, which is also consistent with current observations, with columnar pseudostratified epithelium lining the disseminated pars. The solid and luminal secretory ends of the pars disseminata remained undifferentiated until three years of age. In this research, similar results were found at this stage of foetal development as the prostatic epithelium showed distinctive features. The results also indicated that the formation of the main definitive structural features of the prostates as peripheral reproductive organs occur in camels during the foetal period of stepwise ontogeny. Similarly, results from Xia *et al* (1990) mentioned the same suggestions for other mammals. However, the prostatic body does not appear in pigs in textbook descriptions. In that regard, Lossi *et al* (2016) reported that the disseminated Pars appears in the early postnatal period, but a narrower band of white glandular tissue with a glandular mass was found in camels. It is reported that this band has a condensed structure and is deeply buried in the muscle tissue of the urethra during the foetal period of the animal with a progressive increase in the degree of structure and differentiation of the different organ components. Consequently, the lobular structure of mammalian prostates is formed during the foetal period and the base of each component of these organs is a distinct gland. The number of which in some prostates is genetically determined (Angelsen *et al*, 1997). The body of the prostate is a glandular mass surrounded by a distinct fibrous capsule and exposed by the urethral muscle. The fibroreticular capsule of the gland also contains abundant smooth muscle cells. Similar findings have also been described by the authors and this remark has been found in prostates of mouse (Oliveira *et al*, 2016).

While interlobular connective tissue decreases in quantity with age, the remaining glandular

Table 1. Morphometric parameters of the prostate in a mature one-humped camel (x ± SD, n =8).

Months	Interlobular connective tissue thickness	Secretory tips			Secretory ducts	
		Diameter of illuminated acini	Diameter of non-illuminate acini	Height of epithelium	Duct diameter	Height of epithelium
Antenatal period						
9-10	26,61±3,21	21,65±3,87	14,57±2,36	4,57±2,45	13,57±2,07	4,68±3,14
11-12	45,41±3,15	38,25±2,74	32,47±3,74	16,58±2,67	26,12±3,47	12,35±3,78
Postnatal period						
0-12	33,45±2,54	36,14±3,18	42,68±2,24	15,57±2,47	43,14±3,14	11,24±2,11
48-60	11,34±3,41	49,54±2,54	45,73±3,22	21,24±3,28	49,47±3,14	18.24±3,87

SD: Standard Deviation of the mean



components increase. This result may be linked to the increased production of male sex hormones androgens, which stimulate the growth and differentiation of the glandular cells of the prostate. It should be noted that the main excretory duct of the gland is lined with stratified cuboid epithelium. In comparison, the epithelium in the prostate of rams is characterised by a simple cuboid to cylindrical structure with the presence of some basal cells (Odabaş and Kanter, 2008). This is probably due to lower androgen levels in this species. Rams also have a shorter reproductive cycle than dromedaries, which may also contribute to the difference in epithelium. The ducts were more abundantly branched with age and terminated in more luminal secretory ends. Initially, they were mainly mucous, but serous components were added by differentiation of the non-luminous secretory tips later in foetuses and neonates as confirmed by Krill *et al* (1999). Rodger and Hughes (1973) reported the presence of numerous tubular compounds in the prostate. Thus, the free surface of the glandular epithelium is clearly delineated and the epithelium is bilayered. Again, the proportion of glandular tissue to connective tissue is roughly the same in the late foetus. Nevertheless, the epithelial cells are larger, although still slightly smaller than in camel foetuses. In this study, we noted that whereas the prostatic ducts start to appear from the 7th month of gestation, they develop and come into place in cattle between 1 and 15 days after birth and 2 weeks after birth according to Groot and Biolatti (2004).

Studies performed on Bengal goat prostates by Gofur (2019) showed that only luminous acini were considered for micrometry, although some non-luminous acini were observed in the prostatic body of day-old goats. In contrast, the diameter of the acini found with a value of  $44.98 \pm 1.31 \mu\text{m}$  and that of our results, maximum sums of  $49.54 \pm 2.54 \mu\text{m}$  for luminous acini and  $45.73 \pm 3.22 \mu\text{m}$  for non-luminous acini of 48-60 month old neonates, the height of the epithelial cells found in the same breed was  $12.89 \pm 0.28 \mu\text{m}$ , whereas in our research, we found a sum of  $18.24 \pm 3.87 \mu\text{m}$  in neonates aged 48-60 months. We conclude that these indices vary according to the animal species by its body structures and its state of evolution.

The most important development of the prostate parenchyma growth took place during the prepubertal phase and spanned from birth to 18 months. These results are consistent with the observations made by Higgins and Gosling (1989) regarding the growth of parenchymal components

in human newborns, which persisted until the age of 12 months before reaching a static phase until 60 months.

## Conclusion

Histological differentiation and complete formation of prostatic tissue have been observed in the dromedary throughout its life, from the ante-natal period to adulthood, in three main phases:

– Embryonic phase (7<sup>th</sup> month of gestation): formation of the first epithelial cords and glandular tubules, lined with an undifferentiated stratified epithelium.

– Neonatal phase (8<sup>th</sup> month-end of gestation): development of the parietal and caudal parts of the gland. The parenchyma is represented by epithelial cords derived from the mucosa of the urogenital canal.

– Postnatal phase (1<sup>st</sup>-3 years): maturation of the gland, with increase in size, glandular and connective tissue and decrease in muscular tissue.

## Acknowledgement

The authors would thank the staff of the laboratory of histology in the Institute of Agriculture and Veterinary Sciences, University of Souk Ahras, Algeria for their comments that greatly improved the manuscript.

## Competing interest statement

The authors state that there is no competing interest in this research.

## References

- Abou-Elmagd A and Wrobel KH. The periurethral glandular complex in the water buffalo: an ultrastructural, histological and lectin-histochemical study. *Archives of Histology and Cytology*. 1989; 52(5):501-512. <https://doi.org/10.1679/aohc.52.501>
- Abrahamsson PA, Falkmer S, Fält K and Grimelius L. The course of neuroendocrine differentiation in prostatic carcinomas: an immunohistochemical study testing chromogranin A as an "endocrine marker". *Pathology - Research and Practice*. 1989; 185(3):373-380. [https://doi.org/10.1016/S0344-0338\(89\)80016-0](https://doi.org/10.1016/S0344-0338(89)80016-0)
- Ali HA, Moniem KA and Tingari MD. Some histochemical studies on the prostate, urethral and bulbourethral glands of the one-humped camel (*Camelus dromedarius*). *The Histochemical Journal*. 1976; 8:565-578. <https://doi.org/10.1007/BF01003958>
- Angelsen A, Mecsei R, Sandvik AK and Waldum H. Neuroendocrine cells in the prostate of the rat, guinea pig, cat and dog. *The Prostate*. 1997; 33(1):18-25. [https://doi.org/10.1002/\(SICI\)1097-0045\(19970915\)33:1<18::AID-PROS4>3.0.CO;25](https://doi.org/10.1002/(SICI)1097-0045(19970915)33:1<18::AID-PROS4>3.0.CO;25)

- Biancardi MF, Dos Santos FC, de Carvalho HF, Sanches BD and Taboga SR. Female prostate: historical, developmental and morphological perspectives. *Cell Biology International*. 2017; 41(11):1174-1183. <https://doi.org/10.1002/cbin.10759>
- Cardiff RD, Miller CH and Munn RJ. Manual hematoxylin and eosin staining of mouse tissue sections. *Cold Spring Harbor Protocols*. 2014; (6):655-8. doi: 10.1101/pdb.prot073411.
- Degen AA and Lee DG. The male genital tract of the dromedry (One-Humped) camel (*Camelus dromedarius*): gross and microscopic anatomy. *Anatomia, Histologia, Embryologia*. 1982; 11(3):267-282. <https://doi.org/10.1111/j.1439-0264.1982.tb00995.x>
- Gibson E, Gaed M, Gómez JA, Moussa M, Pautler S, Chin JL, Crukley C, Bauman GS, Fenster A and Ward AD. 3D prostate histology image reconstruction: Quantifying the impact of tissue deformation and histology section location. *Journal of Pathology Informatics*. 2013; 4(1):31. <https://doi.org/10.4103/2153-3539.120874>
- Gofur MR. Effect of testosterone on penile and urethral development in Black Bengal goat. *Bangladesh Journal of Veterinary Medicine (BJVM)*. 2019; 17(2):133-139. <https://doi.org/10.33109/bjvmjd19m3>
- Groot MJ and Biolatti B. Histopathological effects of boldenone in cattle. *J. Vet.* 2004; 51(2):58-63. <https://doi.org/10.1111/j.1439-0442.2004.00606.x>
- Hafez ESE and Hafez B. Reproductive parameters of male dromedary and bactrian camels. *Archives of Andrology*. 2001; 46(2):85-98. <https://doi.org/10.1080/01485010120937>
- Higgins JRA and Gosling JA. Studies on the structure and intrinsic innervation of the normal human prostate. *The Prostate*. 1989; 15(S2):5-16. <https://doi.org/10.1002/pros.2990150503>
- Kadim IT, Mahgoub O and Purchas RW. A review of the growth and of the carcass and meat quality characteristics of the one-humped camel (*Camelus dromedarius*). *Meat Science*. 2008; 80(3):555-569. <https://doi.org/10.1016/j.meatsci.2008.02.010>
- Krill D, Stoner J, Konety BR, Becich MJ and Getzenberg RH. Differential effects of vitamin D on normal human prostate epithelial and stromal cells in primary culture. *Urology*. 1999; 54(1):171-177. [https://doi.org/10.1016/S0090-4295\(99\)00103-X](https://doi.org/10.1016/S0090-4295(99)00103-X)
- Lopes WD, Santos TR, Luvizotto MCR, Sakamoto CAM, Oliveira GP and Costa AJ. Histopathology of the reproductive system of male sheep experimentally infected with *Toxoplasma gondii*. *Parasitology Research*. 2011; 109:405-409. <https://doi.org/10.1007/s00436-011-2268-9>
- Lossi L, D'Angelo L, De Girolamo P and Merighi A. Anatomical features for an adequate choice of experimental animal model in biomedicine: II. Small laboratory rodents, rabbit and pig. *Annals of Anatomy-Anatomischer Anzeiger*. 2016; 204:11-28. <https://doi.org/10.1016/j.aanat.2015.10.002>
- Luo Y, Li H, Lv Y, Xu S, Liu Y, Zhang N, Wang D, Shao B and Wang J. Anatomical and histochemical features of the bulbourethral glands in bactrian camel (*Camelus bactrianus*). *Journal of Camel Practice and Research* 2016; 23(1):121-125. <https://doi.org/10.5958/2277-8934.2016.00020.5>
- Mahmud MA, Josephat O, Abdullahi SS, Aminu UM, Abdurrahman B, Akawu SH, Abubakar D and Shehu S. Species variation on gross morphology and gross morphometry of accessory sex glands in one-humped camel bull (*Camelus dromedarius*), Uda Ram and Red Sokoto Buck. *World's Veterinary Journal*. 2016; 6(2):53-58. <http://hdl.handle.net/123456789/2867>
- Mohammed OB, Omer SA and Sandouka MA. The efficacy of Ivermectin and Levamisole against natural *Nematodirus spathiger* infection in the Arabian sand gazelle (*Gazella subgutturosa marica*) and the Arabian mountain gazelle (*Gazella gazella*) in Saudi Arabia. *Veterinary parasitology*. 2007; 150(1-2):170-173. <https://doi.org/10.1016/j.vetpar.2007.09.009>
- Odabaş Ö and Kanter M. Histological investigation of testicular and accessory sex glands in Ram lambs immunized against recombinant GNRH fusion proteins. *Electronic Journal of General Medicine*. 2008; 5(1):21-26. <https://doi.org/10.29333/ejgm/82570>
- Oliveira DS, Dzinic S, Bonfil AI, Saliganan AD, Sheng S and Bonfil RD. The mouse prostate: a basic anatomical and histological guideline. *Bosnian Journal of Basic Medical Sciences*. 2016; 16(1):8. <https://doi.org/10.17305/bjbm.2016.917>
- Rodger JC and Hughes RL. Studies of the accessory glands of male marsupials. *Australian Journal of Zoology*. 1973; 21(3):303-320. <https://doi.org/10.1071/ZO9730303>
- Saber ASM, Abd-Elnaeim MM and Shoghy KM. Radiological Study on the Osteogenesis of the Dromedary Skull (*Camelus dromedarius*). *Journal of Veterinary Anatomy*. 2020; 13(1):1-19. <https://doi.org/10.21608/JVA.2020.109648>
- Sakairi A, Tsukise A, Nara T, Habata I and Yamada K. Histochemical analysis of glycoconjugates in the disseminate prostate gland of the wild boar (*Sus scrofa riukiuanus*). *Acta Histochemica et Cytochemica*. 2003; 36(5):455-463. <https://doi.org/10.1267/ahc.36.455>
- Shaaeldin AH and Tingari MD. Ultrastructural and morphometric studies on the prostate gland of the one-humped camel (*Camelus dromedarius*). *Journal of Camel Practice and Research*. 2019; 26(1):21-27. <https://doi.org/10.1267/ahc.36.455>
- Soliman SM, Mazher KM and Abdelrazek AH. Light and electron microscopic studies of the prostate gland of adult one humped camel (*Camelus dromedarius*). *Journal of Veterinary Medicine and Research*. 2010; 20(1):44-51. <https://doi.org/10.21608/JVMR.2020.77579>
- Xia T, Blackburn WR and Gardner WA. Foetal prostate growth and development. *Pediatric Pathology*. 1990; 10(4):527-537. <https://doi.org/10.3109/15513819009067141>

## CAMEL CONSERVATION AND WELL-BEING



The workshops held in Nagaur, Rajasthan, India, 16-17 February 2024 focusing on Camel Conservation and Well-Being, shed light on the precarious state of the region's camel population. Despite being revered as the state animal, camels have faced a troubling decline exacerbated by legislative measures aimed at their protection, such as the Rajasthan camel law enacted in 2015. This legislation, while well-intentioned, has led to unintended consequences, including increased suffering and mortality rates among camels. Against this backdrop, stakeholders gathered to

discuss strategies for improving camel welfare, enhancing economic opportunities, and preserving traditional camel husbandry practices.

One significant aspect addressed by the workshops was the empowerment of women stakeholders in the camel husbandry sector. Working Group 1 emphasised the importance of women's active participation and leadership in the industry, advocating for training programs, ownership policies, and the creation of fair value chains. By providing women with the skills and resources to engage in camel husbandry and milk-based product production, these initiatives not only enhance economic viability but also promote gender equality and community development.

Another critical focus area was the expansion of market reach for camel milk, recognised for its nutritional value and potential to support sustainable agriculture. Working Group 2 devised a multi-faceted strategy to increase consumer awareness, engage health professionals, and diversify distribution channels. By positioning camel milk as a beneficial dietary supplement and advocating for ethical production practices, stakeholders aim to elevate its profile and consumption, thereby enhancing economic opportunities for camel pastoralists.

In addition to these efforts, Working Group 3 concentrated on improving the well-being of camels in Rajasthan through initiatives such as rehabilitating grazing lands, establishing camel sanctuaries, and addressing mistreatment issues. These measures aim to create suitable habitats for camel herds, promote conservation efforts, and ensure access to proper care and nutrition for all camels, regardless of their past treatment.

Lastly, Working Group 4 proposed strategies for enhancing camel grazing access in Rajasthan, emphasizing the conservation of grazing lands, establishment of camel sanctuaries, and creation of connecting corridors between traditional grazing grounds. These efforts seek to safeguard the well-being of camels while maintaining ecological balance and promoting sustainable land management practices.

Overall, the workshops underscored the importance of collaborative efforts to address the complex challenges facing Rajasthan's camel population. By empowering women stakeholders, expanding market opportunities for camel products, improving camel welfare, and enhancing access to grazing lands, stakeholders aim to ensure the sustainable coexistence of agriculture, animal husbandry, and natural ecosystems in the region. The insights and recommendations gleaned from these workshops offer a roadmap for charting a sustainable path forward for camel conservation and livelihoods in Rajasthan and beyond.

(Courtesy: Ilse Kohler Röllefson and Hanwant Singh, LPPS, Sadri, India)



# HAEMATOLOGICAL PROFILE VARIATIONS IN FEMALE DROMEDARY CAMELS DURING DRY PERIOD, EARLY PREGNANCY, LACTOGENESIS AND GALACTOPOIESIS IN ALGERIAN DROMEDARY CAMELS

Maria Chikha<sup>1</sup>, Tarek Khenenou<sup>1</sup>, Djalel Eddine Gherissi<sup>2</sup>, Soundes Akriche<sup>3</sup>, Mohammed Saci Ferdia<sup>4</sup>, Rabah Mayouf<sup>5</sup> and Sabry Mohamed El-Bahr<sup>6,7</sup>

<sup>1</sup>Laboratory of Science and Techniques for Living, <sup>2</sup>Laboratory of Animal Production, Biotechnologies and Health, Institute of Agricultural and Veterinary Sciences,

Mohamed-Cherif Messaadia University - Souk Ahras, BP 41000 Souk-Ahras, Algeria

<sup>3</sup>National Research Ogarev Mordovia State University, Saransk, Russia

<sup>4</sup>Department of Governance and Finance, Faculty of Economics and Management University of Sfax, Airport Road Km 4 Sfax 3018, Tunisia

<sup>5</sup>Faculty of NLS, Department of Agronomy Echahid Hamma Lakhdar University, The new group BP 789, 39000 El Oued, Algeria

<sup>6</sup>Department of Biomedical Sciences, College of Veterinary Medicine, King Faisal University, 31982 Al-Ahsa, Saudi Arabia

<sup>7</sup>Department of Biochemistry, Faculty of Veterinary Medicine, Alexandria University, 21526 Alexandria, Egypt

## ABSTRACT

The present study investigated the effects of different reproductive stages on haematological parameters of mature local dairy female dromedary camels (n=52) in El Oued region in southeastern Algeria under a semi-intensive breeding system. The haematological parameters analysed included white blood cell count (WBC), lymphocyte count (LYM), lymphocyte percentage (LYM%), monocyte count (MON), monocyte percentage (MON%), granulocyte count (GRAN), granulocyte percentage (GRAN%), red blood cell count (RBC), haemoglobin (HGB), haematocrit (HCT), mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC), and platelet count (PLT). The study focused on four distinct stages: dry period, early pregnancy, lactogenesis, and galactopoiesis. Significant increases ( $P<0.001$ ) were observed in MON% and GRAN% during the dry period and early pregnancy. MCV values were significantly higher ( $P<0.05$ ) during the dry period than early pregnancy. However, GRAN during these stages was significantly lower ( $P<0.001$ ) than during galactopoiesis. Lactogenesis and galactopoiesis stages were characterised by significant elevations ( $P<0.001$ ) in WBC, LYM, and LYM%. Additionally, HGB, MCHC, and MCH values were significantly higher ( $P<0.001$ ) during these stages. Conversely, RBC, HCT, and PLT were significantly lower ( $P<0.001$ ) during lactogenesis and galactopoiesis. Interestingly, the MON during galactopoiesis was significantly lower ( $P<0.001$ ) than during the dry period. These findings demonstrate that haematological parameters in female dromedary camels undergo significant alterations across different reproductive stages. This knowledge can contribute to improved reproductive management strategies and provide insights into the physiological adaptations of camels during various reproductive phases in semi-intensive breeding systems.

**Key words:** Camel, dry period, haematology, galactopoiesis, lactogenesis, pregnancy

Assessing the physiological, nutritional and pathological conditions of animals and their productivity, including camels, is often evaluated through haematologic profiles (Hafez, 2006; Tharwat *et al*, 2015). Various factors such as breed, sex, age, nutrition and seasonal changes have been observed affecting haematologic values (Tornquist, 2010; Farooq *et al*, 2011). The reproductive status of animals can also influence these parameters due to the effects of the endocrine system on multiple organs (Tharwat *et al*, 2015; Jalali *et al*, 2018). Dromedary camels exhibit

seasonal breeding patterns, with breeding seasons during winter and spring, while summer and autumn are typically non-breeding seasons for females (Tibary and Anouassi, 1996). Additionally, they are induced ovulators (Eman, 2019) and their pregnancy lasts approximately 387 days (Wilson and Payne, 1999).

Some studies have investigate the impact of different reproduction periods on haematological parameters (Hafez, 2006; Muhammad *et al*, 2011; Tharwat *et al*, 2015; Jalali *et al*, 2018; Ebissy *et al*, 2019; Elkhair, 2019; Chikha *et al*, 2024). However, there

SEND REPRINT REQUEST TO MARIA CHIKHA [email: m.chikha@univ-soukahrass.dz](mailto:m.chikha@univ-soukahrass.dz)

remains a lack of documented data on the impact of various reproductive periods, including the dry period, early pregnancy, lactogenesis and galactopoiesis on haematological parameters in semi-intensive breeding system of camel population in the south east of Algeria. Therefore, this study aimed to evaluate the effects of different reproductive periods on haematological attributes in one-humped female camels (*Camelus dromedarius*) in the southeast of Algeria.

## Materials and Methods

In this research, 52 adult female dromedary camels were used. The camels kept in a semi-intensive breeding system situated in El Oued region, southeastern Algeria. Throughout the study duration, the camels were found to be free of both internal and external parasites and with good health status.

### Blood Collection and analysis

In this study, blood samples were collected from a total of 52 animals, out of which 15 samples were obtained during the dry period, 13 during early pregnancy (first month), 12 during lactogenesis (last month of pregnancy) and 12 during early galactopoiesis (first month of lactation). The samples were collected and transported to the laboratory for haematological analysis. Haematological parameters including white blood cell count (WBC), lymphocyte count (LYM), monocyte count (MON), granulocyte count (GRAN), lymphocyte percentage (LYM%), monocyte percentage (MON%), granulocyte percentage (GRAN%), red blood cell count (RBC), haemoglobin concentration (HGB), haematocrit (HCT), mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC), platelet count (PLT) and mean platelet volume (MPV) were determined using a MINDRAY haematology auto analyser (BC-3000Plus, China) as used in the study of Chikha *et al* (2024).

### Statistical analysis

The haematological parameters data were analysed using IBM SPSS Statistics, version 25.0, based in Armonk, NY, USA. The findings were expressed as mean values  $\pm$  standard deviation (SD). To assess group differences across various lactation stages, a one-way ANOVA test was applied, followed by Tamhane T2 test as a post hoc analysis. The significance level was established at  $p < 0.05$ .

## Results and Discussion

The results of white blood cell (WBC), red blood cell (RBC) and platelet parameters are summarised

in Table 1 and Table 2, respectively. As depicted in Table 1, there was no significant difference between the WBC count in early pregnancy and the dry period, which was notably lower compared to the values during lactogenesis and galactopoiesis. Consistent with these findings, previous work (Atta *et al*, 2021) reported lower values of these parameters during the dry period and early pregnancy. This alteration was linked to physiological adaptations associated with foetal growth and development throughout pregnancy (Muhammad *et al*, 2011). Conversely, other studies (Saeed *et al*, 2011); Ebissy *et al*, 2019) did not demonstrate any notable variance induced by pregnancy. Early pregnancy was associated with a significantly lower lymphocyte count than both the lactogenesis and galactopoiesis phases as well as the dry period. Similar results (Atta *et al*, 2021) were reported where the number of lymphocytes significantly increased in lactating camels and decreased significantly in dry and pregnant ones. While the monocyte (MON) count during galactopoiesis was significantly lower than during the dry phase, it was not statistically different from the counts during early pregnancy and lactogenesis. In addition, a significant increase was observed during lactation in MON (Atta *et al*, 2021). In the current study, the granulocyte count during dry period and early pregnancy was significantly lower than during galactopoiesis. Atta *et al* (2021) also reported higher significant values of granulocyte count during lactation. The current findings indicated that there was a considerable variation in LYM% between the various times; the value was low during early pregnancy and extremely significant during galactopoiesis in comparison to other periods. Parallel to the current findings, a significant decrease in lymphocyte percentage was noted during late pregnancy compared to early lactation (Elkhair, 2019). However, no significant difference was reported between the dry period and early pregnancy, but a high value was reported during lactation (Atta *et al*, 2021). Data of the current study indicated that, during the dry period MON% was significantly higher compared to the early period, lactogenesis and galactopoiesis, with the lowest value observed during galactopoiesis. Consistent to the current study, a significant increase in monocyte percentage was observed during early lactation compared to late pregnancy periods (Elkhair, 2019). In the current study, the percentage of granulocytes showed no significant difference between the different periods. A significant difference between the dry period, pregnancy and lactation was reported earlier (Atta *et al*, 2021). The significant

changes observed in the percentages of lymphocytes, monocytes and granulocytes could be linked to the release of cortisol and ACTH in response to the stresses of pregnancy and lactation (Ahmed, 2017; Jainudeen and Hafez, 2000).

In the current study, RBC and HCT were significantly lower during lactogenesis and galactopoiesis compared to the dry period and early pregnancy. A significant increase in the number of RBCs was reported in lactating camels compared to the dry period (Atta *et al*, 2021). The variation in erythrocyte parameters according to physiological status may be attributed to the heightened demand for oxygen consumption and the need for a higher metabolic rate to support growth, particularly during late pregnancy and early lactation (Elkhair and Minawy, 2018). The current study reported that, Haemoglobin (HGB) levels were significantly higher during lactogenesis and galactopoiesis compared

to dry period and early pregnancy. Similar results (Tharwat *et al*, 2015; El Zahar *et al*, 2017), indicated that higher values of Hb were reported during late pregnancy and early lactation in camels. Contrarily, other study (Atta *et al*, 2021) reported a slight increase in HGB levels in dry camels compared to postpartum and lactating ones. The higher levels of haemoglobin during lactogenesis and galactopoiesis could be attributed to the combination of increased elevated metabolic rate and hormonal changes during these periods. In the current study, Mean Corpuscular Volume (MCV) values were significantly higher during dry period than early pregnancy and no significant difference was observed between lactogenesis and galactopoiesis. Similar results (Ebissy *et al*, 2019) were reported earlier. The low value of MCV during early pregnancy could be the result of hormonal changes that prioritise resources towards supporting foetal growth and development.

**Table 1.** White blood cell parameter values in female camels during different reproductive periods (Dry, Early Pregnancy, Lactogenesis, Galactopoiesis).

Parameters	Period				P value
	Dry period (n=15)	Early pregnancy (n=13)	Lactogenesis (n=12)	Galactopoiesis (n=12)	
WBC (×10 <sup>9</sup> /L)	12.50±2.87 <sup>a</sup>	11.65±2.30 <sup>a</sup>	27.06±8.95 <sup>b</sup>	65.77±19.17 <sup>c</sup>	0.000
LYM (×10 <sup>9</sup> /L)	3.60±1.40 <sup>a</sup>	1.23±0.35 <sup>b</sup>	35.30±13.45 <sup>c</sup>	47.38±15.59 <sup>c</sup>	0.000
MON (×10 <sup>9</sup> /L)	2.58±1.28 <sup>ab</sup>	1.71±1.03 <sup>b</sup>	2.22±0.77 <sup>b</sup>	2.64±1.44 <sup>a</sup>	0.000
GRAN(×10 <sup>9</sup> /L)	6.34±3.12 <sup>a</sup>	8.76±1.71 <sup>a</sup>	9.54±3.44 <sup>ab</sup>	12.40±3.60 <sup>b</sup>	0.000
LYM%	29.06±8.60 <sup>a</sup>	12.16±2.54 <sup>b</sup>	63.60±8.00 <sup>c</sup>	73.041±7.38 <sup>d</sup>	0.000
MON%	15.16±3.31 <sup>a</sup>	10.80±3.03 <sup>b</sup>	9.17±2.38 <sup>b</sup> <sup>c</sup>	6.94±1.69 <sup>c</sup>	0.000
GRAN%	52.85±12.21 <sup>a</sup>	75.084±5.64 <sup>b</sup>	31.86±9.68 <sup>c</sup>	17.30±3.19 <sup>d</sup>	0.000

a,b,c; Values within a column that have different superscripts vary significantly from each other (p<0.05). WBC; Number of white blood cells, LYM; Lymphocyte, MON; Monocyte, GRAN; Granulocyte, LYM%; Per cent Ratio of Lymphocyte, MON%; Per cent Ratio of Monocyte, GRAN%; Per cent ratio of Granulocyte.

**Table 2.** Red blood cell and platelets parameter values in female camels during different reproductive periods (Dry, Early Pregnancy, Lactogenesis, Galactopoiesis).

Parameters	Stage of lactation				P value
	Dry period (n=15)	Early pregnancy (n=13)	Lactogenesis (n=12)	Galactopoiesis (n=12)	
HGB (g/dl)	11.06±1.39 <sup>a</sup>	14.23±1.34 <sup>b</sup>	124.38±9.44 <sup>c</sup>	121.41±6.27 <sup>c</sup>	0.000
RBC (×10 <sup>12</sup> /L)	5.38±0.62 <sup>a</sup>	5.34±0.54 <sup>a</sup>	4.60±0.43 <sup>b</sup>	4.23±0.58 <sup>b</sup>	0.000
HCT%	23.36±2.58 <sup>a</sup>	22.39±2.17 <sup>a</sup>	19.80±1.65 <sup>b</sup>	19.01±2.43 <sup>b</sup>	0.000
MCV fL	43.21±0.47 <sup>a</sup>	42±0.68 <sup>bc</sup>	43.15±1.43 <sup>ac</sup>	42.98±1.65 <sup>ac</sup>	0.029
MCH pg	23.34±1.85 <sup>a</sup>	26.05±1.97 <sup>b</sup>	28.25±4.19 <sup>b</sup>	28.72±3.82 <sup>b</sup>	0.000
MCHC g/dL	53.98±4.40 <sup>a</sup>	63.86±7.24 <sup>b</sup>	655.33±73.58 <sup>c</sup>	594±32.80 <sup>c</sup>	0.000
PLT (×10 <sup>9</sup> /L)	211.73±36.45 <sup>a</sup>	154.74±13.24 <sup>b</sup>	137.16±19.05 <sup>b</sup> <sup>c</sup>	139.91±12.38 <sup>c</sup>	0.000
MPV fL	6.62±0.56	6.33±0.84	6.47±0.75	7.058±0.83	0.110

a,b,c; Values within a column that have different superscripts vary significantly from each other (p<0.05). RBC; Number of red blood cells, HGB; Haemoglobin concentration, HCT; Haematocrit, MCV; Mean corpuscular volume, MCH; Mean corpuscular haemoglobin, MCHC; Mean corpuscular haemoglobin concentration, PLT; Number of Platelets, MPV; Mean Platelet Volume.



The current findings indicated that high significant values of Mean Corpuscular Haemoglobin (MCH) and Mean Corpuscular Haemoglobin Concentration (MCHC) were observed during lactogenesis and galactopoiesis. However, no significant difference was observed between lactogenesis and galactopoiesis in the current study. These results were in the contrary to the results that reported earlier (Ebissy *et al*, 2019), whereas, a significant difference was reported between antepartum and postpartum days. The high values of MCH and MCHC could be the result of adaptations to optimise oxygen transport and metabolism, including alterations in erythropoiesis and haemoglobin synthesis to meet the increased metabolic demands of milk production. In the current study, a low significant level of platelets (PLT) was observed during lactogenesis and galactopoiesis compared to dry period and early pregnancy. This low value could be attributed to the physiological stress associated with lactation and the demands of milk production, which can impact platelet levels. The current findings indicated that there was no significant difference in Mean Platelet Volume (MPV) values during the different periods.

In conclusion, the current data illustrates how several time periods such as the dry period, early pregnancy, lactogenesis and galactopoiesis affect the haematological parameters of dromedary camels and may serve as a useful guide for she-camels.

### Conflict of interests

The authors have not declared any conflict of interests.

### References

- Ahmed M. Effect of Selenium and Vitamin E Injection During Transition Period on Physiological Performance of Camels (*Camelus dromedarius*) and their Neonates Reared Under -Semi-intensive System. MVSc Thesis. 2017; Khartoum University. Sudan.
- Atta AS, El-Shourbagy I, Gawish HA, El-Metwaly HA, Younis FE and Salem ML. Farming and pastoral systems shape the physiological and immunological indices of the dromedary she-camels. *International Journal of Clinical Biochemistry and Research*. 2021; 5(1):49-63
- Chikha M, Khenenou T, Gherissi DE, Mayouf R, Rahmoun DE, Al-Gabri N, Ramdani N, Abu-Sulik AA, Aidoudi H and El-Bahr SA. Lactation related changes of haematological parameters of female dromedary camels reared under semiintensive farming system in algerian extreme arid region. *Journal of Camel Practice and Research*. 2024; 31(1):25-28
- Ebissy EA, El-Sayed AA, Ragab HM. Haematological and biochemical profile in female camels (*Camelus dromedarius*) during the transition period. *Slovenian Veterinary Research*. 2019; 56(22):571-7
- Elkhair NM and Minawy AA. Effect of the physiological status on the haematological parameters of female camels (*Camelus dromedarius*). *Journal of Veterinary Medicine and Animal Production*. 2018; 9(2):154-162
- Elkhair NM. Monitoring of the Haematological Parameters in Camels (*Camelus dromedarius*) during Late Pregnancy and Early Lactation. *EC Veterinary Science*. 2019; 4(8):675-680
- El-Zahar H, Zaher H, Alkablawy A and Swelum AS. A monitoring the changes in certain haematological and biochemical parameters in camels (*Camelus dromedarius*) during postpartum period. *Journal of Fertility Biomarkers*. 2017; 1(1):47-54.
- Eman E, El-Sayed A, Hassan Mohamed R. Haematological and biochemical profile in female camels (*Camelus dromedarius*) during the transition period. *Slovenian Veterinary Research*. 2019; 56(22):571-7.
- Farooq U, Samad HA, Khurshid A and Sajjad S. normal reference haematological values of one-humped camels (*Camelus dromedarius*) kept in cholistan desert. *The Journal of Animal & Plant Sciences*. 2011; 21(2): pp157-160
- Hafez YM. Haematological and blood plasma biochemical constituents of one humped camel (*Camelus dromedarius*) in relation to some physiological, nutritional and parasitological conditions. *Egyptian Journal of Animal Production*. 2006; 43(1):157-168
- Jainudeen M and Hafez E. Gestation, Prenatal Physiology and Parturition. *Reproduction in Farm Animals*, 7<sup>th</sup> Edition. Wiley. 2000; pp 140-155
- Jalali SM, Hasani YN, Darabifard A, Mavadati AH. A study of haematologic and biochemical profile in female dromedary camels during the breeding and non-breeding seasons. *Comparative Clinical Pathology*. 2018; 27:655-659
- Muhammad B, Aliyu D, Njidda A and Madigawa I. Some haematological, biochemical and hormonal profile of pregnant and nonpregnant she-camels (*Camelus dromedarius*) raised in a Sudan savanna zone of Nigeria. *Journal of Camel Practice and Research*. 2011; 18(1): 73-77
- Saeed A, Khan IA and Hussein M. Change in haematological profile of pregnant camels (*Camelus dromedarius*) at term. *Comparative Clinical Pathology*. 2011; 20(1):53-55
- Tharwat M, Ali A, Al-Sobayil F, Selim L and Abbas H. Haemato-biochemical profile in female camels (*Camelus dromedarius*) during the periparturient period. *Journal of Camel Practice Research*. 2015; 22(1):101-106
- Tibary A and Anouassi A (1996) Ultrasonographic changes of the reproductive tract in the female camel during the follicular cycle and pregnancy. *Journal of Camel Practice Research*. 1996; 3:71-90
- Tornquist SJ. Haematology of camelids. In: Schalm's *Veterinary Haematology*. Wiley-Blackwell Publishing. 2010; pp 910-917
- Wilson RT and Payne JA. An introduction to Animal Husbandry in The Tropic Blackwell Science, Ltd. 1999; pp 1-120.

# PREVALENCE INVESTIGATION OF GASTROINTESTINAL NEMATODES IN SONID BACTRIAN CAMELS

Bin Hou, Mila Tai and Surong Hasi

College of Veterinary Medicine, Inner Mongolia Agricultural University; Key Laboratory of Clinical Diagnosis and Treatment Technology in Animal Disease, Hohhot, Inner Mongolia 010018, China

## ABSTRACT

To investigate the prevalence, intensity and predominant species of gastrointestinal nematode infections in Sonid Bactrian camels, Inner Mongolia, a survey on the prevalence of gastrointestinal nematodes was conducted on 307 naturally grazed Bactrian camels from 5 regions in Inner Mongolia using faecal egg, larvae and adult nematode identification methods. The results showed that gastrointestinal nematodes were quite severe in Sonid Bactrian camels, with an overall infection rate of 87.6% (269/307) and an average infection intensity of 1,981 (EPG). Among them, the infection rates of Bactrian camels were generally higher in Abaga Banner, West Urumqi Banner, Durbed Banner, Sonid East Banner and Sonid West Banner, which were 100% (15/15), 100% (15/15), 98.6% (71/72), 98.1% (101/103) and 65.6% (67/102), respectively with infection intensities ranging from 50 to 10,350 (EPG). *Trichostrongylus* spp. was the predominant species, followed by *Ostertagia ostertagi*. This result was validated through PCR testing of DNA extracted from 46 faecal samples collected from Sonid Bactrian camels, confirming the findings. Therefore, the present study provides preliminary epidemiological survey data on the major nematode infections prevalent in Sonid Bactrian camel herds in Inner Mongolia, contributing to the understanding of gastrointestinal nematodiasis in Sonid Bactrian camels.

**Key words:** Bactrian camels, epidemiological investigation, gastrointestinal nematodes, PCR

The gastrointestinal nematodes of Bactrian camels mainly include species such as *Haemonchus contortus*, *Trichostrongylus* spp., *Ostertagia ostertagi*, *Nematodirus* spp. and *Marshallagia* spp. Studies have reported that in Algeria, the infection rate of gastrointestinal parasites in dromedaries is 48.26%, with parasites from 12 genera identified, among which the infection by nematodes of the *Strongyloides* spp. is the most severe, while those of the *Cooperia* spp. were the least prevalent (Bouragba *et al*, 2020). Researchers conducted gastrointestinal nematode examinations on 144 dromedaries in Iran, identifying 26 species of nematodes with an infection rate of 86.3%, primarily by *Haemonchus contortus* and *Trichostrongylus colubriformis* (Anvari-Tafti *et al*, 2013). The population of Bactrian camels is relatively small, accounting for less than 10% of the total old world camelids population, most studies on gastrointestinal nematodes in camels have focused on dromedaries. Sonid Bactrian camels are mainly distributed in Inner Mongolia Autonomous Region, including Xilin Gol League, Ulanqab City and eastern Hulunbuir, with a focus on Sonid West Banner, Sonid East Banner, Abaga Banner and West Urumqi Banner in Xilin

Gol city (He, 2003). The modern Bactrian camel breeding industry is gradually shifting from simple labour purposes to comprehensive utilisation such as milk, meat and fur, with the breeding population showing an increasing trend year by year. However, nematode infections, mainly affecting Sonid Bactrian camels, are prevalent, significantly impacting the productivity of camel herds and hindering the healthy development of the local camel breeding industry. Moreover, the majority of herders have insufficient understanding of the harmfulness economic losses caused by gastrointestinal nematodes, leading to a lack of scientific and targeted approaches in the prevention and control of nematode diseases and a prevalence of indiscriminate drug administration, resulting in a significant lag in parasitic disease prevention and control efforts (Eyeledege, 2018).

In this study, a systematic investigation of gastrointestinal nematodes in Sonid Bactrian camels in 5 different pastoral counties in Inner Mongolia was conducted, aiming to understand the dominant species, epidemiological status and infection intensity of gastrointestinal nematodes in Sonid Bactrian camels.

SEND REPRINT REQUEST TO SURONG HASI [email: baohaas@163.com](mailto:baohaas@163.com)

Materials and Methods

Ethics Statement

Animal procedures were performed in accordance with the National Standard Guideline for Ethical Review of Animal Welfare (GB/T 35892-2018) and approved by the Animal Care and Use Committee of Inner Mongolia Agricultural University.

Collection of Sample

From January 2022 to December 2023, a total of 307 Bactrian camel fresh faecal samples were collected in Durbed Banner, Sonid West Banner, Sonid East Banner, Abaga Banner and West Uzumqin Banner of Inner Mongolia Autonomous Region, China, with 72, 102, 103, 15 and 15 samples, respectively. Samples were collected by individuals wearing disposable gloves, who directly collected faeces from the rectum or the centre of faecal balls not in contact with the ground (10g-50g). The samples were then placed in self-sealing bags, labeled with the Bactrian camel’s basic information and transported back to the laboratory under refrigeration for examination. Faecal samples that could not be examined immediately were stored in a 4°C refrigerator to prevent hatching of eggs that could affect the interpretation of the results.

Identification of nematode eggs

The identification of gastrointestinal nematode species is based on the morphological structure and

characteristic features of different eggs, referring to the *Colour Atlas of Morphological Classification of Parasites of Domestic Animals in China*. Additionally, the identification of larvae structures cultured from faeces and the confirmation of adult worms obtained through dissecting deceased camels was done.

Faecal eggs quantitative examination

Faecal egg counts were undertaken using the modified McMaster technique, as described in Veterinary Clinical Parasitology (Anne and Gary, 2012).

PCR identification

Forty-six faecal samples of Bactrian camels were collected from Sonid West Banner, categorised into 5 groups (I, II, III, IV and V) based on the collection locations, with EPG (eggs per gram) values ≥500. PCR detection was conducted on faecal DNA samples extracted from each group using specifically designed primers for 8 common gastrointestinal nematodes of Bactrian camels (some specific primers can identify down to the species level, while other specific primers can only identify down to the genus level), negative control without DNA template.

DNA extraction was performed using the Stool DNA Kit D4015-01 from OMEGA Bio-Tek (USA), following the manufacturer’s instructions and the extracted DNA was stored at -20°C. Specific primers (Appendix) for eight gastrointestinal nematodes

Appendix. Sequences of primers specific for 8 gastrointestinal nematodes of the bactrian camel.

Nematode species	Primer	Primer sequence (5'-3')	Target fragment(bp)	Annealing temperature
<i>Haemonchus contortus</i> (Zou <i>et al</i> , 2023)	Upstream primer	ATTGTTCTGCTCAAATGGCA	270	53°C
	Downstream primer	AGTTTCTTTTCTCCGCT		
<i>Nematodirus</i> spp. (Jia, 2014)	Upstream primer	GTAGGTGAACCTGCGGAAGGATCATT	800	45°C
	Downstream primer	TTAGTTTCTTTTCTCCGCT		
<i>Trichuris ovis</i> (Zhang <i>et al</i> , 2013)	Upstream primer	TTTGATATCTTTTACCTTACCATT	900	55°C
	Downstream primer	AGGGCTTATTGCTATGTGGTTA		
<i>Parabronema skrjabini</i> (Zheng, 2015)	Upstream primer	TTTACAAGAGGGATACGCC	551	50°C
	Downstream primer	GGTATCACAACTTATCGGG		
<i>Strongyloides</i> spp.	Upstream primer	CACCTCTTCAGGGACAT	433	47°C
	Downstream primer	TTTGGAGCATTTGGAT		
<i>Ostertagia ostertagi</i> (Qu, 2013)	Upstream primer	CGCTTAGAGTGGTAAAAATTTGAAC	342	57°C
	Downstream primer	TTAGTTTCTTTTCTCCGCTAAATG		
<i>Chabertia</i> spp. (Zhao <i>et al</i> , 2013)	Upstream primer	TTTTTTGGGCATCCTGAGGTTTAT	450	55°C
	Downstream primer	TAAAGAAAGAACATAATGAAAATG		
<i>Trichostrongyle</i> spp. (Li, 2020)	Upstream primer	TTGTCGAAACCAACACATGG	173	58°C
	Downstream primer	GGGAACCTTCGCATGAACAAT		




synthesised from literature were utilised for PCR amplification (Table 1).

After PCR amplification, 5 µL of the product was taken for 1% agarose gel electrophoresis. The voltage was set to 110 V and electrophoresis was conducted for 30 minutes. Subsequently, the gel was placed into a UV gel imaging system for result observation and photographing for documentation.


### Result

#### The infection status of gastrointestinal nematodes of Sonid Bactrian camels

According to Table 2, the overall infection rate of gastrointestinal nematodes in 307 faecal samples of Bactrian camels from Sonid is 87.6% (269/307), with 269 samples testing positive. The infection rates of gastrointestinal nematodes in Durbed Banner, Sonid West Banner, Sonid East Banner, Abaga Banner and West Uzumqin Banner were 98.6% (71/72), 65.6% (67/102), 98.1% (101/103), 100% (15/15) and 100% (15/15), respectively.



**Fig 1.** Eggs of *Trichostrongylus* spp. and *Nematodirus* spp. (100X).




**Fig 2.** Third-stage (L3) larvae of *Trichostrongylus* spp. (400X).


**Table 1.** The PCR reaction system by special primers (25 µL)

Components	Volume
Upstream primer	1 µL
Downstream primer	1 µL
Premix Taq	12.5 µL
Template DNA	2 µL
dH <sub>2</sub> O	8.5 µL
Total volume	25 µL

The highest EPG of gastrointestinal nematodes in Durbed Banner, Sonid West Banner, Sonid East Banner, Abaga Banner and West Uzumqin Banner



**Fig 3.** Third-stage (L3) larvae of *Ostertagia ostertagi* (400X).




**Fig 4.** Female anterior end and vulva of *Trichostrongylus* spp. (100X).



**Fig 5.** Female anterior end and reproductive tract of *Nematodirus* spp. (100X)

**Table 2.** Nematode infection in the digestive tract of Sonid Bactrian camels of Inner Mongolia.

Areas	No. Samples	No. Positives	Rate of infection (%)	Infection intensity (EPG)	Mean intensity of infection (EPG)
Durbed Banner	72	71	98.6	50-10,350	1,810
Sonid West Banner	102	67	65.6	50-1,800	396
Sonid East Banner	103	101	98.1	50-7,850	1,409
Abaga Banner	15	15	100	1,000-6,050	3,246
West Uzumqin Banner	15	15	100	250-5,750	3,043
Total	307	269	87.6	50-10,350	1,981



**Fig 6.** Electropherogram of PCR amplification results. M: DNA Maker DL2000; A. *Trichostrongylus* spp. B. *Ostertagia ostertagi*. C. *Nematodirus* spp. D. *Chabertia* spp. E. *Haemonchus contortus*. F. *Trichuris ovis*. G. *Parabronema skrjabini* H. *Strongyloides* spp.; I, II, III, IV and V are subgroups; N: negative control.

were 10 350, 1 800, 7 850, 6 050 and 5 750, with average infection intensities of 1 810, 396, 1 409, 3 246 and 3 043, respectively.

### Results of egg, larval hatching and adult worm dissection and identification

According to microscopic examination results, the most prevalent nematode species observed were

from the *Trichostrongylidae*, mainly *Trichostrongylus* spp. (Figs 1,2,4) and *Ostertagia ostertagi* (Fig 3), followed by the *Nematodirus* spp (Figs 1,5).

The eggs of *Trichostrongylus* spp. measure 76-92×37-46 µm, with relatively well-filled contents. The eggs of *Nematodirus* spp. measured 165-175×76-86 µm and contained 6-8 embryonated cells (Fig 1).

The larvae have a relatively stout body, with a clearly visible excretory pore and 16 distinct triangular intestinal cells. The tail is thick and blunt, with either one or two segments or an indistinct rounded tail tip and a very short tail sheath. Measurements indicate a total length of 622-706 µm and a length from caudal end to tip of caudal sheath 31-39 µm.

The larvae were of moderate size, with two refractile bodies or a bright bar between the mouth and esophagus visible. The excretory pore was clearly visible and the tail was relatively thick with a short tail sheath.

### Results of PCR Test

PCR testing was conducted on 46 faecal samples from Bactrian camels in Sonid, revealing 4 species of nematodes tested positive (Fig 6). *Trichostrongylus* spp. had the highest infection rate, followed by *Ostertagia ostertagi*. Among the 46 faecal samples from Sonid Bactrian camels, 41 were positive for *Trichostrongylus* spp., resulting in a total infection rate of 89.1% (41/46). The infection rates for Groups I, II, III, IV and V were 100%, 100%, 100%, 44.4% and 100%, respectively.

*Ostertagia ostertagi*. had a total infection rate of 76.1% (35/46), with infection rates in Groups I, II, III, IV and V being 100%, 100%, 100%, 44.4% and 25%, respectively.

*Nematodirus* spp. had a total infection rate of 54.3% (25/46), with infection rates in Groups I, II, III,

IV and V being 100%, 20%, 40%, 66.6% and 37.5%, respectively.

*Chabertia* spp. had a total infection rate of 13% (6/46), with infection rates in Groups I, II, III, IV and V being 44.4%, 10%, 0%, 0% and 10%, respectively.

Nematodes such as *Haemonchus contortus*, *Trichuris ovis*, *Parabronema* spp. and *Strongyloides* spp. all tested negative.

## Discussion

Due to the rampant prevalence of gastrointestinal nematodes in Bactrian camels in the Inner Mongolia autonomous region of China in recent years, the productivity and product quality of Bactrian camels have declined, causing serious economic losses to the camel husbandry industry. This study conducted a preliminary investigation and analysis of the prevalence of gastrointestinal nematodes in Sonid Bactrian camels in Durbed Banner of Ulanqab City and Sonid West Banner, Sonid East Banner, Abaga Banner and West Uzumqin Banner of Xilin Gol city in Inner Mongolia. The results showed that the overall infection rate of gastrointestinal nematodes in 307 faecal samples from Sonid Bactrian camels was as high as 87.6% (269/307), indicating a severe situation of gastrointestinal nematodes in Sonid Bactrian camels in Inner Mongolia. The infection rates of Bactrian camels in the 5 investigated areas, namely Durbed Banner, Sonid West Banner, Sonid East Banner, Abaga Banner and West Uzumqin Banner, were 98.6% (71/72), 65.6% (67/102), 98.1% (101/103), 100% (15/15) and 100% (15/15), respectively. The highest EPG values were 10 350, 1 800, 7 850, 6 050 and 5 750, respectively. Morphological identification of eggs, infective larvae and adult nematodes revealed that the most commonly infected nematode was *Trichostrongylus* spp., followed by *Ostertagia ostertagi*. and *Nematodirus* spp.

In previous studies, the prevalence of gastrointestinal nematodiasis in Bactrian camels was highly common, with both high infection rates and intensity (Abubakr *et al*, 2000; Fraser and Craig, 1997; Abd El-Wahed, 2005). *Trichostrongylus* spp. was identified as the predominant species, which aligns closely with the findings of this study. For instance, El-Alfy *et al* (2019) using different molecular markers, identified infection rates of 26%, 65.2%, 60.8% and 95.6% for *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Cooperia oncophora* and *Haemonchus contortus*, respectively (El-Alfy *et al*, 2019) in faecal samples from 101 single-humped camels in Egypt. Similarly, researchers conducted

PCR amplification and sequencing on faecal samples from 160 camels in central Iraq, revealing an infection rate of 18.13% (29/160) (Rasool *et al*, 2021). Research on gastrointestinal nematodes in Bactrian camels in China has primarily focused on the *Parabronema skrjabini* (Zhou, 2021; Li *et al*, 2020; Zhao *et al*, 2012). An epidemiological survey by Zheng (2015) reported a significant decrease in the prevalence of *Parabronema skrjabini* infection in Bactrian camels in Sonid, Xilin Gol League, from 4,696 worms per camel in 1988 to 16 worms per camel. In this study, *Parabronema skrjabini* was not detected in the genetic analysis of 46 faecal samples from Bactrian camels in Sonid West Banner, indicating some effectiveness in the prevention and control of *Parabronema skrjabini* disease in the Xilin Gol area over the past two decades.

In conclusion, the prevalence of gastrointestinal nematodes in Bactrian camels in Sonid, Inner Mongolia, is relatively common, especially with severe infections of *Trichostrongylus* spp., which significantly hinder the healthy development and economic benefits of camel husbandry in the region. The epidemiological findings of this study are of great significance for understanding the prevention and control of gastrointestinal nematodes in Sonid Bactrian camels, providing scientific evidence and technical support for the protection of Bactrian camel health and the promotion of sustainable development in animal husbandry.

## Conflict of interests

The authors declare no conflict of interests.

## Acknowledgements

This work was supported by National Natural Science Foundation of China (32060815) and Natural Science Foundation of Inner Mongolia (2020MS03011).

## Reference

- Abd El-Wahed MM. Incidence of *Nematodirus* spp. and their differentiation through the infective third stage larvae among Egyptian camels. *Journal of the Egyptian Society of Parasitology*. 2005; 35(2):447.
- Abubakr IM, Nayel MN, Mohamed FE, Abdelrahman AO, Abuobeida SA and Elgabara YM. Prevalence of gastrointestinal parasites in young camels in Bahrain. *Revue D' Elevage et de Medecine Veterinaire des Pays Tropicaux*. 2000; 53(3):267-270.
- Anvari-Tafti M, Sazmand A, Hekmatimoghaddam S and Moobedi I. Gastrointestinal helminths of camels (*Camelus dromedarius*) in centre of Iran. *Tropical Biomedicine*. 2013; 30(1):56-61.
- Bouragba M, Laatamna A, Cheddad FE, Baroudi D, Houali K and Hakem A. Gastrointestinal parasites of dromedary



- camel (*Camelus dromedarius*) in Algeria. *Veterinary World*. 2020; 13(8):1635-1640.
- El-Alfy ES, Abu-Elwafa S, Abbas I, Al-Araby M, Al-Kappany Y, Umeda K and Nishikawa Y. Molecular screening approach to identify protozoan and trichostrongylid parasites infecting one-humped camels (*Camelus dromedarius*). *Acta Tropica*. 2019; 197:105060.
- Eyeledege. Research on the epidemiology of major helminthic diseases in sheep and investigation of deworming effect in Uttarakhand. Inner Mongolia Agricultural University. 2018.
- Fraser A and Craig PS. Detection of gastrointestinal helminth infections using coproantigen and molecular diagnostic approaches. *Journal of Helminthology*. 1997; 71(2):103-108.
- He XM. Atlas of Camel Resources in China. Changsha: Hunan Science and Technology Press. 2003;72-73.
- Jia YQ. Molecular characterisation and genetic variation of sheep nematode. Northwest Agriculture and Forestry University. 2014.
- Jirimutu, Wang Z, Ding G, *et al.* Genome sequences of wild and domestic bactrian camels. *Nature Communications*. 2012; 3:1202.
- Li RQ. Analysis of the prevalence of elk parasitosis and the molecular evolution of major species in Beijing Elk Park in 2018-2019. Shanxi Agricultural University. 2020.
- Li Z, Zhao XY, Tian WL, Chen JZ, Guo YT, Hu XY, Li SN, Tian RX, Dong WL, Su ZQ, Yao G, Rang DL, Fu Q and Shi HJ. Investigation on the infection situation of gastrointestinal parasites in Bactrian Camels in some areas of Xinjiang. *Heilongjiang Animal Husbandry and Veterinary Medicine*. 2020; (15):98-100+104+174.
- Qu GG. Establishment and application of detection method of *Ostreococcus australis* nematode and its macrophage transfer inhibitory factor function identification. Jilin University. 2013.
- Rasool AA, Issa AF and Rasool S F. Prevalence and molecular identification of *Nematodirus helvetianus* in camels in Iraq. *Veterinary World*. 2021; 14(5):1299-1302.
- Zhang Y W, Xu Y, Wang Y, Zhu XQ and Liu GH. Cloning and evolutionary analysis of partial sequence of mitochondrial rrnL gene in woolly tail nematode. *Journal of Hunan Agricultural University: Natural Science Edition*. 2013; (1):4.
- Zhao L, Jiang HH, Zhao GH, Liu AG, Zhu XQ and Qian AD. Cloning and sequence analysis of the mitochondrial pcox1 gene in the nematode *Schabertia lamblia*. *Journal of Hunan Agricultural University: Natural Science Edition*. 2013; 39(5):4.
- Zhao ZG, Yang XY, Yang LR, Wang R, Wang HY, Zhao LL, Ao WH and Guo TZ. Epidemiological investigation of *Subulura* spp. disease in Bactrian camels in Inner Mongolia. *Chinese Veterinary Science*. 2012; 42(12):6.
- Zheng WQ. Epidemiologic investigation and establishment of prebiotic PCR diagnostic method for *S. parvum* nematodes in camels. Inner Mongolia Agricultural University. 2015.
- Zheng WQ. Epidemiological Investigation of *Subulura* ssp. Disease in Camels and Establishment of Pre-mortem PCR Diagnostic Methods. Inner Mongolia Agricultural University. 2015.
- Zhou Q. Investigation on the infection situation of gastrointestinal parasites in Bactrian camels in some areas of Xinjiang. Tarim University. 2021.
- Zou M, Yao HZ, Yang BK, Zhou LL, Zhao GM, Yang WH, Wu C and Lin Q. Establishment of a PCR method for the detection of twisted blood spear nematode and its application. *Northwest Journal of Agriculture*. 2023; 32(1):139-144.

# DETERMINATION AND IDENTIFICATION OF REFERENCE INTERVALS FOR SELECTED BLOOD PARAMETERS DURING DIFFERENT PHYSIOLOGICAL STATES IN FEMALE DROMEDARY CAMELS

Nawal Mohamed Elkhair<sup>1,2</sup>

<sup>1</sup>Department of Physiology, Faculty of Veterinary Medicine, University of Khartoum, 13314, Shambat, Sudan

<sup>2</sup>Department of Biomedical Sciences, College of Veterinary Medicine, King Faisal University, 31982, Al-Ahsa, Saudi Arabia

## ABSTRACT

One hundred eighty female Arabian camels aged 3–16 years were used to determine the reference intervals for selected blood parameters during different physiological states. Venous blood was collected for determining erythrocyte and leukocyte parameters, blood glucose, protein fractions,  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$  and Pi. The reference ranges for TEC, PCV, Hb, MCV, MCH, MCHC, TLC, neutrophils, eosinophils, basophils, lymphocytes and monocytes were 4.06–5.03 $\times 10^6/\mu\text{L}$ , 4.8–11.2 g/dL, 24.8–30.6%, 29.7–39.4 fL, 25.2–44.4 pg, 19.6–39.8 g/dL, 11–13 $\times 10^3/\mu\text{L}$ , 41–49%, 5–6%, 1–3%, 45–50% and 1–2%, respectively and for blood glucose, TP, albumin,  $\alpha_1$ ,  $\alpha_2$ ,  $\beta_1$ ,  $\beta_2$  and  $\gamma$ -globulins, A/G ratio,  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$  and Pi were 67–100 mg/dL, 29–72 g/L, 46–78%, 1–7%, 2–10%, 5–14%, 3–14%, 10–27%, 1–15 and 5–9, 1–2, 4–6 mmol/L, respectively. The values of Hb were lower in the growing and early lactating camels than in dry ones, whereas TEC, monocytes percentage and serum  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$ , Pi were higher ( $P < 0.05$ ). The TEC, TP and albumin were higher in late pregnant camels than in dry and early lactating ones ( $P < 0.05$ ). The  $\alpha_1$ ,  $\alpha_2$  and  $\beta_1$  were lower in late pregnant camels than in dry females, whereas they showed higher values in early lactating ones ( $P < 0.05$ ). Neutrophil and lymphocyte percentages showed higher and lower values ( $P < 0.05$ ) in late pregnant and early lactating camels than in dry females, respectively. The blood parameters of female camels varied significantly depending on their physiological state. The findings are most likely applied to evaluate physiological and metabolic adaptations to growth, pregnancy and lactation.

**Key words:** Blood parameters, dromedary camel, physiological state, reference interval

Blood parameters are commonly utilised to assess the physiological state of the animals (Onasanya *et al*, 2015). Age, sex, breed, season, exercise, nutrition, stress, transport and diseases have been shown to alter haematological and serum biochemical parameters in all animal species (Jain, 1998; Faye and Bengoumi, 2018). In camels, significant differences have been found in these parameters in relation to age and during different physiological states (Tharwat *et al*, 2015; Elkhair and Elmgboul, 2015; Elkhair, 2016; Ahmed, 2017; Elkhair and Minawy, 2018; Ahmed and Elkhair, 2019; Islam *et al*, 2019; Waziri *et al*, 2019; Faraz *et al*, 2021; Mohamed *et al*, 2021; Martín-Barrasa *et al*, 2023).

It has been demonstrated that metabolic processes change significantly during pregnancy and lactation in camels; therefore, many investigators

used blood parameters to evaluate the metabolic profile of pregnant and lactating camels (Ayoub *et al*, 2003; Ahmed, 2017; Axay *et al*, 2017; El Zahar *et al*, 2017; Abd El-Salaam and Arafa, 2018; Jalali *et al*, 2018; Elkhair, 2019; Ebissy *et al*, 2019; Atta *et al*, 2021).

Age, growth phase, pregnancy and lactation are physiologically critical states associated with significant metabolic alterations in all animal species. Regarding female camels, few data are available concerning the reference intervals of blood parameters as influenced by age, growth phase, late pregnancy and early lactation and to describe the haematology and serum biochemical profile of growing, pregnant and lactating camels reared under semi-intensive system. Therefore, the study was aimed to establish reference ranges for selected blood parameters of female camels under various physiological states.

SEND REPRINT REQUEST TO NAWAL MOHAMED ELKHAIR [email: nelkhair@kfu.edu.sa](mailto:nelkhair@kfu.edu.sa); nawalelkhair@yahoo.com

## Materials and Methods

The study was conducted with utmost adherence to ethical guidelines, ensuring the welfare of the animals at all times. It was carried out at the Camel Research Centre of the Faculty of Veterinary Medicine, University of Khartoum (Sudan), after receiving approval from the Faculty Research Ethics Committee. The animal care protocol and sample procedures were conducted in strict accordance with the ethics regulations set by the same body and the guidelines established by the ARRIVE standards (<https://arriveguidelines.org>) for animal experiments.

### Animals and Management

The selection process for the animals was rigorous and thorough. One hundred eighty clinically healthy female Arabian camels (ages: 3-5 years for growing females and 8-16 years for dry, late pregnant and early lactating camels, number of parities: 2-3) were randomly divided into four groups of 45 each. The females were raised in a semi-intensive system, browsing and grazing shrubs and trees; water was offered *ad libitum*. The females were fed a daily ration of fresh Abu 70 (*Sorghum biocolor* L.) and concentrate supplements (a mixture of crushed *Sorghum bicolor* grains 30%, wheat bran 40%, groundnut cake 25% and NaCl salt 5%). The pregnant animals were meticulously observed one month *prepartum* up to one month *postpartum* and were chosen based on mating records for each individual participant in the trial, which showed the predicted delivery.

### Collection of blood sample

The experimental animals were sampled twice at two-week intervals in the morning before feeding. Approximately 7 mL of venous blood was obtained at 8:00 am via jugular venipuncture using disposable plastic syringes (Jiangsu Kangsu Medical Instrument Co., Ltd, China). A volume of 2 mL was placed into EDTA vacutainers for haematological analysis. Another 2 mL was drawn into sodium fluoride vacutainers for glucose determination. The remaining blood sample was drawn into plain plastic syringes (Pirmveta®, Laboratory Technique, GmbH, Germany), centrifuged at 3000 rpm for 10 min (Gallenkamp Junior, UK). Free-hemolysis sera were placed into sterile vials (Eppendorf® Tube) and stored at -20 °C for serum biochemistry measurements.

### Laboratory analysis

**Haematological parameters:** The URIT-3010 VET Haematology Analyser (URIT medical

electronics) was employed to determine Hb, PCV, TEC, MCV, MCH, MCHC and TLC. The DLC was performed in thin Giemsa-Grunewald stained blood smears (Weiss and Wardrop, 2010).

**Blood glucose and serum Ca<sup>++</sup>, Mg<sup>++</sup> and Pi concentrations:** Standard spectrophotometric methods and Bio-systems kits (S. A. Barcelona, Spain) were used for determining blood glucose (Caraway, 1987), Ca<sup>++</sup> (Gindler and King, 1972), Mg<sup>++</sup> (Chromya *et al*, 1973) and Pi (Gamst, 1980).

**Total protein (TP) and capillary electrophoresis:** The Biuret standard spectrophotometric method described by Weichselbaum (1946) was performed for TP determination. Capillary electrophoresis was used to perform the percentages of albumin,  $\alpha_1$ ,  $\alpha_2$ ,  $\beta_1$ ,  $\beta_2$  and  $\gamma$ -globulins using an Automated MINICAP-SEBIA and MINICAP CDT kits (MINICAP-SEBIA Instrument, France). The instrument was checked for precision and accuracy following the guidelines of Sebia-MINICAP manual, Ref. 2208.

### Statistical analysis

The IBM SPSS Statistics 20 (Inc. SPSS, 2011), a comprehensive and reliable tool, was used for data analysis. Resource Equation Approach was performed for computing the sample size (Mead, 1988; Arifin and Zahiruddin, 2017). The reference values for blood parameters were determined using the minimum sample size in the survey ( $n_{Min}$ ) according to the formula adopted by Werner (1992).

$$n_{Min} \geq \left( \frac{U_{1-\alpha/2}}{\Delta} \right)^2 \cdot s^2 \quad (1)$$

The Kolmogorov-Smirnov test was applied to determine the distribution of the individual values; the data were normally distributed. General linear model procedures for analysis of variance ANOVA tests and descriptive statistics were applied to evaluate the differences between the groups; the differences were separated at  $P < 0.05$  using LSD.

## Results

### Overall reference range

Tables 1 and 2 demonstrate the overall reference ranges of 180 female camels, i.e. TEC (4.06–5.03×10<sup>6</sup>/μl), Hb (4.8–11.2 g/dL), PCV (24.8–30.6%), MCV (29.7–39.4 fl), MCH (25.2–44.4 pg), MCHC (19.6–39.8 g/dL), TLC (11–13×10<sup>3</sup>/μl), neutrophils (41–49%), lymphocytes (45–50%), monocytes (1–2%), eosinophils (5–6%) and basophils (1–3%). The established reference ranges of blood glucose, serum TP, albumin,  $\alpha_1$ ,  $\alpha_2$ ,  $\beta_1$ ,  $\beta_2$ ,  $\gamma$ -globulins, A/G ratio,



Ca<sup>++</sup>, Mg<sup>++</sup> and Pi were 67–100 mg/dL, 29–72 g/dl, 46–78%, 1–7%, 2–10%, 5–14%, 3–14%, 10–27%, 1–15 and 5–9, 1–2 and 4–6 mmol/L, respectively.

*Blood parameters of the growing females*

The TEC, monocytes percentage, serum Ca<sup>++</sup>, Mg<sup>++</sup> and Pi in the growing females were significantly

**Table 1.** Statistical data of the haematological parameters in female camels (*Camelus dromedarius*) during different physiological states.

Parameter	Statistical values	Growing	Dry	Late pregnant	Early lactating
TEC (×10 <sup>6</sup> /μl)	$\bar{x} \pm s$	5.00±0.68 <sup>A</sup>	4.63±0.96 <sup>Bb</sup>	5.03±0.65 <sup>a</sup>	4.06±0.96
	Reference range <sup>1</sup>	4.2–5.5	3.6–5.5	4.4–5.7	3.8–4.9
	Median	5.2	4.8	5.0	4.7
	(1.-3. Quartile)	4.2–5.5	3.6–5.5	4.4–5.7	3.0–4.8
[Hb] (g/dl)	$\bar{x} \pm s$	7.80±2.7 <sup>B</sup>	9.40±1.9 <sup>Ab</sup>	9.90±0.89 <sup>a</sup>	8.02±1.2 <sup>c</sup>
	Reference range <sup>1</sup>	4.8–10.2	7.4–11.2	8.9–10.6	6.8–9.2
	Median	8.3	9.6	10.2	8.1
	(1.-3. Quartile)	4.8–10.2	7.4–11.2	8.9–10.6	8.9–9.2
PCV (%)	$\bar{x} \pm s$	27.76±4.7	29.4±4.9	29.9±4.89	28.02±4.82
	Reference range <sup>1</sup>	24.8–30.2	27.4–31.2	28.9–30.6	26.8–29.2
	Median	28.3	29.6	30.2	28.1
	(1.-3. Quartile)	24.8–30.2	27.4–31.2	28.9–30.6	26.9–29.2
MCV (fl)	$\bar{x} \pm s$	32.93±2.8	35.3±2.8	36.03±2.9	34.34±2.3
	Reference range <sup>1</sup>	29.7–34.7	32.7–38.2	34–39.4	32.5–38.2
	Median	14.4	15	14.7	13.7
	(1.-3. Quartile)	9.7–14.7	12.7–18.2	14–19.4	12.5–18.2
MCH (pg)	$\bar{x} \pm s$	21.67±4.35	20±0.62	28.67±0.6	20.76±2.7
	Reference range <sup>1</sup>	25.2–41.5	39.5–40.7	38.1–39.3	38.1–44.4
	Median	38.3	39.8	38.6	41.3
	(1.-3. Quartile)	15.2–41.5	39.5–40.7	38.7–39.7	38.2–43.1
MCHC (g/dl)	$\bar{x} \pm s$	26.47±11.5	20.27±0.3	19.63±0.9	21.10±1.88
	Reference range <sup>1</sup>	19.6–39.8	20–20.6	18.6–20.2	19.1–23.4 21.8
	Median	20	20.2	20.1	19.2–22.7
	(1.-3. Quartile)	19.6–39.8	20.0–20.6	18.6–20.2	
TLC (×10 <sup>3</sup> /μl)	$\bar{x} \pm s$	11.93±4.9	11.97±4.85	12.3±4.82	12.22±4.9
	Reference range <sup>1</sup>	11–12.8	11.8–12.1	12.1–12.5	11–13.3
	Median	12	12	12.3	12.5
	(1.-3. Quartile)	11–12.8	11.8–12.1	12.1–12.5	11.3–13
Neutrophils (%)	$\bar{x} \pm s$	42.5±2.1	41.67±2.5 <sup>b</sup>	41.5±0.7 <sup>c</sup>	47.5±2.1 <sup>a</sup>
	Reference range <sup>1</sup>	41–44	39–44	41–42	46–49
	Median	42.5	42	42	47.5
	(1.-3. Quartile)	30.8–33	39–44	30.8–31.5	34.5–36.8
Lymphocytes (%)	$\bar{x} \pm s$	48.50±2.1	46.50±2.1 <sup>c</sup>	50.33±1.5 <sup>a</sup>	49.50±2.1 <sup>b</sup>
	Reference range <sup>1</sup>	47–50	45–48	49–52	48–51
	Median	48.5	46.5	50	49.5
	(1.-3. Quartile)	35.25–37.5	33.75–33	49–52	36–38.5
Monocytes (%)	$\bar{x} \pm s$	2±0.02 <sup>A</sup>	1±0.01 <sup>B</sup>	1±0.01	1.33±0.58
	Reference range <sup>1</sup>	2–2.1	1–1.2	1–1.2	1–2
	Median	2	1	1	1
	(1.-3. Quartile)	1.5–1.5	0.75–0.75	0.75–0.75	1–2
Eosinophils (%)	$\bar{x} \pm s$	5±0.04	3.5±0.7	5±1.4	4.3±0.58
	Reference range <sup>1</sup>	5–5.2	3–4	4–6	4–5
	Median	5	3.5	5	4
	(1.-3. Quartile)	3.75–3.75	2.3–3.0	3–4.5	4–5
Basophils (%)	$\bar{x} \pm s$	2±0.03	1.5±0.7	3±1.4	2.67±0.58
	Reference range <sup>1</sup>	2–2.2	1–2	2–4	2–3
	Median	2	1.5	3	3
	(1.-3. Quartile)	1.5–1.5	0.75–1.5	1.5–3	1.5–3

<sup>1</sup>m± s.d.×1.96 indicated the lower and the upper limits, Brackets ([ ]) denote concentration.

Aa, Bb Overall means within the same row bearing different superscripts are significantly different at P≤0.05.

**Table 2.** Statistical data of selected serum biochemical parameters in female camels (*Camelus dromedarius*) during different physiological states.

Parameter	Statistical values	Growing	Dry	Late pregnant	Early lactating
Blood glucose (mg/dL)	$\bar{x} \pm s$ Reference range <sup>1</sup> Median (1.-3. Quartile)	71.67±0.58 71–72 72 71–72	78.33±11 67–89 79 67–89	82.33±10.4 72–100 75 72–100	76.80±11.1 66–97 73 68–87
Serum-[TP] (g/l)	$\bar{x} \pm s$ Reference range <sup>1</sup> Median (1.-3. Quartile)	42.67±10.8 35–55 38 35–55	43.67±5.1 <sup>c</sup> 39–49 36 29–39	59.67±8.5 <sup>a</sup> 47–72 60 47–72	50.40±8.8 <sup>b</sup> 38–61 53 40–59
Albumin (%)	$\bar{x} \pm s$ Reference range <sup>1</sup> Median (1.-3. Quartile)	62.62±8.4 46–70 63 46–70	59.85±6.2 <sup>b</sup> 47–77 58 47–75	63.33±9 <sup>a</sup> 46–78 66 46–78	57.25±9.6 <sup>b</sup> 47–70 56 48.75–67
$\alpha_1$ -globulins (%)	$\bar{x} \pm s$ Reference range <sup>1</sup> Median (1.-3. Quartile)	2.5±2.07 2–3 2.5 2–3	2.71±2.13 <sup>a</sup> 2–3 2.5 2–3	1.33±0.58 <sup>b</sup> 1–2 1 1–2	3.75±3.4 <sup>a</sup> 2–7 3 2–6.3
$\alpha_2$ -globulins (%)	$\bar{x} \pm s$ Reference range <sup>1</sup> Median (1.-3. Quartile)	4.88±3.13 2–9 5 2–9	4.86±3.38 <sup>a</sup> 2–9 5 2–9	2.67±1.2 <sup>b</sup> 2–4 2 2–4	6.50±3.7 <sup>a</sup> 2–10 7 2.8–9.8
$\beta_1$ -globulins (%)	$\bar{x} \pm s$ Reference range <sup>1</sup> Median (1.-3. Quartile)	9±3.2 7–10 9 7–10	9.29±3.4 <sup>a</sup> 7–10 9 7–10	7±2.6 <sup>b</sup> 5–10 6 5–10	11±2.9 <sup>a</sup> 8–14 11 8.3–13.8
$\beta_2$ -globulins (%)	$\bar{x} \pm s$ Reference range <sup>1</sup> Median (1.-3. Quartile)	6.71±4.8 4–12 6.5 3.5–12.8	7.5±4.8 6–14 7.5 5.5–11.8	9.5±4.9 6–13 9.5 4.5–9.8	6.5±5.1 3–14 4.5 3.3–11.8
$\gamma$ -globulins (%)	$\bar{x} \pm s$ Reference range <sup>1</sup> Median (1.-3. Quartile)	15.63±6.7 11–18 16 11–18	17.14±5.6 11–19 17 11–19	19.67±7.5 12–27 20 12–27	15.25±3.8 10–18 16.5 11.3–18
A/G ratio	$\bar{x} \pm s$ Reference range <sup>1</sup> Median (1.-3. Quartile)	3.7±4.71 3–4 3.5 3–4	3.71±5.8 3–4 3.5 3–4	2.33±1.5 1–4 2 1–4	4.75±6.8 1–15 1.5 1–11.8
serum-[Ca <sup>++</sup> ] (mmol/l)	$\bar{x} \pm s$ Reference range <sup>1</sup> Median (1.-3. Quartile)	8.33±1.2 7–9 9 7–9	5.67±1.2 5–7 5 5–7	5.67±0.58 5–6 6 5–6	7.2±1.8 5–9 7 5–9
serum-[Mg <sup>++</sup> ] (mmol/l)	$\bar{x} \pm s$ Reference range <sup>1</sup> Median (1.-3. Quartile)	2±0.01 2–2.1 2 2–2.1	1.67±0.58 1–2 2 1–2	2±0.01 2–2.1 2 2–2.1	1.80±0.44 1–2 2 1.5–2
serum-[Pi] (mmol/l)	$\bar{x} \pm s$ Reference range <sup>1</sup> Median (1.-3. Quartile)	6±0.04 6–6 6 6–6.2	5±1.0 4–6 5 4–6	5.67±0.58 5–6 6 5–6	4.8±0.45 4–5 5 4–5

<sup>1</sup>m± s.d.×1.96 indicated the lower and the upper limits, Brackets ([ ]) denote concentration.

Aa, Bb Overall means within the same row bearing different superscripts are significantly different at P≤0.05.

higher (P<0.05) than in dry adult ones, while Hb was lower (P<0.05). The PCV, MCV, MCH, MCHC, TLC, the percentage of neutrophils, lymphocytes, eosinophils and

basophils, the blood glucose, serum TP, albumin and  $\alpha_1$ ,  $\alpha_2$ ,  $\beta_1$ ,  $\beta_2$  and  $\gamma$ -globulins did not change significantly in the growing females compared to the dry ones.

Fig 1 illustrates a serum protein electrophoretic pattern with six fractions. Albumin fraction exhibited approximately a similar pattern in growing (a) and dry females (b), while  $\alpha_1$ ,  $\alpha_2$ ,  $\beta_1$ ,  $\beta_2$  and  $\gamma$  fractions differed from those in dry adults.

### ***Blood parameters of late pregnant and early lactating camels***

The TEC and Hb were significantly higher ( $P<0.05$ ) in late pregnant camels than in dry ones, while Hb was lower ( $P<0.05$ ) in early lactating camels than in late pregnant and dry ones. Neutrophils percentage was significantly higher ( $P<0.05$ ) in early lactating camels than in dry females. In contrast, their percentage was lower ( $P<0.05$ ) in late pregnant camels than in early lactating and dry ones. Lymphocytes percentage was significantly higher ( $P<0.05$ ) in late pregnant camels than in early lactating and dry ones. The PCV, MCV, MCH, MCHC and TLC values and the percentage of eosinophils and basophils did not change significantly in late pregnant and early lactating camels.

Serum TP was significantly higher ( $P<0.05$ ) in late pregnant and early lactating camels than in dry ones (Table 2). The albumin fraction percentage was significantly higher ( $P<0.05$ ) in late pregnant camels than in dry and early lactating ones; however, the percentage was significantly lower ( $P<0.05$ ) in early lactating animals than in dry females. The percentage of  $\alpha_1$ ,  $\alpha_2$  and  $\beta_1$ -globulin fractions were lower ( $P<0.05$ ) in late pregnant than in dry and early

lactating camels, while the highest significant ( $P<0.05$ ) percentage was observed during early lactation. Camels in late pregnancy and early lactation revealed no substantial change in blood glucose,  $\beta_2$ - and  $\gamma$ -globulins, A/G ratio and serum  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$  and Pi.

Fig 2 shows albumin fraction exhibited approximately similar pattern in dry (a), late pregnancy (b) and early lactation (c). However,  $\alpha_1$ ,  $\alpha_2$ ,  $\beta_1$ ,  $\beta_2$  and  $\gamma$ - globulins displayed different patterns in response to the physiological state.

### **Discussion**

Determination of the reference ranges for blood parameters is essential for monitoring health, prognosis, diagnosis and treatment of diseases. To the author's knowledge, there is relatively little published data on the reference intervals of blood parameters in female camels. Therefore, we established a reference range for these parameters during various physiological states, which might be utilised for the clinical monitoring of females health. In dromedary camels, most prior reference values were obtained on undefined and relatively small groups, with no consideration given to the physiological states (Faye and Bengoumi, 2018). In clinical practice, it is critical to assess the number of animals required for establishing representative reference ranges.

In this study, we applied equation (1), as established by Werner (1992), to determine the minimum number of samples ( $n_{Min}$ ) required. Equation (1) clearly demonstrated that the reference




Fig 1. Serum protein capillary electrophoresis pattern of the growing (a) and dry adult female camels (b).



values for these parameters are mostly determined by the standard deviation of their individual values. The confidence level of  $\Delta = 1 \mu\text{l, g/dL, mmol/L}$  and  $\text{g/L}$  indicates that the average  $\delta$  of  $x$  values in the population with an error probability of  $\alpha = 0.05$  should not deviate more than  $1 \mu\text{l, g/dL, mmol/L}$ , or  $1 \text{ g/L}$  from the mean value of the selected groups. The PCV, TLC and blood glucose demonstrated no significant differences between the experimental females (Tables 1 and 2); therefore, we computed a mean value for the standard deviation ( $s$ ) of all groups for these parameters and substituted the estimated values of  $s = 4.8$  as  $\delta$  in Equation (1):

$$nMin \geq \left(\frac{1.96}{1}\right) \cdot (4.8)^2 = 45 \text{ animals}$$

The determined minimum number of camels ( $nMin$ ) equal to or less than 45 represents the precise number of females in each group; hence, the reference ranges for each parameter suggest reference values under defined physiological conditions. The  $nMin$  for all parameters studied using the same equation was comparable to or less than the minimum number of animals employed (Tables 1 and 2).

The mean and reference interval for the erythrocytes parameters (PCV, Hb, TEC, MCV, MCH and MCHC) reported for the growing, dry, late pregnant and early lactating camels were similar to the reference range of dromedary camels (Faye and Bengoumi, 2018; Waziri *et al*, 2019; Islam *et al*, 2019; Martín-Barrasa *et al*, 2023).

The current study demonstrated significantly lower TEC and Hb in growing females than in dry ones, while PCV, MCH, MCV and MCHC exhibited no change. Conversely, many investigators reported significant changes in PCV, MCV, MCH and MCHC with age in camels (Al-Sultan, 2008; Elkhair and Elmgboul, 2015; Tharwat *et al*, 2015). Significant lower

values of TEC and Hb have been reported by Yagoub (1988), who concluded that camels aged one to five years had lower TEC values than camels aged more than five years. Furthermore, Al-Busadah and Osman (2000) found that age influenced the erythrocytes parameters, with younger camels exhibiting lower TEC and Hb values than the older camels. In contrast, young camels showed higher TEC and Hb values than the older ones (Hussein *et al*, 1992; Martín-Barrasa *et al*, 2023; Monaco *et al*, 2024). Moreover, camel calves had significantly lower values for TEC and Hb than lactating females (Al-Rammahi *et al*, 2016).

The pattern of erythrocytes parameters (higher TEC and Hb) observed during late pregnancy could be explained by the increased metabolic demand for oxygen consumption required for rapid foetal growth and lactogenesis and suggesting a potential adaptation to the increased metabolic demands during pregnancy and redistribution of resources towards lactation. Conversely, TEC and Hb manifested significantly lower values in late pregnant and early lactating camels compared to early lactating and dry ones (Hussein *et al*, 1992; Al-Busadah and Osman, 2000; Elkhair, 2019; Mohamed *et al*, 2021). Saeed *et al* (2011) reported lower TEC, Hb and PCV in pregnant than those in non-pregnant camels. Other investigators reported higher values of Hb, PCV, TEC, MCV and MCH in late pregnant camels compared to non-pregnant (Abd El-Salaam and Arafa, 2018; Abdul-Rahaman *et al*, 2018), whereas TEC and MCV were significantly lower in lactating camels than those in non-lactating (Mohamed *et al*, 2021). Significantly higher values of PCV, Hb and TEC were recorded in pregnant and lactating camels and the transition period (Getnet and Abebe, 2005; Tharwat *et al*, 2015; Abd-El-Rahman *et al*, 2017). The non-significant variations in PCV and haematological induces (MCV,




Fig 2. Serum protein capillary electrophoresis pattern of dry (a), late pregnant (b) and early lactating (c) camels.

MCH and MCHC) values recorded are comparable to those reported in late pregnant and early lactating camels and transition period (El Zahar *et al*, 2017; Elkhair and Minawy, 2018; Monaco *et al*, 2024).

The TLC in female camels corresponded with the reference range demonstrated in the literature for dromedary camels (Faye and Bengoumi, 2018; Waziri *et al*, 2019; Islam *et al*, 2019; Martín-Barrasa *et al*, 2023; Monaco *et al*, 2024). The TLC and eosinophils and basophils percentages did not change significantly in the growing females. The findings are comparable to previous results documented by many investigators in camels (Elkhair and Elmgboul, 2015; Omer *et al*, 2016; Ahmed, 2017). However, other investigators reported significant changes in TLC with age (Hussein *et al*, 1992; Al-Sultan 2008; Saeed and Hussein, 2008; Tharwat *et al*, 2015; Islam *et al*, 2019; Gaashan *et al*, 2020). Furthermore, TLC, eosinophils and basophils percentages did not differ significantly in late pregnant and early lactating animals; however, Axay *et al* (2017) reported marked leukocytosis in early lactating camels compared to late pregnant ones. Moreover, Al-Busadah and Osman (2000) reported that TLC of young camels was significantly higher compared to lactating camels. In other studies, TLC remained unchanged in late pregnant and lactating compared to dry camels (Ahmed, 2017; Elkhair and Minawy, 2018; Mohamed *et al*, 2021). The slight non-significant higher values of TLC recorded during early lactation compared to the dry females is attributed to neutrophilia recorded (Table 1).

The significant monocytosis obtained for the growing females contradicts the previous reports (Al-Sultan, 2008; Elkhair and Elmgboul, 2015; Omer *et al*, 2016). On the other hand, the significant neutrophilia and lymphocytosis observed during early lactation and late pregnancy, respectively, compared to the dry ones, potentially indicate an immune response to the stress of pregnancy, parturition and lactation, as well as the release of ACTH and cortisol (Getnet and Abebe, 2005; Omer *et al*, 2016; Ahmed, 2017; Mohamed *et al*, 2021). Neutrophilia and lymphocytosis during early lactation and late pregnancy were consistent with previous findings in camels (Muhammad *et al*, 2011; Tharwat *et al*, 2015; El Zahar *et al*, 2017; Ebissy *et al*, 2019; Elkhair, 2019).

The blood glucose concentration in camels, a key indicator of metabolic health, is comparable to the reference range established by Faye and Bengoumi (2018). The non-significant variation in glucose concentration in growing camels, specifically those between the ages of 1-5 years, is particularly

significant as it aligns with a previous study by Ghodsian *et al* (1978), which found no significant difference in camels up to five years old. However, it also revealed a contrasting trend, with many investigators reporting that young camels, typically under the age of 5 years, were more hyperglycemic than adults (Elias and Yagil, 1984; Faye and Mulato, 1991; Souilem *et al*, 1999; Ben Romdhane *et al*, 2003; Roba *et al*, 2023). These findings underscore the importance of age as a factor in blood glucose regulation in camels.

It is widely known that the blood glucose in camels increases with progressive pregnancy and decreases during the first two weeks of lactation (Kelanemer *et al*, 2015; Souilem *et al*, 1999). The current study found no significant differences in glucose concentration in late pregnant and early lactating camels. The results are consistent with previous findings reported in late pregnant, early lactating camels and the transition period, such as those by Tharwat *et al* (2015), El Zahar *et al* (2017), Ebissy *et al* (2019), Faraz *et al* (2021) and Mohamed *et al* (2021). The non-significant hypoglycemia reported in early lactating camels compared to late pregnant ones is most likely attributable to volume expansion combined with increased consumption of large amounts of blood glucose for lactose synthesis (Jainudeen and Hafez, 1992; Kaneko *et al*, 2008). Many investigators concluded that the potential synthesis of milk lactose was linked to increased glucose uptake by the mammary glands (Afshar and Fathi, 2012; Zhao, 2014).

It is well established that protein requirement increases with progressive pregnancy and the onset of lactation, as indicated by decreased maternal serum protein concentration (Muhammad *et al*, 2011; Saeed *et al*, 2011; Roba *et al*, 2023). Farm animals consume more amino acids from the maternal bloodstream for protein synthesis due to the progressive increase in foetal growth (Jainudeen and Hafez, 1994). Compared to dry ones, the considerable hyperproteinemia observed in late pregnant camels combined with marked variations in albumin and  $\alpha 1$ ,  $\alpha 2$  and  $\beta 1$ -globulins, indicates the critical metabolic changes during these stages required foetal growth and lactogenesis. The findings reveal unique metabolic changes in camels during late pregnancy and early lactation, such as hyperproteinemia and variations in albumin and globulin fractions, which are more pronounced in late pregnant and early lactating animals compared to dry females. These changes suggest a potential role of albumin and globulins in

regulating dynamic balance and immune response, respectively.

Camels in late pregnancy and early lactation revealed no substantial change in globulin fractions ( $\beta_2$ - and  $\gamma$ ) and A/G ratio, confirming the stability of these parameters during these stages. Generally, the significant changes in serum protein profile observed in late pregnancy ( $\uparrow$ TP and albumin ratio and  $\downarrow\alpha_1$ ,  $\alpha_2$  and  $\beta_1$ -globulins) followed by a marked hypoproteinemia in early lactating camels than those in the dry females reflects the maternal requirement of proteins for foetal growth or to cope with colostrum and milk synthesis and production and providing immunoglobulins. Many investigators explained the changes in serum protein profile during late pregnancy and early lactation by the increased maternal requirement of proteins for foetal growth, colostrum and immunoglobulins synthesis (Kaneko *et al*, 2008; Tharwat *et al*, 2015; Elkhair, 2019). In contrast, pregnant and lactating camels showed a marked decrease in serum protein parameters (Elkhair and Hartmann, 2014; Kelanemer *et al*, 2015; Tharwat *et al*, 2015; Elkhair, 2016; 2019; El Zahar *et al*, 2017; Elkhair and Minawy, 2018; Elkhair *et al*, 2018; Ebissy *et al*, 2019; Mohamed *et al*, 2021).

Serum protein capillary electrophoretic pattern is widely used to indicate health status and may act as crucial diagnostic and prognostic biomarkers for numerous pathogenic disorders (Alberghina *et al*, 2011; Yang *et al*, 2012). The current results validated the application of capillary electrophoresis for serum protein fractionation in female camels during different physiological states. Many researchers found that age and growth phase have a significant influence on serum proteins electrophoresis components: albumin,  $\alpha_1$ ,  $\alpha_2$ ,  $\beta_1$ ,  $\beta_2$ ,  $\gamma$ -globulins and A/G ratios in camels (Ahmadi-hamedani *et al*, 2014; Elkhair and Hartmann, 2014; Abdoslam *et al*, 2018). Serum protein capillary electrophoretic pattern obtained produced six fractions, including albumin and five globulin fractions, consistent with previous camel results (Chaudhary *et al*, 2003; Abdoslam *et al*, 2018). The progressive, non-significant increase in globulin fractions observed in dry adult females compared to the growing camels can provide an age-related explanation. The significant decrease in  $\alpha_1$ ,  $\alpha_2$  and  $\beta_1$  percentages during early lactation is probably due to the decrease in various  $\alpha$ -fraction molecules and the passage of  $\beta_1$ -fraction molecules from the bloodstream to the mammary glands for milk and colostrum biosynthesis (Kaneko *et al*, 2008). The significant increase in albumin fraction percentage combined

with a significant decrease in  $\alpha_1$ -,  $\alpha_2$ - and  $\beta_1$ -globulins has been observed in lactating camels, dairy cows and ewes (Piccione *et al*, 2011, 2012; Elkhair and Hartmann, 2014; Ebissy *et al*, 2019; Tóthová *et al*, 2018; Mohamed *et al*, 2021; Adam and Elkhair, 2023). In the present study,  $\gamma$ -globulins and A/G ratio showed non-significant changes during late pregnancy and early lactation. Conversely, significant higher values of  $\gamma$ -globulins accompanied by lower A/G ratio in lactating camels (Elkhair and Hartmann, 2014).

All animal species require mineral for growth, reproduction and lactation (Underwood and Suttle, 1999). The serum mineral profile obtained for  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$  and Pi concentrations is comparable to the reference values for camels (Elkhair, 2016; Faye and Bengoumi, 2018). In the growing females, the significant hypercalcemia, hypermagnesemia and hyperphosphatemia observed could be attributable to their higher mineral requirements for growth. This finding raises intriguing questions about the unique mineral needs of growing female dromedary camels and the potential implications for their health and development. Goff (2015) stated that  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$  and Pi were essential for blood and muscles functions, enzymatic reactions, cellular components synthesis and energy transferring molecules. Extensive research has consistently shown that young camels are hypercalcemic (Elias and Yagil, 1984; Rezakhani *et al*, 1997; Al-Busadah 2003, 2010; Ben Romdhane *et al*, 2003; Saeed *et al*, 2004; Barri *et al*, 2005; Tajik *et al*, 2015, Martín-Barrasa *et al*, 2023), hypermagnesemic (Barri *et al*, 2005; Tharwat *et al*, 2015; Elkhair, 2019; Martín-Barrasa *et al*, 2023) and hyperphosphatemic (Saeed *et al*, 2004; Al-Busadah, 2010; Elkhair, 2016; Martín-Barrasa *et al*, 2023) than adults. However, other researchers concluded that age had no remarkable influence on  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$  and Pi (Faye and Mulato, 1991; Saeed *et al*, 2004; Tajik *et al*, 2015; Tharwat *et al*, 2015; Elkhair, 2016).

The serum concentrations of  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$  and Pi in late pregnant and early lactating animals remained stable, a finding that aligns with previous reports in lactating camels (Singh *et al*, 2015; Tharwat *et al*, 2015; Ebissy *et al*, 2019). However, the observation of non-significant hypercalcemia and hypermagnesemia during these critical periods challenges the previously reported considerable hypocalcaemia and hypomagnesemia in camels. These conditions were attributed to excessive losses through urine and colostrum or impaired absorption of these minerals from the alimentary tract (Eltohamy *et al*, 1986; Kuria *et al*, 2006). The lack of statistical significance in serum



Pi during late pregnancy or early lactation also echoes the findings of several camel researchers (Khadjeh, 1998; Saeed *et al*, 2009; Ahmed, 2017; Faraz *et al*, 2021). However, a contrasting study by Mohamed *et al* (2021) suggests that hyperphosphatemia during early lactation is most likely caused by enhanced growth hormone (GH) activity combined with intestinal absorption and renal reabsorption of Pi, which stimulates GH synthesis by the alveolar cells of the mammary glands. This finding could have significant implications for our understanding of camel health and the management of late pregnancy and early lactation.

## Conclusion

Blood parameters were affected significantly by the age, growth phase, late pregnancy and early lactation in female camels reared under a semi-intensive system. Critical alterations in these parameters concerning the corresponding physiological state could be associated with growth requirements, intensive foetal growth and lactogenesis. The data could be employed to evaluate the metabolic profile of growing, pregnant and lactating camels and to develop efficient management strategies for females in a semi-intensive system.

## Abbreviations

Hb: Hemoglobin; PCV: Packed cell volume; TEC: Total erythrocytes count; MCV: Mean corpuscular volume; MCH: Mean corpuscular haemoglobin; MCHC: Mean corpuscular haemoglobin concentration; TLC: Total leukocytes count; Serum-[TP]: Serum total protein concentration;  $[Ca^{++}]$ ,  $[Mg^{++}]$  and  $[Pi]$ : calcium and magnesium and inorganic phosphate concentration.

## Funding

The research project was funded by the Ministry of Higher Education and Scientific Research (Sudan) and the Deanship of Scientific Research, Vice Presidency for Graduate Studies and Scientific Research, King Faisal University, Saudi Arabia (Grant No. 5, 960). Nawal M. Elkhair was a recipient of the research fund. The funder had no role in study design, data collection and analysis, publication decisions, or manuscript preparation.

## Acknowledgements

The author would like to thank and acknowledge the Deanship of Scientific Research, Vice Presidency for Graduate Studies and Scientific Research, King Faisal University (Saudi Arabia) and

the technician team of the Department of Physiology, Faculty of Veterinary Medicine, University of Khartoum (Sudan) for financial support and laboratory assistance.

## Conflict of Interest

The author declares no conflict of interest.

## Data Availability data Statement

The data is available in the manuscript and from the corresponding author upon reasonable request.

## References

- Abd El-Salaam AM and Arafa MM. Postpartum haematological, biochemical, mineral and hormonal changes in blood of Maghrebian she-camels with different parity orders under Egyptian condition. *IOSR Journal of Agriculture and Veterinary Science*. 2018; 11(2):68-78.
- Abd-El-Rahman HMA, Ibrahim MA and Elmetwaly HA. Hormonal profile, antioxidant status and some biochemical parameters during pregnancy and periparturient period in dromedary she camel. *Egypt. Egyptian Journal of Veterinary Sciences*. 2017; 48(2):81-94 <https://doi.org/10.21608/ejvs.2017.2040.1022>
- Abdoslam O, Bayt-Almal M, Almghrbe A and Algriany O. Serum protein electrophoretic pattern in one-humped camels (*Camelus dromedarius*) in Tripoli, Libya. *Open Veterinary Journal*. 2018; 8:1-4. <https://doi.org/10.4314/ovj.v8i1.1>
- Abdul-Rahaman A, Mnati A and Mohammed WM. Effect of different gestation periods on some physiological aspects of Iraqi female dromedary camel (*Camelus dromedarius*). *Iraqi Journal of Agricultural Sciences*. 2018; 49(4):494-707. <https://doi.org/10.36103/ijas.v49i4.80>
- Adam AA and Elkhair NM. Serum protein capillary electrophoretic pattern during lactation period in camels (*Camelus dromedarius*). *Comparative Clinical Pathology*. 2023; 32:599-605. <https://doi.org/10.1007/s00580-023-03468-z>
- Afshar M and Fathi H. Lactose in ruminants feeding: A review. *Annals of Biological Research*. 2012; 3(1):645-650.
- Ahmadi-hamedani M, Ghazvinian K, Kokhaei P, Barati M and Mahdavi A. Comparison of effects of age and sex on serum protein electrophoretic pattern in one-humped camels (*Camelus dromedarius*) in Semnan, Iran. *Open Veterinary Journal*. 2014; 4:4-8. PMID: PMC4629590
- Ahmed MH. Effects of selenium and vitamin E injection during transition period on physiological performance of camels (*Camelus dromedarius*) and their neonates reared under semi-intensive system. Thesis (MVSc), University of Khartoum, Sudan. 2017.
- Ahmed MH and Elkhair NM. Effects of the physiological status and selenium and vitamin E injection during transition period on serum proteins parameters in camels (*Camelus dromedarius*) reared under semi-intensive system. *International Journal of Research in Medical and Basic Sciences*. 2019; 5(1):14-25.

- Alberghina D, Giannetto C, Vazzana I, Ferrantelli V and Piccione G. Reference intervals for total protein concentration, serum protein fractions and albumin/globulin ratios in clinically healthy dairy cows. *Journal of Veterinary Diagnostic Investigation*. 2011; 23:111-114. <https://doi.org/10.1177/104063871102300119>
- Al-Busadah KA and Osman TEA. Haematological parameter of adult dry, lactating and camel calves in Saudi Arabia. *Pakistan Journal of Biological Sciences*. 2000; 3(10):1749-1751. <https://doi.org/10.3923/pjbs.2000.1749.1751>
- Al-Busadah KA. Some aspects of calcium and magnesium metabolism in camel calves. *Journal of Camel Practice and Research*. 2003; 10(2):111-114.
- Al-Busadah KA. Serum concentration of aluminum, calcium, magnesium and phosphorous in camels. *Scientific Journal of King Faisal University*. 2010; 11(1):161-167.
- Al-Rammahi HM, Al-Jebory HA and Abed Al-Sattar H. Some normal haematological values of Arabian camels reared in western desert of Al-Najaf governorate/Iraq. *MRVSA 5 (Special issue) 1<sup>st</sup> Iraqi Colloquium on Camel Diseases and Management*, 2016; pp 34-38.
- Al-Sultan SI. Studies on haematological and certain serum biochemical values in young Magaheim dromedary camel at Al-Ahsa province. *Veterinary Research*. 2008; 2(3-4): 34-37. <https://medwelljournals.com/abstract/?doi=vr.2008.34.37>
- Arifin WN and Zahiruddin WM. Sample size calculation in animal studies using resource equation approach. *Malaysian Journal of Medical Sciences*. 2017; 24(5):101-105. <https://doi.org/10.21315/mjms2017.24.5.11>
- Atta A, El-Shourbagy I, Gawish H, El-Metwaly H, Younis F and Salem M. Farming and pastoral systems shape the physiological and immunological indices of the dromedary she-camels. *Egyptian Journal of Cancer and Biomedical Research*. 2021; 5(1):49-63. <https://doi.org/10.21608/jcbr.2020.49275.1092>
- Axay J, Haque F, Lateef A, Patel A, Patel P and Bhalakiya N. Study on blood metabolites and leukocyte indices of Kutchi camels during different stages of lactation. *Journal of Animal Health and Production*. 2017; 5(3):92-96. <http://dx.doi.org/10.17582/journal.jahp/2017/5.3.92.96>
- Ayoub MA, El Khoully AA and Mohamed TM. Some haematological and biochemical parameters and steroid hormone levels in the one-humped camel during different physiological conditions. *Emirates Journal of Food Agriculture*. 2003; 15(1):44-55. <https://doi.org/10.9755/ejfa.v15i1.5281>
- Barri MES, Al-Busadah KA and Homeida AM. Comparative calcium and magnesium status in adult and young camels. *Scientific Journal of King Faisal University*. 2005; 6:151-158.
- Ben Romdhane S, Romdane MN, Feki M, Sanhagi H, Kaabachi N and M'Bazaa A. Blood biochemistry parameters in dromedary (*Camelus dromedarius*). *Revue de Médecine Vétérinaire*. 2003; 154(11):695-702.
- Caraway WT. Carbohydrates. In: Tietz NW. *Fundamentals of Clinical Chemistry*. 1987; Saunders, Philadelphia.
- Chaudhary ZI, Iqbal J and Rashid J. Serum protein electrophoretic pattern in young and adult camels. *Australian Veterinary Journal*. 2003; 81(10):625-626 <https://doi.org/10.1111/j.1751-0813.2003.tb12508.x>
- Chromya V, Svoboda V and Stepanova I. Spectrophotometric determination of magnesium in biologic fluids with xylidyl blue. *Biochemical Medicine*. 1973; 7(2):208-217. [https://doi.org/10.1016/0006-2944\(73\)90075-6](https://doi.org/10.1016/0006-2944(73)90075-6)
- Ebissy EA, El-Sayed AA and Mohamed RH2. Haematological and biochemical profile in female camels (*Camelus dromedarius*) during the transition period. *Slovenian Veterinary Research*. 2019; 56(22):571-577. <https://doi.org/10.26873/SVR-794-2019>
- El Zahar H, Zaher H, Alkablawy A, Al Sharifi S and Swelum A. Monitoring the changes in certain haematological and biochemical parameters in camels (*Camelus dromedarius*) during postpartum period. *Journal of Fertility Biomarkers*. 2017; 1(1):47-54. <https://doi.org/10.14302/issn.2576-2818.jfb-17-1595>
- Elias E and Yagil R. Haematological and serum biochemical values in lactating camels and their newborn. *Refuah Veterinary Journal*. 1984; 41:7-13.
- Elkhair NM and Hartmann H. Serum protein capillary electrophoretic pattern in camels (*Camelus dromedarius*): influence of age and sex. *Journal of Life Sciences*. 2014; 8(1):78-81.
- Elkhair NM and Elmgboul WE. Haematological parameters of neonate camels (*Camelus dromedarius*) during the first week of life. *University of Khartoum Journal of Veterinary Medicine and Animal Production*. 2015; 6(2):103-110.
- Elkhair NM, Ahmed MH and Alameen AO. Monitoring of certain serum biochemical parameters during transition period in camels (*Camelus dromedarius*). *University of Khartoum Journal of Veterinary Medicine and Animal Production*. 2018; 9(1):27-35.
- Elkhair NM and Minawy AA. Effect of the physiological status on the haematological parameters of female camels (*Camelus dromedarius*). *University of Khartoum Journal of Veterinary Medicine and Animal Production*. 2018; 9(2):154-162.
- Elkhair NM. Influence of age and sex on certain serum biochemical parameters of dromedary camels (*Camelus dromedarius*). *Nova Journal of Medical and Biological Sciences*. 2016; 5(3):1-8.
- Elkhair NM. Monitoring of the haematological parameters in camels (*Camelus dromedarius*) during late pregnancy and early lactation. *EC Veterinary Science*. 2019; 4(8): 675-680.
- Eltohamy MM, Salama A and Youssef AEA. Blood constituents in relation to the reproductive state in she-camel (*Camelus dromedarius*). *Beitrage zur Tropischen Landwirtschaft und Veterinarmedizin*. 1986; 24:425-430. <https://pubmed.ncbi.nlm.nih.gov/3566702>
- Faraz A and Waheed A, Tauqir NA, Mirza RH, Ishaq HM, Nabeel MS. Effect of pregnancy on blood biochemical profile of semi-intensive kept Marecha (*Camelus dromedarius*) Camel. *International Journal of*

- Agriculture and Biology. 2021; 25:241-248. <https://doi.org/10.17957/IJAB/15.1662>
- Faye B and Mulato C. Facteurs de variation des paramètres protéo-énergétiques, enzymatiques et minéraux dans le plasma chez le dromadaire de Djibouti. *Revue d'élevage et de médecine vétérinaire des pays tropicaux*. 1991; 44:325-334. <https://doi.org/10.19182/remvt.9174>
- Faye B and Bengoumi M. Haematology. In: *Camel Clinical Biochemistry and Haematology*, Springer International Publishing. 2018; pp 13-45.
- Gaashan MM, Al-Mubarak AIA and Hussien, J. Leukocyte populations and their cell adhesion molecules expression in newborn dromedary camel calves. *Veterinary World*. 2020; 13:1863-1869. <https://doi.org/10.14202/VETWORLD.2020.1863-1869>
- Gamst O and Try K. Determination of serum-phosphate without deproteinisation by ultraviolet spectrophotometry of the phosphomolybdic acid complex. *Scandinavian Journal of Clinical Laboratory Investigation*. 1980; 40(5):483-486. <https://doi.org/10.3109/00365518009101872>
- Getnet AM and Abebe W. The influence of late pregnancy and excitement on blood parameters of Issa type dromedaries in Eastern Ethiopia. *Israel Journal of Veterinary Medicine*. 2005; 60:117-120. <https://studylib.net/doc/7706468>
- Ghodsian I, Nowrouzian I and Schels HF. A study of some haematological parameters in the Iranian camel. *Tropical Animal Health and Production*. 1978; 10:109-110. <https://doi.org/10.1007/BF02235320>
- Gindler EM and King JD. Rapid colorimetric determination of calcium in biologic fluids with methyl thymol blue. *American Journal of Clinical Pathology*. 1972; 58:376-382. <https://doi.org/10.1093/ajcp/58.5.376>
- Goff JP. Minerals, bones and joints. In: Reece, W.O., Erickson, H.H., Goff, J.P., Uemura, E.E. (Eds.), *Duke's Physiology of Domestic Animals*. 13th ed. John Wiley and Sons Inc. 2015; pp 568-578.
- Hussein MF, Liman MM, Mogawer HH, Bakkar MN and Garel Nabi AR. The haematology of growing camels (*Camelus dromedarius*) during the first year of life. *Asian-Australasian Journal of Animal Sciences*. 1992; 5(3):519-525. <https://doi.org/10.5713/ajas.1992.519>
- Hussein MF, Salah MS, Mogawer HH and Gar ElNabi AR. Effect of lactation on the haemogram and certain blood constituents of the dromedary camel. *Journal of Applied Animal Research*. 1992; 1:43-50. <https://doi.org/10.1080/09712119.1992.9705907>
- Inc SPSS. IBM SPSS Statistics Base 20, SPSS Inc. Chicago, IL, USA. 2011.
- Islam S, Ferdous J, Rahman MK, Akter S, Hassan MM and Islam A. Reference values for haematological and serum biochemical parameters of dromedary camel (*Camelus dromedarius*) in sub-tropical climate of Bangladesh. *Advances in Animal and Veterinary Sciences*. 2019; 10:232-237. <https://doi.org/10.17582/journal.aavs/2019/7.4.232.237>
- Jain NC. *Essentials of Veterinary Haematology*, Second ed. Lea & Febiger, Philadelphia, USA. 1998; pp 65-68.
- Jainudeen MR and Hafez ESE. Gestation, prenatal physiology and parturition. In: Hafez, E.S.E., (Ed.), *Reproduction in Farm Animals*. Lippincott, Williams and Wilkins. 1994; pp 247-283.
- Jalali SM, Hasani YN, Darabifard A and Mavadati AH. A study of hematologic and biochemical profile in female dromedary camels during the breeding and non-breeding seasons. *Comparative Clinical Pathology*. 2018; 27(3):655-659. <https://doi.org/10.1007/s00580-018-2645-1>
- Kaneko JJ, Harvey JW and Bruss ML. *Clinical Biochemistry of Domestic Animals*, Sixth ed. Elsevier Academic Press, London. 2008.
- Kelanemer R, Antoine-Moussiaux N, Moula N, Abu-median A, Hanzen C and Kaidi R. Effect of nutrition on reproductive performance during the peripartum period of female camel (*Camelus dromedarius*) in Algeria. *Journal of Animal Veterinary Advances*. 2015; 14:192-196. <https://doi.org/10.36478/javaa.2015.192.196>
- Khadjeh GH. Concentration of serum electrolytes in pregnant and non-pregnant camels (*Camelus dromedarius*). *Journal of Camel Practice and Research*. 1998; 5(2):289-290.
- Kuria SG, Gachui CK, Wahome RG and Wanyoike MM. Mineral profile in the plasma of free ranging camels (*Camelus dromedarius*) in Kenya. *Indian Journal of Animal Sciences*. 2006; 76(12):1068-1070. <http://www.lrrd.org/lrrd25/8/kuri25149.htm>
- Martín-Barrasa JL, Tejedor-Junco MT, Cabrera S, Morales M, Melián A and Corbera JA. Haematological and biochemical blood reference values for Canary Island camels (*Camelus dromedarius*), an endangered dromedary species. *Saudi Journal of Biological Sciences*. 2023; 30(6):103677. <https://doi.org/10.1016/j.sjbs.2023.103677>
- Mead R. *The design of experiments: Statistical Principles for Practical Applications*. Cambridge University Press, Cambridge. UK.1988.
- Mohamed RH, Ali F and Nour SY. Comparative studies on ovarian hormones and hematobiochemical profile of lactating and non-lactating she-camels (*Camelus dromedarius*). *Comparative Clinical Pathology*. 2021; 30:415-420. <https://doi.org/10.1007/s00580-021-03232-1>
- Monaco D, Mariella J, Fusi J, Osman TK, Abdel Rauf A and Probo M. Haematological reference intervals in newborn dromedary calves in the first week after birth: Age and sex-related variations. *Research in Veterinary Science*. 2024; 169:105172. <https://doi.org/10.1016/j.rvsc.2024.105172>
- Muhammad BF, Aliyu D, Njidda AA and Madigawa IL. Some haematological, biochemical and hormonal profile of pregnant and non-pregnant she-camels (*Camelus dromedarius*) raised in a Sudan savanna zone of Nigeria. *Journal of Camel Practice and Research*. 2011; 18(1):73-77.
- Omer SA, Khougali SME, Agab H and Samad GHA. Studies on some biochemical and haematological indices of Sudanese camels (*Camelus dromedarius*) 2016). [http://www.sustech.edu/staff\\_publications/20090615152717900.pdf](http://www.sustech.edu/staff_publications/20090615152717900.pdf)



- Onasanya GO, Oke FO, Sanni TM and Muhammad AI. Parameters influencing haematological, serum and biochemical references in livestock animals under different management systems. *Open Journal of Veterinary Medicine*. 2015; 5:181-189. <https://doi.org/10.4236/ojvm.2015.58025>
- Piccione G, Messina V, Schembari A, Casella S, Giannetto C and Alberghina D. Pattern of serum protein fractions in dairy cows during different stages of gestation and lactation. *Journal of Dairy Research*. 2011; 78:421-425. <https://doi.org/10.1017/S0022029911000562>
- Piccione G, Alberghina D, Marafioti S, Giannetto C, Casella S, Assenza A and Fazio F. Electrophoretic serum protein fraction profile during the different physiological phases in Comasina Ewes. *Reproduction in Domestic Animal*. 2012; 47:591-595. <https://doi.org/10.1111/j.1439-0531.2011.01925.x>
- Rezakhani A, Habibabadi, SN, Ghogh MM. Studies on normal haematological and biochemical parameters of Turkmen camel in Iran. *Journal of Camel Practice and Research*. 1997; 4(1):41-44
- Roba WJ, Yisehak KK, Asrat AG. Effects of season, productive state, age and agro-ecology on blood biochemical characteristics of dromedary camels (*Camelus dromedarius*) in natural browsing environment. *Asian Journal of Dairy and Food Research*. 2023; DOI: 10.18805/ajdfr.DRF-330
- Saeed A, Hussain MM, Khan IA, Chand G, El-Yousuf RA. Effect of sex and age on blood biochemical profile in camel. *Journal of Camel Practice and Research*. 2004; 11(1):73-77.
- Saeed A and Hussein MM. Change in normal haematological values of camels (*Camelus dromedarius*): Influence of age and sex. *Comparative Clinical Pathology*. 2008; 17:263-266. <https://doi.org/10.1007/s00580-008-0736-0>
- Saeed A, Khan IA and Hussein MM. Change in biochemical profile of pregnant camels (*Camelus dromedarius*) at term. *Comparative Clinical Pathology*. 2009; 18:139-143. <https://doi.org/10.1007/s00580-008-0763-x>
- Saeed A, Khan IA and Hussein MM. Change in haematological profile of pregnant camels (*Camelus dromedarius*) at term. *Comparative Clinical Pathology*. 2011; 20:53-55. <https://doi.org/10.1007/s00580-009-0933-5>
- Singh S, Dedar RK, Legha RA, Bala PA and Patil NV. Minerals and electrolytes profile in Lactating and pregnant Indian camels. *Journal of Camel Practice and Research*. 2015; 22(1):121-124. <https://doi.org/10.5958/2277-8934.2015.00019.3>
- Souilem O, Chine O, Alguemi C and Gogny M. Glycaemia in the camel (*Camelus dromedarius*) in Tunisia: preliminary results. *Revista de Medicina Veterinaria*. 1999; 150(6):537-542.
- Tajik J, Sazmand A, Hekmatimoghaddam S, Rasooli A and Mohammadzadeh YA. Serum concentrations of some ions in clinically healthy camels (*Camelus dromedarius*). *Eurasian Journal of Veterinary Sciences*. 2015; 31(4):204-208. [https://www.eurasianjvetsci.org/pdf/pdf\\_EJVS\\_1099.pdf](https://www.eurasianjvetsci.org/pdf/pdf_EJVS_1099.pdf)
- Tharwat M, Ali A, Al-Sobayil F, Selim L and Abbas H. Hemato-biochemical profile in female camels (*Camelus dromedarius*) during the periparturient period. *Journal of Camel Practice and Research*. 2015; 22(1):101-106. <https://doi.org/10.5958/2277-8934.2015.00016.8>
- Tharwat M, Al-Sobayil F and El-Deeb W. Haematology and serum biochemistry profiles in dromedary camel calves during the first month of age. *Journal of Camel Practice and Research*. 2015; 22:107-112. <https://doi.org/10.5958/2277-8934.2015.00017.X>
- Tóthová C, Nagy O, Nagyová V and Kováč G. Serum protein electrophoretic pattern in dairy cows during the periparturient period. *Journal of Applied Animal Research*. 2018; 46:33-38. <https://doi.org/10.1080/09712119.2016.1256293>
- Underwood EJ and Suttle NF. *The Mineral Nutrition of Livestock*, Third ed. Center for Agriculture and Biosciences International, United Kingdom. 1999.
- Waziri A, Hassan SU and Igbokwe IO. Haematological reference values of dromedary camels in northern Nigeria. *Comparative Clinical Pathology*. 2019; 28:1769-1777. <https://doi.org/10.1007/s00580-019-03018-6>
- Weichselbaum IE. An accurate and rapid method for the determination of proteins in small amounts of blood serum and plasma. *American Journal of Clinical Pathology*. 1946; 16(3):40-49. [https://doi.org/10.1093/ajcp/16.3\\_ts.40](https://doi.org/10.1093/ajcp/16.3_ts.40)
- Weiss DJ and Wardrop KJ. *Schalm's Veterinary Haematology*, Sixth ed. A John Wiley and Sons, Ltd., Publication. 2010.
- Werner J. *Biomathematik und Medizinische Statistik*. Urban & Schwarzenberg, München, Germany. 1992; pp 249-259.
- Yagoub IA. Studies on haematological parameters and some biochemical blood constituents of Sudanese camels (*Camelus dromedarius*). *Acta Veterinaria*. 1988; 38:99-106.
- Yang YX, Wang JQ, Bu DP, Li SS, Yuan TJ, Zhou LY, Yang JH and Sun P. Comparative proteomics analysis of plasma proteins during the transition period in dairy cows with or without subclinical mastitis after calving. *Czech Journal of Animal Science*. 2012; 57:481-489. <https://doi.org/10.17221/6348-CJAS>
- Zhao FQ. Biology of glucose transport in the mammary gland. *Journal of Mammary Gland Biology and Neoplasia*. 2014; 9:3-17. <https://doi.org/10.1007/s10911-013-9310-8>

# RADIOGRAPHIC EVALUATION AND SOFT PALATE DIVERTICULECTOMY OF IMPACTED DULAA IN TWO CAMELS (*Camelus dromedarius*)

Ruchi Patwa, Mahendra Tanwar, P. Bishnoi, A.K. Bishnoi, Shivangi Diwedi and Balkrishan Yadav

Department of Veterinary Surgery and Radiology, College of Veterinary and Animal Science,  
Rajasthan University of Veterinary and Animal Sciences, Bikaner-334 001, Rajasthan

## ABSTRACT

Two camels of impacted dulaa (soft palate diverticulum) were presented with a history of inability of soft palate protrusion, anorexia and dyspnoea. Clinical examination with radiographic evaluation was done. Clinical examination of throat region revealed pain and dyspnoea on palpation. Lateral radiographic view of cranial cervical region was taken and radiograph showed increased radiodensity of irregular soft tissue with scattered gaseous radiolucency. Complete clinical and radiographic examination revealed accumulation of food and pus in the soft palate and surgical resection of impacted dulaa was planned. Soft palate diverticulectomy was done by intraoral approach under general anaesthesia. Both the animals recovered uneventfully. The conclusion of the clinical case report was that, in case of impacted dulaa, pharyngeal region radiography helps delineating the larynx, proximal trachea, oesophagus, naso and oropharynx. Enlarged size and radiodense dulaa (impacted by feed material) were seen radiographically and helped planning the surgery.

**Key words:** *Camelus dromedarius*, impacted dulaa, radiographic evaluation, soft palate

The soft palate diverticulum (dulaa) is an oroventral projection of soft palate and is injured by the camel's own canine teeth or by other camels while fighting, or by thorns and wires commonly during breeding season. The signs may include oedema, perforation and haematoma depending upon the causal agent and degree of trauma (Gahlot and Chouhan, 1992; Tanwar *et al*, 2016; Kumar *et al*, 2017). The food sometimes becomes impacted in the dulaa. The size of impacted dulaa may sometimes go up to the size of a football. The mass is very hard to touch. In the case of larger mass, the resection of the organ is required. Tracheotomy is also before resection (Choudhry *et al*, 2017). Clinical findings and surgical exteriorisation of dulaa has been reported following various types of dulaa injuries (Sadan and El-Shafaey, 2020). Radiography is scarcely reported for the diagnosis of injured dulaa in camels. The present study described radiographic evaluation of impacted dulaa and its surgical resection in two dromedary camels.

## Materials and Methods

Two camels aged 7 and 12 years, were presented at Veterinary Clinical Complex, Rajasthan University of Veterinary and Animal Sciences,

Bikaner, with the history of anorexia, dyspnoea, inability of dulaa protrusion and swelling at cervical region. Clinical examination along with aseptic fine needle aspiration of swollen area of throat region and radiographic evaluation was done. Clinical examination revealed pain and swelling on palpation and while straightening the neck (Fig 1). Complete clinical and radiographic examination revealed accumulation of food and pus in the soft palate and surgical resection of impacted dulaa or soft palate diverticulectomy was planned. Animals were kept off feed and water for 24 hours. Camels were restrained in sternal recumbency and surgical procedure was performed under general anaesthesia including (xylazine @ 0.2 mg /kg and ketamine @ 1mg / kg body weight). After induction of anaesthesia, a gag was placed between the upper and lower jaws and an attempt was made to pull the soft palate out of the oral cavity using a hook while the assistant was instructed to massage the proximal area of the neck with the palm towards the oral cavity and to squeeze the neck region but all went in vain. The surgeon's hand was inserted inside the oral cavity till the pharyngeal region and the distended soft palate was pulled out manually after removing the adhesions by holding the organ with a towel.

SEND REPRINT REQUEST TO RUCHI PATWA [email: ruchipatwa611@gmail.com](mailto:ruchipatwa611@gmail.com)

The exteriorised soft palate was found gangrenous and the pus was drained out along with the feed material (Fig 2A&B). The soft palate was excised using long straight Mayo scissors by pulling it outwards and twisting it with a towel. The oral cavity was flushed daily with light potassium paramanganate solution for 5-7 day. Postoperatively streptopenicillin 7.5 g for 5 days and meloxicam @ 0.3 mg/kg body weight for 3 days were administered intramuscularly. The camels were given soft fodder for two weeks.



**Fig 1.** Swelling and pain on palpation and while straightening of neck.

## Results and Discussion

Clinical signs, dyspnoea, i.e. pain and swelling were suggestive of severe inflammation throat region, even were straightening the neck. Aseptic fine needle aspiration of swelling at throat region revealed straw coloured fluid which was suggestive of presence of inflammatory fluid together with trapped feed. Lateral radiograph of cranial cervical region in both cases showed increased radiodensity of irregular soft tissue with scattered gaseous radiolucency close the epiglottis (Fig 3). Oral cavity examination under general anaesthesia found impacted soft palate with feed material. Soft palate was found gangrenous and contained pus (Fig 4). Soft palate diverticulectomy was performed and the camels recovered uneventfully (Fig 5).

Dulaa gets injured with its own teeth, biting of other camel and by external trauma (Gahlot, 2000 and Tanwar *et al*, 2016). Injury occurs commonly during rut season and injured dullla either hangs out or remain entrapped *in situ*. The affected animals usually show discomfort when swallowing, dysphagia, dyspnoea and are unable to extrude the dulaa due to lacerated wounds, gangrene, ulcers, abscesses or impaction with feed material (Ramadan, 2013). In animals of present study, dysphagia, dyspnoea and pain on palpation at throat region and while





**Fig 2A.** Gangrenous soft palate (arrow).



**Fig 2B.** Pus drained into oral cavity.






**Fig 3.** Lateral radiographic view of cranial cervical region was taken and radiograph showed increased radiodensity of irregular soft tissue with scattered gaseous radiolucency (arrow) in the camels.



**Fig 4.** Gangrenous soft palate.



**Fig 5.** Surgical wound after 15 days of excision of soft palate.

straightening the neck were observed. Rollefson *et al* (2001) also reported inability of dulaa ballooning as it remains entrapped condition or stays hanging outside the mouth. In present case report, the dulaa was entrapped and camel couldn't take feed and water in both the cases.

Radiography can be used for detection of impacted dulaa in pharyngeal region when the dulaa cannot be extruded by the animal. However, when

the dulaa gets impacted with feed particles its size and density increases. It is pushed caudally to press against the epiglottis thus interfering with breathing (Ramadan, 2013). In present clinical case report dulaa was impacted with feed material and pus which appeared as radiopaque soft tissue mass in cranial cervical region radiography (Fig 3). Sadan *et al* (2023) reported that radiography is advanced non-invasive diagnostic technique for the diagnosis of oesophageal

obstruction in addition to exploratory surgery to confirm the condition. Dorsoventral and lateromedial standard radiographs of affected camels were obtained under light sedation and radiographs were subjectively interpreted before surgical interventions. Moreover, successful surgical resection of the dulaa could be used as curative treatment of such disorders in camels (Sadan *et al*, 2020).

Ramadan (2013) also secured camels in sternal position and sedated with 2% xylazine hydrochloride (0.2 mg/kg body weight) together with 10% ketamine hydrochloride at the dose of 1 mg/kg body weight given, intravenously for soft palate resection (Ramadan, 2013) and local infiltration analgesia using lidocaine HCL 2% which was infiltrated at the root of the dulaa (Al-Sobayil and Ahmed, 2011; Ramadan, 2013). Similarly, in present case report xylazine @ 0.2 mg /kg body weight and ketamine @ 1mg / kg body weight were used in soft palate diverticulectomy without local anaesthesia in both cases.

In present case report resection of soft palate was performed by intraoral route which is simple, less time consuming and doesnot subject the animal to the risk of general anaesthesia. However, Ramadan (2013) described resection of soft plate by pharyngostomy, where in dulla was pulled out through pharyngeal opening between intermandibular space. A temporary tracheotomy tube or a suitable endotracheal tube was inserted down the trachea in pharyngostomy approach.

In present study, soft palate found was gangrenous and impacted with feed material. Similar injuries of soft palate leading to gangrene and haematoma of the soft palate have been reported previously (Gahlot and Chouhan, 1992; Sadan and El-Shafaey, 2020).

In present clinical case report was dulaa resected after pulling the soft palate out with a cotton cloth and two long artery forceps were used to hold and crush the major blood vessels close to the origin of the dulaa. A purse string suture using polydioxanone no. 2 was applied approximately 1 cm distal to the location of the artery forceps. The dissection was performed distal to the suture. The

dulaa was removed and the stump was closed by cushioning suture pattern using polydioxanone no. 2. Al-Sobayil and Ahmed (2011) stated that in some cases, interrupted, overlapping horizontal mattress suture pattern was used before amputation and the wound was further closed by simple continuous sutures. The severed edges were cauterised with strong potassium permanganate solution. Haemostasis occurred in couple of minutes (Gharu *et al*, 2016).

## Conclusion

The impacted dulaa can be diagnosed early with radiography which can help in deciding the extent of pathology before proceeding for its surgery.

## References

- Al-Sobayil FA and Ahmed AF. Surgery of the injured dulla in dromedary camels (*Camelus dromedarius*). Iranian Journal of Veterinary Surgery. 2011; 6(1-2):17-22.
- Chaudhry Z. Camel and its Disease. 1<sup>st</sup> Ed. Lap Lambert Academic Publishing, Mauritius. 2017; pp 6-42.
- Gahlot TK and Chouhan DS. Camel Surgery. Gyan Prakash Mandir, Bikaner, India. 1992; pp 47-52.
- Gahlot TK. Selected Topics on Camelids. The Camelid Publishers. Bikaner, India. 2000; 318-319.
- Gharu S, Singh D, Khosa JS, Kachwaha, K, Bamniya MK Qureshi SM and Gahlot TK. Surgical management of entrapped soft palate in camels (*Camelus dromedarius*). Indian Veterinary Journal. 2016; 93(11):67-68.
- Kumar P, Purohit NR and Gahlot TK. Clinical study on surgical affections of soft palate in camels (*Camelus dromedarius*). Intas Polivet. 2017; 18(1):222-225.
- Ramadan RO. Surgery of the dulaa in the camel (*Camelus dromedarius*). Journal of Agricultural and Marine Sciences. 2013; 18:25-29.
- Rollefson KI, Mundy P and Mathias E. A Field Manual of Camel Diseases. ITDG Publishing. London, UK. 2001; pp 106-107.
- Sadan M and El-Shafaey ES. Diagnosis and surgical repair of soft palate (dulla) injuries in camel (*Camelus dromedarius*). Journal of Camel Health. 2020; 1(2):38-42.
- Sadan M, El-Khodery S, Almatroodi S, Alsobayi F and El-Shafaey ES (2023). Diagnosing and treating oesophageal obstruction in camels (*Camelus dromedarius*). Veterinary World. 2023; 16(4):735.
- Tanwar Mahendra, Bishnoi P, Palecha S, Jhirwal SK and Singh S. Soft palate excision in camel (*Camelus dromedarius*)-report of 8 cases. Intas Polivet. 2016; 17(2):567-568.

# WEIGHT, GROWTH AND GENETICS IN SHAMI CAMELS (*Camelus dromedarius*)

Khaled A Al-Najjar<sup>1</sup>, Moammar Dayoub<sup>2</sup>, Azzam N. Al-Yacoub<sup>3</sup> and Widya P.B. Putra<sup>4</sup>

<sup>1</sup>General Commission for Scientific Agricultural Research, GCSAR, 12573, Damascus, SYRIA

<sup>2</sup>Turku University, Faculty of Technology, Computing Department, 20014, Turku, FINLAND

<sup>3</sup>Umm Al-Qura University, Faculty of Applied Science, Biology Department, 24382, Makkah, SAUDI ARABIA

<sup>4</sup>Research Centre for Applied Zoology-National Research and Innovation Agency Bogor-Jakarta Rd. Km. 46 Cibinong, Bogor, West Java, 16911, INDONESIA

## ABSTRACT

This study examines Shami camel calf rearing in semi-arid regions of Syria, focusing on weight and growth (birth to 1 year), heritability and breeding values of birth weight for breeding purposes. Birth weight, weight at 12 months and growth rate from birth to 12 months differed depending on the birth year. However, weight at 6 and 9 months and growth from birth to these ages did not show significant variation. The differences in weight at 6 and 12 months and growth from birth to 6 and 12 months, were highly significant according to calf sex, while differences at birth, at 9 months and in growth from birth to 9 months were not significant. Shami camel weights were correlated across growth stages, especially between weight at 6 months and growth rate from birth to 6 and 9 months. Male Shami camel calves and calves born to multiparous she-camels had slightly higher birth weights. Sire affected birth weight, with significant variation among them. The high heritability (0.80) of birth weight in Shami camels was found. Male camels were ranked according to the estimated breeding value of their birth weight, with the top sire (ID 126) expected to produce heavier calves. This study found that birth weight and growth rates varied by birth year and sex, emphasising the importance of Shami camel male selection in breeding programmes aimed at improving weights as early growth strongly influences later weight gain.

**Key words:** Breeding values, growth rates, heritability, live weights, Shami camel

The distinctive single hump of the Shami camel is adapted for fat storage, complemented by features like long legs, flat feet, bushy eyebrows, thin nostrils, thick lips and efficient water retention mechanisms (Osterloff, 2024).

Research estimates the Shami camel's growth and predicted weights from birth to 4 years of age. Genetic studies have identified unique variations that enhance water conservation, metabolism and overall body weight, improving their survival in desert conditions (Ali *et al*, 2019; Joana *et al*, 2021).

Comprehensive knowledge of the species' weights, growth rates and genetic characteristics is essential for developing effective breeding programmes and conservation strategies. This supports genetic enhancement and custom breeding, securing Shami camels' sustainable future (Fraser, 2008; Balasundaram *et al*, 2023).

Genetic factors influencing growth patterns include traits like lean meat and lower fat content, with younger Shami camels producing superior meat

quality (Al-Owaimer *et al*, 2014; Al-Saiady *et al*, 2015; Faraz *et al*, 2020; Suliman *et al*, 2020).

Research into camel breeding and management provides insights into optimal practices for preserving the Shami camel breed. Key considerations include improving reproduction, boosting productivity, developing fodder resources and training producers, supporting the sustainability of this valuable animal resource (Padalino *et al*, 2015; Sofiane *et al*, 2023).

Shami camel breeders' high-quality early care see faster growth and higher economic returns. Strong weight-growth rate links highlight the importance of breeding programmes for improved birth weights, herd productivity and market value. Utilising camels with high EBVs further maximises economic gains. Selective breeding and high-quality early care can significantly enhance breeder profitability. Therefore, this research was aimed to study the weights, growth rates and genetic aspects of the Shami camels.

SEND REPRINT REQUEST TO KHALED A AL-NAJJAR [email: khnajj2011@yahoo.com](mailto:khnajj2011@yahoo.com)



## Materials and Methods

### The herd

This research was conducted at the Deir Al-Hajar station, east of Damascus, Syria, for Shami camels adapted to semi-arid environments (Fig 1). The area receives annual rainfall between 100 to 125 mm, with coordinates at latitude: 33.5533 and longitude: 36.6339.

Camels were housed in concrete barns at night and during harsh weather for protection. They were vaccinated to prevent infectious diseases and spent 8-10 hours a day grazing on *Atriplex Salty* or *Solsola* rigid plants. Additionally, they received supplemental barley and bran (when available) and had free access to water. Breeding season occurs from November to March, with males allocated to females based on their reproductive efficiency. Both sexes can mate naturally when they reach 4-5 years old, depending on their weight. Calving happens from February to May and calves are weighed immediately after birth. Weaning occurs around 6 months of age, based on the calves' weight.

### Data Analyses

The data set was covered a period from 2002 to 2007, which included 217 records of Shami camel weights that were analysed by SAS (2012).

The General Linear Model was used to analysis data weights and growth rates as following model,

$$Y_{ijk} = \mu + R_i + X_j + e_{ijk} \quad (1)$$

Where,  $Y_{ijk}$ = Birth weight (BW), Weight at 6 months old (W6), Weight at 9 months old (W9), Weight at 12 months old (W12), Growth rate from

BW until W6 (BW6), Growth rates from BW until W9 (BW6), Growth rates from BW until W12 (BW12) of  $ijk^{th}$  observations.  $\mu$ = overall mean.  $R_i$ = Birth year (1= 2002, 2= 2003 ... and 6= 2007).  $X_j$ = Sex of calf (1= male and 2= female).  $e_{ijk}$ = Random error of  $ijk^{th}$  observations with mean = 0 and variance=  $\sigma_e^2$ .

Partial correlation coefficients were estimated from the SSCP errors matrix with their corresponding probability values denoted by (Prob.> |r|) according to equation (1).

$$Y_{ijk} = \mu + P_i + X_j + e_{ijk} \quad (2)$$

Where,  $Y_{ijk}$ = Birth weight of  $ijk^{th}$  observations.  $P_i$ = Parity (i = 1, 2... 6). The remaining symbols are shown in the previous models.

Heritability ( $h^2$ ) value was computed by paternal half-sib (Method=TYPE1) with this model:

$$Y_{ij} = \mu + S_i + e_{ij} \quad (3)$$

Where,  $Y_{ij}$ = Birth weight of  $ij^{th}$  observations.  $S_i$ = Sire (i = 1, 2... 9). The prior models display the remaining symbols.  $e_{ij}$ = Effect of environmental and genetic deviation related to individual in a group of sire. Therefore, the  $h^2$  value was computed in accordance to the Becker (1992).

$$h^2 = 4t, \quad t = \frac{V_s}{V_s + V_w}, \quad k = \frac{1}{s-1} \left\{ N - \frac{\sum N_i^2}{N} \right\}, \quad (3)$$

$$SE(h^2) = 4 \sqrt{\frac{2(1-t)^2(1+(k-1)t)^2}{k(k-1)(s-1)}}$$

Where,  $h^2$ = heritability value,  $V_s$ = Variance component of sire,  $V_w$ = Variance component of an individual, t and k are the constant,  $SE(h^2)$ = Standard



Fig 1. Female and male camles at Deir Al-Hajar station, Damascus countryside (Syria).

error of heritability,  $N$ = Total number of progeny,  $N_i$ = Number of progeny per sire.  $S$ = Number of sires.

$$EBV = \frac{N_i h^2}{4 + (N_i - 1)h^2} (P_{prog.} - P_{pop.}) \quad (4)$$

Estimated breeding values (EBV) were computed in accordance to the Hardjosubroto (1994).

Where, EBV= Estimated breeding value,  $N_i$ = Number of progeny per sire,  $h$ = Root of heritability value,  $P_{prog.}$ = Average trait of progeny,  $P_{pop.}$ = Average birth weight of the population. The previous models showed the remaining symbols.

Results

Table 1 displays notable disparities in birth weight (BW), weight at 12 months old (W12) and growth rates from birth to 12 months (BW12) across different birth years. Conversely, weight at 6 months old (W6), 9 months old (W9), growth rates from birth to 6 months (BW6) and growth rates from birth to 9 months (BW9) demonstrated no significant differences. When considering calf sex, (W6), (W12), (BW6) and (BW12) exhibited significant distinctions, while (BW), (W9) and (BW9) did not show significant variations.

Table 2 shows partial correlation coefficients for weights and growth rates to evaluate the relationships between response variables after controlling for the effects of other variables. It illustrates strong positive

correlations between weights at different stages of development in Shami camels, particularly between early growth (6 months) and later weight gain (9 and 12 months).

Table 3 presents mean birth weights (BW) and standard errors for Shami camels based on sex, parity and sire ID. Key findings included that male calves had a slightly higher mean BW than females, while multiparous mothers had higher BWs than primiparous mothers did. Sire ID had a notable influence, with sire 126 producing the heaviest calves and sire 243 producing the lightest. These findings highlight the importance of considering maternal parity and sire genetics to improve birth weight of calves in Shami camels.

Table 4 shows a high heritability rate (0.8) for birth weight in Shami camels suggesting a genetic influence. Sire selection in breeding programmes can improve birth weight, potentially leading to better weight outcomes for offspring.

Table 5 shows the top six sire ranked by their estimated breeding values for birth weight. The first-ranked sire (ID 126) has an estimated breeding value of +2.07 kg, indicating that his offspring tend to be heavier than average. The second-ranked sire (ID 111) has an estimated breeding value of +1.79 kg and so on. The lower-ranked sires have negative estimated breeding values, indicating that their offspring tend to be lighter than average.

**Table 1.** Least square means ± Standard Errors and variance of analysis of weights and growth rates of Shami camels under Semi-intensive rearing (Equation 1).

Source of variance	BW	W6	W9	W12	BW6	BW9	BW12
μ	33.6±1.07	172.2±7.60	198.1±6.85	238.0±6.35	23.1±1.28	18.2±0.75	17.0±0.53
Birth year	**	Ns	Ns	**	Ns	Ns	**
Sex	Ns	*	Ns	*	*	Ns	*
Residuals	16.388	830.352	674.448	579.630	23.750	8.162	4.052

μ= Grand mean. BW= Birth weight; W6= Weight at 6 months old; W9= Weight at 9 months old; W12= Weight at 12 months old. BW6= Growth rates from BW until W6; BW9= Growth rates from BW until W9; BW12= Growth rates from BW until W12. Coefficient of Variations for BW, W6, W9, W12, BW6, BW9 and BW12 = 11.74, 17.56, 13.21, 10.31, 22.56, 15.86, 12.1, respectively.

**Table 2.** Partial Correlation Coefficients for weights and growth rates of Shami camels under Semi-intensive breeding from the Error SSCP Matrix/Prob.>|r|, (Equation 1).

Variables	W6	W9	W12	BW6	BW9	BW12
BW	-0.04 <sup>ns</sup>	0.14 <sup>ns</sup>	0.06 <sup>ns</sup>	-0.17 <sup>ns</sup>	-0.01 <sup>ns</sup>	-0.10 <sup>ns</sup>
W6		0.50**	0.17 <sup>ns</sup>	0.99**	0.51**	0.17 <sup>ns</sup>
W9			0.45**	0.47**	0.90**	0.44**
W12				0.16 <sup>ns</sup>	0.46**	0.99**
BW6					0.51**	0.18 <sup>ns</sup>
BW9						0.46**

BW= Birth weight; W6= Weight at 6 months old; W9= Weight at 9 months old; W12= Weight at 12 months old. BW6= Growth rates from BW until W6; BW9= Growth rates from BW until W9; BW12= Growth rates from BW until W12.

Discussion

Table 1 reveals significant variations in birth weight (BW), weight at 12 months (W12) and growth rates from birth to 12 months (BW12) across different birth years of Shami camels reared under semi-intensive conditions. In contrast, weight at 6 months (W6), 9 months (W9), growth rates from birth to 6 months (BW6) and growth rates from birth to 9 months (BW9) did not show significant differences across birth years, suggesting stable early growth rates. In addition, significant differences in W6, W12, BW6 and BW12 were observed based on calf sex, suggesting that sex-specific management practices are necessary. These have implications for breeding strategies, nutritional and management practices and sex-specific management to improve the growth of Shami camels. Camels at birth weigh about 35 kilograms, but this can vary significantly between breeds and within breeds depending on regions (Kadim *et al*, 2008). No significant difference exists between the birth weights of male and female camels (Al-Momani and Al-Najjar, 2020). However, although the birth weight differences between sexes were minimal, male camels tended to be heavier than female camels as they grew older (Njanja and Oba, 2011). Significant weight differences emerge at 6, 12 and 18 months, with males typically being

heavier (Bakheit *et al*, 2017). Camel weights increased steadily from birth until age 4 under semi-intensive care conditions (Faris *et al*, 2022) and living weights continue to increase significantly as camels age (Seid *et al*, 2016). The year of birth also affects camel weight (Al-Momani and Al-Najjar, 2020). Camel growth rates are influenced by sex, nutrition, stress and health. These rates vary between strains and are affected by growth rates before and after weaning, which in turn affect body weights (Faraz, 2022). Birth weight differences between male and female camels are minimal; various factors such as sex, nutrition, health and environmental conditions significantly influence their growth rates and weights as they age. Understanding these factors can help optimise camel breeding and care practices to promote healthier and more robust growth.

Table 2 shows strong positive correlations between weights at different stages of Shami camel development, especially between early growth (6 months) and later weight gain (6 and 12 months). This indicated that early growth significantly influenced later weight gain. Conversely, the partial correlation coefficients between birth weight and later weights and growth rates were low and not statistically significant, suggesting birth weight had little impact on later development. These findings implied that

Table 3. Means ± standard error for birth weight base on sex and parity and sire of Shami camels (Equation 2).

Trait	Parity						Sex		Sire ID					
	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>th</sup>	4 <sup>th</sup>	5 <sup>th</sup>	6 <sup>th</sup>	M	F	111	118	126	230	238	243
BW (kg)	29.87±6.21	32.35±4.74	32.66±4.63	33.41±4.85	34.00±5.08	33.00±4.70	32.25±5.54	31.78±5.14	34.10±5.66	32.04±4.48	34.26±4.12	30.31±4.84	27.95±5.88	23.71±5.62

BW= Birth weight, M= Male, F= Female.

Table 4. Variance component and heritability ± standard error for birth weight of Shami camels (Equation 3).

Component	N <sub>s</sub>	N <sub>Prog.</sub>	V <sub>s</sub>	V <sub>w</sub>	k	t	h <sup>2</sup>
Values	6	217	7.38	29.58	33	0.20	0.80 ± 0.12

N<sub>s</sub>= Number of sire, N<sub>Prog.</sub>= Number of progeny, V<sub>s</sub>= Variance component of sire, V<sub>w</sub>= Variance component of individual within sire, k and t= Constant, h<sup>2</sup>= Heritability.

Table 5. Estimated breeding values for body weight of Shami camel males (Equation 4).

Sire rank	Sire ID	N <sub>prog.</sub>	BWp (kg)	EBV (kg)
1 <sup>st</sup>	126	65	34.10	+2.07
2 <sup>nd</sup>	111	29	32.04	+1.79
3 <sup>rd</sup>	118	56	34.26	-0.02
4 <sup>th</sup>	230	35	30.31	-1.57
5 <sup>th</sup>	238	20	27.95	-3.43
6 <sup>th</sup>	243	12	23.71	-5.31

N<sub>prog.</sub>= Number of progeny, BWp= Average birth weight of progeny, EBV= Breeding value.



optimising early growth was crucial for effective breeding, nutrition and management practices in Shami camels. According to these results, we should evaluate birth weight from an environmental and genetic perspective. Fatih *et al* (2021) showed a strong relationship between birth weight and mature weight. Furthermore, Njanja and Oba (2011) emphasised the important relationship between dam weights and calf birth weights. These underscore the importance of understanding the interaction between birth weight and subsequent growth and highlight potential implications for management practices in animal husbandry and husbandry programmes.

Shami camel birth weight (Table 3) shows trends, i.e. males were heavier than females and multiparous mothers exceeded primiparous ones. Sire genetics matter with sire 126 excelling. This highlights the importance of maternal experience and sire selection for improving birth weight. According to Almutairi *et al* (2010), parity significantly affects the birth weight of camels, indicating the importance of understanding maternal factors in camel breeding programs. Fatih *et al* (2021), who emphasised the importance of birth weight as a critical measurement in camels, suggesting its significance in monitoring camel weight, further supported this finding. Additionally, Bene *et al* (2020) confirmed that directing attention to the she-camel and optimising the environment could enhance the calf weight at birth, providing insights into practical strategies for improving calf weights in camel farming.

The heritability value of  $0.80 \pm 0.12$  indicates that approximately 80% of the variation in birth weight can be attributed to genetic factors. Sire selection would have an impact on improving birth weight compared to other traits (Table 4). Camel growth studies show varying genetic influences on weight. Some researchers found birth weight heritability high (Almutairi *et al*, 2010) or low (Bene *et al*, 2020). Later weight and gain showed moderate heritability (Al-Sobayil *et al*, 2006; Kaleri *et al*, 2017).

Table 5 shows sire's genetic merit for Shami camel birth weight using EBVs. Sires with positive EBVs (like 126 and 111) were predicted to have heavier offspring, while those with negative EBVs were predicted to have lighter offspring. This information helps breeders improve birth weight in future Shami camel generations. Estimating breeding values is crucial for evaluating performance and guiding breeding operations to achieve sustainable trait development using genetic models, as emphasised by Al-Mutairi *et al* (2010), who noted

a gradual improvement in weights and growth rates in camel breeding. Significant differences in breeding values for birth weight were found among male camels (Bene *et al*, 2020). Al-Sobayil *et al* (2006) assessed breeding values ranging from 25.3 to 115.1 for body weight and 0.270 to 0.638 kg for daily gains. The purposes are to enhance the selection process for superior camels, leading to improved weight and growth rates, achieving better economic returns and contributing to the sustainability and efficiency of camel breeding programs.

## Conclusion

This study highlights significant disparities in birth weight and growth rates across different birth years and sex-related differences were found. Strong positive correlations were found between weights at different stages, particularly between early growth and later weight gain. Maternal parity, sire genetics and breeding values play a crucial role in improving birth weight. Male calves and multiparous mothers had higher birth weights. A heritability value of 0.80 indicates that genetic factors contribute significantly to birth weight variation and breeding values of sire can improve birth weight.

## Recommendations

The study recommends the importance of taking into account maternal parity and the genetic breeding values of sires to improve birth weight and growth rates in Shami camels.

## Acknowledgements

The authors thank the universities and research institutions that provided essential administrative support for this collaborative research.

## Conflict of Interest

The authors declare no conflicts of interest to disclose.

## References

- Ali A, Baby B and Vijayan R. From desert to medicine. A Review of camel genomics and therapeutic products. *Frontiers in Genetics*. 2019; 10:17. DOI: 10.3389/fgen.2019.00017
- Al-Momani Ahmad and Al-Najjar Khaled. Some productive characteristics of Shami camels (*Camelus dromedarius*). *Journal of Camel Practice and Research*. 2020; 27(3):345-349. DOI:10.5958/2277-8934.2020.00050
- Almutairi SE, Boujenane I, Musaad A et al. Genetic and nongenetic effects for milk yield and growth traits in Saudi camels. *Tropical Animal Health and Production*. 2010; 42:1845-1853. DOI: 10.1007/s11250-010-9647-6
- Al-Owaimer AN, Suliman GM, Sami AS, Picard B and

- Hocquette JF. Chemical composition and structural characteristics of Arabian camel (*Camelus dromedarius*) m. longissimus thoracis. *Meat Science*. 2014; 96(3):1233-1241. DOI: 10.1016/j.meatsci.2013.10.025
- Al-Saiady MY, Mogawer HH, Al-Mutairi SE, Bengoumi M, Musaad A, Gar-Elnaby A and Faye B. Factors affecting feed intake, body weight, testicular size and testosterone, follicle-stimulating hormone (FSH) and luteinising hormone (LH) serum concentrations in peripubertal male camels. *African Journal of Agricultural Research*. 2015; 10(14):1709-1713. DOI:10.5897/AJAR2014.9319
- Al-Sobayil KA, Khalil MH, Al-Jobeile HS, Mohamed KM and Salal SA. Quantitative genetic analysis and evaluation for early growth performance in Saudi camels. In: *International Scientific Conference on Camels 2006*; pp 9-11.
- Bakheit Sallam Abdelfadeil, Faye Bernard, Ibrahim IE and Idris AO. The effect of management system on camel calves growth rate and daily gain. *Journal of Scientific and Engineering Research*. 2017; 4(2):41-47. <https://jsaer.com/archive/volume-4-issue-2-2017/>
- Balasundaram B, Thiruvankadan AK, Murali N, Muralidharan J, Cauveri D and Peters SO. Genetic parameters of growth traits and quantitative genetic metrics for selection and conservation of mecheri sheep of Tamil Nadu. *Animals (Basel)*. 2023; 13(3):454. DOI: 10.3390/ani13030454
- Becker WA. A Fortran Program for testing and analysing multiple comparisons. *Journal of Educational Statistics*. 1992; 17(4):344-353.
- Bene S, Szabó F, Polgár JP, Juhász J and Nagy P. Genetic parameters of birth weight trait in dromedary camels (*Camelus dromedarius*). *Tropical Animal Health and Production*. 2020. 52(5):2333-2340. Doi: 10.1007/s11250-020-02256-z
- Faraz A, Younas M, Waheed A, Tauqir NzA, Nabeel MS, Mirza RH, Ishaq HM and Khan NU. Growth management of marecha camel (*Camelus dromedarius*) in feedlot under desert conditions. *Advances in Animal and Veterinary Sciences*. 2020; 8(11):1114-1119. DOI: 10.17582/journal.aavs/2020/8.11.1114.1119
- Faraz A. Growth assessment in camel (*Camelus dromedarius*): A meta-analysis study. *Journal of Zoological Research*. 2022; 4(1):15-20. DOI: 10.30564/jzr.v4i1.2273
- Faris, Shadi; Al-Najjar, Khaled; Dib, Ali and Nouh Abdulla. Estimation of growth curve prediction equation parameters for weights of shami camel before mature/ Syria. *Arab Journal of Arid Environments*. 2022; 13(1). <https://digitalcommons.aaru.edu.jo/aae/vol13/iss1/9>
- Fatih A, Celik S, Eydurán E, Tirink C, Tariq MM, Sheikh IS, Faraz A and Waheed A. Use of MARS algorithm for predicting mature weight of different camel (*Camelus dromedarius*) breeds reared in Pakistan and morphological characterisation via cluster analysis. *Tropical Animal Health and Production*. 2021; 53:191. DOI: 10.1007/s11250-021-02633-2
- Fraser DJ. How well can captive breeding programmes conserve biodiversity? A review of salmonids. *Evolutionary Applications*. 2008; 1(4):535-86. DOI: 10.1111/j.1752-4571.2008.00036.x
- Hardjosubroto P. Breeding value of sires based on offspring weaning weight as a recommendation for selecting kebumen ongole grade cattle. *Journal of Animal Science*. 1994; 72(12):3456-3463.
- Joana L. Rocha, Raquel Godinho, José C Brito and Rasmus Nielsen, Life in deserts. The genetic basis of mammalian desert adaptation. *Trends in Ecology & Evolution*. 2021; 36(7):637-650. DOI: 10.1016/j.tree.2021.03.007.
- Kadim I, Mahgoub O and Purchas R. A review of the growth and of the carcass and meat quality characteristics of the one-humped camel (*Camelus dromedarius*). *Meat Science*. 2008; 80(3):555-569. DOI: 10.1016/j.meatsci.2008.02.010
- Kaleri RR, Kaleri HA, Baloch SK, Kaleri A, Mangi RA, Bhatti AL, Mari GM, Qureshi FN, Pahnwar MS and Marri A. Heritability estimates for some growth traits of dhatti camel breed in Tharparkar. *Journal of Basic & Applied Sciences*. 2017; 13:556-558.
- Njanja JC and Oba G. The turkana camel breed in North-Western Kenya: Part 2: Productivity characteristics. *Outlook on Agriculture*. 2011; 40(3):273-279. DOI: 10.5367/oa.2011.0053
- Osterloff Emily. How do camels survive in deserts? *Natural History Museum*. 2024; <https://www.nhm.ac.uk/discover/how-do-camels-survive-in-deserts.html>
- Padalino B, Monaco D and Lacalandra GM. Male camel behavior and breeding management strategies: how to handle a camel bull during the breeding season? *Emirates Journal of Food and Agriculture*. 2015; 27: 338-349.
- SAS. The SAS System for Windows. Release 9.1.3. SAS Institute, Cary, North Carolina, USA. 2012.
- Seid A, Kurtu MY and Urge M. Effect of age and body condition on slaughter characteristics of dromedary camels (*Camelus dromedarius*) in Eastern Ethiopia. *Journal of Camelid Science*. 2016; 9:35-53. <http://www.isocard.net/en/journal>
- Sofiane Boudalia, Yassine Gueroui, Rabah Zebba, Tarek Arbia, Ammar Elhassen Chiheb, M'hamed Benada, Zouheyr Hadri, Abdeldjalil Youcefi and Aissam Bousbia. Camel livestock in the Algerian Sahara under the context of climate change: Milk properties and livestock production practices. *Journal of Agriculture and Food Research*. 2023; 11: DOI: 10.1016/j.jafr.2023.100528.
- Suliman GM, Al-Owaimer AN, Hussein EOS, Abulfatah K and Othman MB. Meat quality characteristics of the Arabian camel (*Camelus dromedarius*) at different ages and post-mortem ageing periods. *Asian-Australasian Journal of Animal Sciences*. 2020; 33(8):1332-1338. DOI: 10.5713/ajas.19.0589

# INSTRUCTIONS TO CONTRIBUTORS

(Effective from the year 2024)

(Journal of Camel Practice and Research - triannual -April, August and December issues every year)

The Journal of Camel Practice and Research (JCPR) is a triannual journal (April, August and December issues) published in the English language by the Camel Publishing House, 67, Gandhi Nagar West, Near Lalgargh Palace, Bikaner, 334 001 (India). It is in offset print size of 20.5x27.5 cm in two columns with a print area of 17x22 cm. It will be known as **Journal of Camel Practice and Research** with **Volume number** on yearly basis and **Number** on issues per volume basis (in exceptional cases there can be more than three issues in a volume). The editorial policies of JCPR are established by the editor-in-chief and is detailed in this section. Views expressed in papers published in JCPR represent the opinions of the author(s) and do not necessarily reflect the official policy of the author's affiliated institution, or the editor-in-chief.

**Nature of coverage:** This journal is dedicated to disseminate scientific information about new and old world camelids in form of **Original research** articles in camel science, health, husbandry, pastoralism, sports, specific behaviour, history and socio-economics. **Reports** on unusual clinical case(s) or unreported management of clinical case(s) are also published. Review articles will be accepted on invitation only. **Book review** directly or indirectly related to camels will be reviewed by subject-matter specialists and included if sent to the journal for this purpose. The Journal of Camel Practice and Research will occasionally contain an **invited editorial** commenting on the current research and papers in the issue.

## Retraction guidelines (As per the formal COPE policy)

Editor in Chief of JCPR would consider retracting a publication, if:

- They have clear evidence that the findings are unreliable, either as a result of major error (eg, miscalculation or experimental error), or as a result of fabrication (eg, of data) or falsification (eg, image manipulation)
- It constitutes plagiarism
- The findings have previously been published elsewhere without proper attribution to previous sources or disclosure to the editor, permission to republish, or justification (i.e., cases of redundant publication)
- It contains material or data without authorisation for use
- Copyright has been infringed or there is some other serious legal issue (eg, libel, privacy)
- It reports unethical research
- It has been published solely on the basis of a compromised or manipulated peer review process
- The author(s) failed to disclose a major competing interest (aka, conflict of interest) that, in the view of the editor, would have unduly affected interpretations of the work or recommendations by editors and peer reviewers.

## Publication Ethics and Malpractice Statement

JCPR is a peer-reviewed journal and ensures the highest standards of publication ethics. All parties involved in the act of publishing process (editors, authors, reviewers and the publisher) have to agree upon standards of ethical behaviour. The accepted principles of Publication Ethics and Publication Malpractice Statement based on the Code of Conduct and Best Practice Guidelines for Journal Editors of the Committee on Publication Ethics – COPE (available at <http://publicationethics.org/>) are followed.

**Submission of manuscript:** Manuscripts should be submitted in word files to **Dr. Tarun Kumar Gahlot**, Editor, Journal of Camel Practice and Research at [tkcamelvet@yahoo.com](mailto:tkcamelvet@yahoo.com) or [editorjcpr@camelsandcamelids.com](mailto:editorjcpr@camelsandcamelids.com) by online

submission portals available at [www.indianjournals.com](http://www.indianjournals.com) or [www.camelsandcamelids.com](http://www.camelsandcamelids.com). The figures can be submitted preferably as a high pixel JPEG or other format. The manuscript should be accompanied by a covering note and author consent letter and ethical statement declaration from the author responsible for correspondence. It should also contain a statement that manuscript has been seen and approved by all co-authors. Editor and members of the editorial board are not responsible for the opinions expressed by authors and reserves the right to reject any material or introduce editorial changes. Material will be accepted for publication on the understanding that it has not been published in any other form and is not being considered elsewhere. Any material once accepted for publication may not be republished in any form without prior permission of the author. **Single blind peer review policy** is used for the manuscripts submitted and the quality standards are maintained by the JCPR editorial board and by authors who submit manuscripts for publication. A preprint PDF is provided to the author in correspondence for verification of contents and corrections, if any.

**Ethical declarations** in research form is an important step while submitting a manuscript to the Journal of Camel Practice and Research (JCPR). Author has to look into several questions and statements before submission. These are listed below:

## Pre-submission considerations for authors

- Manuscript should have been read and approved by all the authors. All the authors mentioned in the manuscript should have agreed for authorship and it's order. An author consent letter, duly signed should accompany the manuscript as per the given format (Author in correspondence can do it). (The authorship criteria should be based on the [ICMJE guidelines](#).)
- Full names, institutional affiliations, highest degree obtained by the authors, e-mail address (in some cases, [ORCID ID](#) and social media handles - Facebook, Twitter, or LinkedIn) need to be clearly mentioned on the title page.
- The corresponding author, who takes full ownership for all the communication related to the manuscript, should be designated and his/her detailed institutional affiliation (including the postal address, telephone number, fax number, and e-mail address) should be provided.
- In the research papers, JCPR expects a guarantor (who could be a senior research scientist or one of the author), he/she may be responsible for the integrity of the manuscript (including ethics, data handling, reporting of results, and study conduct), would communicate with the journal if any technical clarifications related to the manuscript are required, and would handle similar responsibilities.

All the authors need to agree on the name(s) included in the Acknowledgement section.

**Manuscript submission related declarations:** JCPR expects authors to declare the following:

- The manuscript in part or in full has **not been submitted or published anywhere**. In other words, the authors



should ensure that the manuscript is not a duplicate publication.

- The manuscript **will not be submitted elsewhere** until the editorial process is completed.
- If any part of the manuscript contains previously published content (figures/tables), authors should submit a **statement of permission to reproduce the material** signed by the author(s) and publishers concerned.
- When submitting material related to commercial products, it may, in some circumstances, be appropriate for the author to forward a copy of the contribution to the manufacturers before publication. This is to verify the correctness of the contents of the section in the manuscript that describes the new device/product.
- Authors should **declare any previous or pending publication** of the manuscript's content in any conference proceedings, letters to journals and brief communications, or as pre-prints on repositories.
- Authors should ensure that if the current study's Abstract has been published in any conferences, it should not be under a **copyright protection** from the publishers. Necessary permission should be sought for re-using the material.
- In case the manuscript is a secondary publication i.e. it is a subsequent republication that will be published in two or more journals (in the same or another language), the authors should explicitly declare this. Moreover, they should **obtain mutual consent of the journal editors** and follow the stipulated guidelines.
- If the manuscript is based on a dataset that has been the basis of another manuscript, authors should **maintain transparency** in such cases and should declare that by an appropriate reference.
- If there is a data set associated with the manuscript, provide information about where the data supporting the results or analyses presented in the paper can be accessed. Where applicable, this should include the hyperlink to publicly archived datasets, DOI, or other persistent identifier associated with the data set(s).

#### Statements of ethical approval for studies involving animals

If your study involves animals, and also if your manuscript includes case reports/case series, you need to provide the following:

- Authors must provide the **name of the ethical approval committee/Institutional Review Board** they have obtained consent from along with approval number/ID.
- Authors must state that **written informed consent was obtained** from the owners of the animals or head of the veterinary institutes where the research was carried out.

#### Declarations specific to article types

We have looked at the declarations related to manuscript submission and when your study involves human (for example-feeding camel milk to the diabetic human patients or autistic patients) or animal subjects. Let us now turn to specific article types and the declarations you need to prepare when submitting them.

1. **Clinical trials and research papers:** All clinical trials and planned research should be approved by the institutional ethics committee and should also be approved through an

appropriate research committee of the institute (It should be depicted in the acknowledgement section).

2. **Reviews:** Reviews do not need any ethical approvals or informed consent. However, JCPR expects that the review should not be older than 4 decades.
3. **Short Communications or Case Reports:** These are rare clinical reports or new diagnoses or a new technique (pilot trials), usually published with or without abstract and keywords. These article types do not exceed 2-3 pages of the journal.

#### Other important declarations related to funding, conflicts of interest, and more

Apart from the declarations we have discussed, there are others that authors need to consider. Let us take a look at them:

1. **Describing new taxa:** Authors must provide relevant documents and unique digital identifier for manuscripts that describe new taxa or species. They should also declare that the relevant guidelines have been followed for algae, fungi and plants, zoological taxa, bacteria, and viruses. Registration numbers for the new species (for e.g. from **Mycobank** for fungi or **ZooBank** for zoological species) should be stated in the manuscript. New virus names should be sent to the relevant study groups for consideration before publication in a journal.
2. **Authors' contribution:** The individual contributions of authors to the research work and writing of the manuscript should be specified in this section; for example, who conceived the study design, who did the data acquisition, who performed the experiments, who did the data analysis, who wrote the manuscript, etc. Authors should check journal-specific guidelines to declare the authors' contribution.
3. **Acknowledgements:** Anyone who does not meet the authorship criteria, such as people who provided technical help, institutional/department head who provided general support, or scientific writers who assisted with the preparation of the manuscript content, should be acknowledged. Even if the authors have no one to acknowledge, usually JCPR expects authors to include this section in the manuscript and write "Not applicable."
4. **Funding:** All sources of funding for the research work and their role (if at all) in the design of the study and collection, analysis, interpretation of data, and in writing the manuscript should be declared. Provide the name(s) of the funding agency/agencies along with the grant number(s). If the study did not receive any funding, report the same.
5. **Competing interests/conflict of interest:** All financial and non-financial competing interests must be declared by the authors. Non-financial competing interests include a declaration of political, personal, religious, ideological, academic, and intellectual competing interests. Authors from pharmaceutical companies, or other commercial organisations that sponsor clinical trials, should declare these as competing interests on submission.

Authors should declare any personal conflict of interest including any association with consultancies; employment details; participation in advocacy groups; stock or share ownership, and any financial details with regard to grants; fees; honoraria, reimbursements royalties, and any registered patents. They should also declare any institutional conflict of interest, i.e. if their employer has any financial interest in or is in conflict with the subject matter or materials discussed in

the manuscript. If there is no disclosure, add the following statement: “No potential conflict of interest was reported by the authors.”

It is mandatory and important that the authors declare all the above-mentioned statements to avoid un-submission of the manuscript. These declarations ensure ethical publication of the manuscript. JCPR expects from all the authors of manuscripts to read these practices involving transparency and integrity in the guidelines from the webpage of [www.camelsandcamelids.com](http://www.camelsandcamelids.com) in the Instructions to Contributors section.

### Initial Checks

The JCPR staff and in-house editorial team perform an initial quality check to identify potential issues such as:

- Competing interests
- Compliance with editorial policies and ethical standards
- Financial disclosures
- Data availability

Submissions may be returned to authors for changes or clarifications at this stage.

### Editorial Review

After completing internal checks, each new submission is assigned to an Academic Editor (Usually Editor in Chief) with relevant subject matter expert. The editor reviews the manuscript against our publication criteria and determines whether reviews from additional experts are required to evaluate the manuscript. The Editor in Chief decides about further handling of the manuscript by usually a member of the Editorial Board of JCPR, but occasionally a Guest Editor is invited to serve instead.

### Peer Review

The Editor in Chief selects reviewers, whether internal or external, based on expertise, publication history, and past reviews, and invites them to provide feedback on the manuscript. After agreeing to review, external peer reviewers typically have 10 days to submit their review. The journal office will follow up with late reviewers and keep you informed if there are any delays. **JCPR uses single-blind peer review policy.** Reviewers remain anonymous in our submission system.

### Editorial Decisions

The handling Academic Editor (Editor in Chief) or member of the editorial board makes the final decision on each manuscript. The time to render a first decision usually takes 2-4 weeks, but times vary depending on how long it takes for the editor to receive and assess reviews.

The Editor in Chief considers reviewer feedback and their own evaluation of the manuscript in order to reach a decision. The following decision types are available:

- Reject
- Major revision
- Minor revision
- Accept

Decisions are communicated to the corresponding author in a formal letter, along with reviewer feedback and any other requirements from the journal office.

### Revisions

If the editor feels that your manuscript has the potential to be published, but requires changes, you'll be invited to revise it. Usually authors are asked to resubmit the revised manuscript for both a major or a minor revision within 4 weeks.

In most cases, the revised manuscript is re-assigned to the original referee. The Editor in Chief may make a new decision based on their own assessment of the revised manuscript and your response to reviewers, or request additional input from external peer reviewers.

### Accepted Manuscripts

**JCPR uses two levels of accept decision. When the handling editor is satisfied with the scientific aspects of the manuscript a provisional acceptance decision is communicated, pending final checks for formatting and technical requirements. Once the final requirements are fulfilled, the journal office will send a formal acceptance and the manuscript will move on to production.**

**Preparation of the manuscript:** Manuscript should be typed in Book Antiqua font size 12 using British English, spellings and generic names of drugs. International Code of Zoological Nomenclature, Nomina Anatomica Veterinaria, International Code of Nomenclature of Bacteria, International Code of Botanical Nomenclature and International Standards should be strictly followed. All terms should be identified by their scientific names and for easy comprehension common terms/names can be used. Population data and geographical distribution of camelids should invariably be avoided in introduction, unless it is warranted. Kindly remain restricted to the relevant subject matter of given title of manuscript only. The review of literature should be up to the submission year of manuscript. Kindly check every reference for its accuracy from the relevant book, journal or through google.

### Each manuscript should have the following sections:

**Title page:** This page should contain title of the article, name of the department/institution where work has been done, present postal address of each author and name of author with email to whom reprint request should be addressed.

Running title should be short, succinct title with not more than 45 keystrokes (characters plus spaces) in length. A title should have the first word and proper nouns capitalised. Species of subject is encouraged. The title should be unique. All authors should have preferably full names (given name, middle initial, family name). Institutions of the authors with location should be denoted with either symbols (\*, †, ‡, §, #, |, and ¶) or Roman numerals behind the author's last name. Affiliation should also include department, city, state, country, and postal code.

Following is the example:

Example:

PROTEOMIC CHARACTERISATION OF SERUM DURING THE BREEDING CYCLE IN MALE BACTRIAN CAMELS

Le Hai<sup>1</sup>, Rendalai Si<sup>2</sup>, Fu-Cheng Guo<sup>1</sup>, Jing HeI, Li Yi<sup>1</sup>, Liang Ming<sup>1</sup>, Jun-Wen Zhou<sup>3</sup>, La Ba<sup>3</sup>, Rigetu Zhao<sup>3</sup> and Rimutu Ji<sup>1,2</sup>

<sup>1</sup>Key Laboratory of Dairy Biotechnology and Bioengineering, Ministry of Education, Inner Mongolia Agricultural University, Hohhot, Inner Mongolia, China

<sup>2</sup>Inner Mongolia Institute of Camel Research, Badanjiran, Inner Mongolia, China

<sup>3</sup>Alxa League Institute of Animal Husbandry, Alxa, Inner Mongolia, China

SEND REPRINT REQUEST TO RIMUTU JI email: yeluotuo1999@vip.163.com

**Abstract and Key words:** A single paragraph of no more than 2,500 keystrokes (characters plus spaces) that summarises the results in an understandable form using statistical evidence (P-values). Abbreviations are defined at first use in the ABSTRACT and again in the body of the manuscript. List up to 6 key words in alphabetical order and separated by a comma. Capitalise only proper nouns. Do not use abbreviations. Place the key words at the end of the Abstract.

**Text:** The proper text of the paper should commence after key words. The text should be divided into sections with headings, introduction, materials and methods, results, discussion, tables/illustrations and references.

**Introduction:** The logic of the introduction is to introduce the specificity and relevance of the topic to the readers. It should include the objective of the work in brief and most important related reference(s). The contents of the introduction should be lucid showing a clear justification for conducting the research with a stated hypothesis and objective(s) is required. The population data of camelids and their importance should be avoided as it is well known and is not relevant to the title in most of the manuscripts. Authors should not write the word, "First report" as later it is found that previous reports exist. The rationale for the experiments should place the work into the context of existing literature. There is NO word limit on the section but brevity is encouraged.

**Materials and Methods:** Should contain details regarding materials and brief account of procedures used. The manuscript must include a statement of institutional animal care and use or ethical committee, or country-specific equivalent, approval of all animal procedures. This statement should appear as the first item in MATERIALS AND METHODS and should specify which publicly available animal care and use standards were followed. A clear description of all biological, analytical and statistical procedures is required with each section denoted by a short descriptive title (i.e., Animals and sampling, Tests used, Treatment, Experimental design and analysis, etc). Materials used must include the product name and vendor at first mention. When a commercial product is used as part of an experiment, the manufacturer name and location must be given parenthetically and the generic name should be used subsequently. No <sup>TM</sup>, ®, or © symbols should be used. Sex, breed, age, species are included in the animal descriptions. Provide evidence of assay validation, or suitable published reference, as well as inter/intra-assay CV, as needed. Appropriate statistical methods should be used with experimental unit defined. Numbers of biological and experimental replicates should be stated. State the threshold for significance ( $P < 0.05$ ) and definition of tendency if used. In case reports, the case record sheet should also be included in materials and methods.

**Results and Discussion** should be presented in logical sequence with implications of findings about other relevant studies. The data or information easily attainable from the tables or graphics need not be repeated in the results. Only important observations need to be summarised. Undue repetition of the text from results to discussion has to be avoided. To preclude it, depending on article, results and discussion can be combined. In discussion only significant results should be discussed. One

should not always stick to the term 'statistically significant' data rather biological importance or significance of any variation should be given due importance in discussion. The section contains the interpretation of the results. It should be clear and concise, address the biological mechanisms and their significance, and integrate the results into existing literature. The Discussion may offer an interpretation that is consistent with the data. Do not include any reference to tables and figures or include P-values in the Discussion. Authors have the option to create a single RESULTS AND DISCUSSION section.

**Tables:** Each tables should be typed on separate sheet. Large tables should be avoided and should not exceed one page. Each table should be numbered in Indo-Arabic numerals according to their sequence in the text that refers to it. In the text it should be referred as proper noun e.g., Table 1. The title of the table should be brief and self-explanatory. Footnotes can be included to enhance understanding ability of the contents of the table.

**Illustrations and Legends:** All illustrations should be submitted about twice the size desired for reproduction that is 17 cm for double column or 8.3 cm for single column. Photographs should be of good quality with adequate contrast and high pixels. All illustrations should be referred as figures in the text and should also be numbered in Indo-Arabic numerals e.g., Fig 1. Legends of all these figures should be typed on a separate sheet. Each legend should be clear, concise and informative. A statement of magnifications or reductions should be given where it is applicable.

#### Disclosures (Conflict of Interest)

The authors must disclose any actual or potential conflicts of interest that may affect their ability to objectively present or review research or data. A succinct statement detailing any perceived conflict of interest is required. If none, please indicate as such.

**References:** Papers in the section must either be published or 'in press'. All references must include the DOI, if available. Authors are encouraged to use the most recent reference style for the Journal of Camel Practice and Research in the reference writing. References should be arranged in alphabetical order. Authors should not modify the title of the references. Mention full name of the journal.

#### Examples of correct forms of references are given below:

**Periodicals:** Shawaf T, El Nahas A, Melegi A, Al Bulushi S, Aiyan AA and Eljalli I. Investigation on biochemical parameters of cerebrospinal fluid in camels with neurological disorders. *Journal of Camel Practice and Research*. 2020; 27(2):165-171.

Wilson R Trevor. The one-humped camel in Eritrea and Ethiopia: a critical review of the literature and a bibliography. *Journal of Camel Practice and Research*. 2020; 27(3):229-262.

**For edited symposium/congress/proceedings:** Abdalla HS. Camel trypanosomiasis in the Sudan. *Proceedings of the First International Camel Conference, Dubai (UAE)*. February 2-6, 1992; pp 401-403.

**Books (Personal authors):** Faye B and Bengoumi M. *Camel Clinical Biochemistry and Haematology*: Springer International Publishing. 2018; pp 275-286.

**Chapter from multiauthored books:** Wernery U, Kinne J and Schuster RK. Unusual arboviruses and other minor viral infections. In: *Camelid Infectious Disorders*. OIE Book. 2014; pp 319-322.



**Thesis:** Rathod Avni. Therapeutic studies on sarcopticosis in camels (*Camelus dromedarius*). Unpublished Masters Thesis (MVSc), Rajasthan Agricultural University, Bikaner, Rajasthan, India. 2006.

**Commercial booklets:** Anonymous/Name. Conray-Contrast Media. IIIrd Edn., 1967; pp 12-15, May and Baker Ltd., Dagenham, Essex, England.

**Magazine articles:** Taylor D. The Constipated Camel. Reader's Digest. Indian Edn. RDI Print & Publishing (P) Ltd., Mehra House, 250-C, New Cross Road, Worli, Bombay, India. 1985; 126:60-64

**News paper articles:** Christina Adams. Camel milk: a miracle cure for children with autism?. Gulf News, Published: April 09. 2014.

**Personal communication:** Hall LW. Reader in Comparative Anaesthesia, Department of Clinical Veterinary Medicine, Madingley Road, University of Cambridge, Cambridge, CB3 0ES, England. 1995.

**Reprints or Publication charges:** There is no provision for free reprints. Author or person in correspondence has to pay INR 4500/- (for Indian Citizens only) or US \$ 450, in advance to receive a final PDF or for 10 reprints for the year 2024. Publication charges for colour and black and white pictures: Author(s) has to pay publication charges of colour plates in his/her manuscript as per the invoice provided to them. Publication charges would be double if a manuscript is more than 10 printed pages in length. The publication charges can be paid into the bank account of Camel Publishing House (using cheque, demand draft or money transfer method) or through a payment portal given in our website [www.camelsandcamelids.com](http://www.camelsandcamelids.com). Author(s) will be sent final pre-published PDF for verification and corrections, if any, along with an invoice and bank account details for making advance payment, before publication of manuscript in JCPR.

**Copyright:** The copyright of the article will remain with the owner, Dr Tarun Kumar Gahlot and will be governed by the Indian Copyright Act.

#### Author Consent Letter

{Note: Kindly take print out of the Author Consent letter form, fill, scan and send along with your manuscript. Without Author Consent letter your manuscript will not be considered for publication}

To  
Editor in Chief  
Journal of Camel Practice and Research  
Camel Publishing House  
Bikaner

**Sub: Submission of manuscript for publication in Journal of Camel Practice and Research reg.**

Dear Sir

I hereby submit the manuscript for publication in 'Journal of Camel Practice and Research'. I assure that this manuscript has neither been published in any other journal nor submitted for publication in any other journal. I also undertake along with the other authors that present study (on animals, human or laboratory animals, if any) was undertaken after the prior approval of relevant country/ institutional ethical committee. I and on behalf of other co-author(s), I declare "No conflict of interest". I assure that this article do not have any plagiarism. Kindly consider the manuscript for publication in your journal. I abide all rules and regulations of the journal. In future if any litigation arises in this article I will cooperate with the editor to resolve the issue. I shall accept the decision of the editor and that would be final.

Thanking You

SIGNATURE of Corresponding Author  
NAME  
ADDRESS

#### Kindly enter the details in the following

Manuscript

TITLE:

SUBJECT:

LABORATORY/RESEARCH CENTRE/INSTITUTE:

#### Ethical Statement Declaration

Hereby, I /insert author name/ consciously assure that for the manuscript /insert title/ the following is fulfilled:

- 1) This material is the authors' own original work, which has not been previously published elsewhere.
- 2) The paper is not currently being considered for publication elsewhere.
- 3) The paper reflects the authors' own research and analysis in a truthful and complete manner.
- 4) The paper properly credits the meaningful contributions of co-authors and co-researchers.
- 5) The results are appropriately placed in the context of prior and existing research.
- 6) All sources used are properly disclosed (correct citation). Literally copying of text must be indicated as such by using quotation marks and giving proper reference.
- 7) All authors have been personally and actively involved in substantial work leading to the paper, and will take public responsibility for its content.

The violation of the Ethical Statement rules may result in severe consequences.

To verify originality, your article may be checked by the originality detection software for plagiarism.

I agree with the above statements and declare that this submission follows the policies of Journal of Camel Practice and Research as outlined in the Guide for Authors and in the Ethical Statement.


Date:

Corresponding author's signature:

# THE CAMEL

## THE ANIMAL OF THE 21<sup>ST</sup> CENTURY

This book authored by Dr Alex Tinson is an acknowledgement to the support and inspiration that His Highness Sheikh Khalifa Bin Zayed Al Nahyan has provided to the centre and to research in general. The last 25 years has been an incredible adventure for us, the noble camel and the people of the U.A.E. Dr Tinson has been involved with many world first's since moving to Abu Dhabi 25 yrs ago. First there was the establishment of pioneering centres in exercise physiology and assisted reproduction. The establishment of the Hilli Embryo Transfer Centre led to five world firsts in reproduction. The world's first successful embryo transfer calf birth in 1990, followed by frozen embryo transfer births in 1994, twin split calves in 1999, pre-sexed embryo births in 2001 and world's first calf born from A.I. of frozen semen in 2013. The hard bound book is spread in 288 pages with 5 chapters. The first chapter involves early history of the centre, world's firsts, world press releases, history of domestication and distribution, evolution of camel racing in the U.A.E. and historical photos the early days. Second chapter comprises camel in health and disease and it involves cardiovascular, haemopoetic, digestive, musculoskeletal, reproductive, respiratory, urinary and nervous systems in addition to the description of special senses. This chapter describes infectious, parasitic and skin diseases in addition to the nutrition. The third chapter is based on Examination and Differential Diagnosis. The fourth chapter is based on special technologies bearing description of anaesthesia and pain management in camels, diagnostic ultrasound and X-Ray, assisted reproduction in camels, drug and DNA testing and surgery. The last chapter entailed future scope of current research.



Author  
Dr Alex Tinson

First Edition : 2017  
© 2017 Camel Publishing House



Publisher:  
**Camel Publishing House**  
67, Gandhi Nagar West, Near Lalgah Palace  
Bikaner-334001, India  
Email : [tkcamelvet@yahoo.com](mailto:tkcamelvet@yahoo.com)

Website:  
[www.camelsandcamelids.com](http://www.camelsandcamelids.com)  
[www.tkgahlotcamelvet.com](http://www.tkgahlotcamelvet.com)

ISBN : 81-903140-5-X  
Printed in India