

# SALIVARY SEX STEROID HORMONES IN FEMALE BACTRIAN CAMELS (*Camelus bactrianus*) DURING DIFFERENT REPRODUCTIVE STAGES

T. Fedorova<sup>1</sup>, K. Brandlová<sup>1</sup>, M. Bičíková<sup>2</sup>, I. Skálová<sup>1</sup> and D. Lukešová<sup>1</sup>

<sup>1</sup>Department of Animal Science and Food Processing, Faculty of Tropical Agriculture Sciences, Czech University of Life Sciences Prague, Prague, 165 21 Czech Republic

<sup>2</sup>Department of Steroids Hormones and Proteohormones, Institute of Endocrinology, Prague, 116 94 Czech Republic

## ABSTRACT

The study aimed to verify the usage of salivary sex steroid hormones monitored in captive bactrian camels (*Camelus bactrianus*) as a new non-invasive method in this species. Saliva of 5 adult female camels housed in the Prague zoological garden were sampled for more than 1 year in maximum interval of 10 days and concentrations of progesterone (P<sub>4</sub>) and oestradiol (E<sub>2</sub>) were measured. The concentrations of P<sub>4</sub> (n = 312) and E<sub>2</sub> (n = 310) were both significantly (p < 0.0001) affected by pregnancy status of animals. Mean (±SE) P<sub>4</sub> concentrations in non-pregnant stages were 2.234 (±0.220) nmol/l, while during the 2<sup>nd</sup> third of pregnancy it was 5.105 (±0.858) nmol/l. E<sub>2</sub> concentrations differed significantly between non-pregnant stages with mean value 0.037 (±0.005) nmol/l and during the 3<sup>rd</sup> third of pregnancy when reached 0.098 (±0.012) nmol/l. The seasonal differences in non-pregnant female camels were also evaluated. While no significant seasonal deviations were found in E<sub>2</sub> concentrations, P<sub>4</sub> values were significantly higher in summer than in spring. The study concluded that salivary P<sub>4</sub> and E<sub>2</sub> measurements are suitable for monitoring different reproductive stages in half-tamed female camels. The autumn and winter seasons seemed to be the best for pregnancy diagnosis in camels bred in Europe.

**Key words:** Camelid, oestradiol, pregnancy, progesterone, saliva

All studies focused on hormonal changes in camels were carried out so far only from blood, e.g. study of Elias *et al* (1984), Zhao *et al* (1994), Skidmore *et al* (1996a, 1996b), or from milk (Abdel Rahim and Elnazier, 1987; Abdel Rahim, 1989). The hormonal changes in serum or plasma have been already well documented in female camels. Oestradiol, also reported as oestradiol-17β, (E<sub>2</sub>) and progesterone (P<sub>4</sub>) concentrations were often used for monitoring the reproductive cycle and pregnancy in dromedary camels (*Camelus dromedarius*) (Skidmore *et al*, 1996a, 1996b; Ayoub *et al*, 2003; Muhammad *et al*, 2011) but also in bactrian camels (*Camelus bactrianus*) (Zhao *et al*, 1994; Zhao *et al*, 1998) with similar results. Nevertheless, camels are seasonal breeders (Al-Hazmi, 2000, Nowshari and Ali, 2005; Ali *et al*, 2008; El-Harairy and Attia, 2010; El-Harairy *et al*, 2010) and hormonal changes in non-pregnant animals can also be influenced by a season of the year (Agarwal and Khanna, 1993; Al-Qarawi *et al*, 2000; El-Harairy and Attia, 2010). The breeding season depends on area of distribution, climate, temperature, humidity, day light length and rainfalls (Bono *et al*, 1989; El-Harairy and Attia, 2010). The average gestation period is between

370 and 385 days in dromedary camels (Skidmore *et al*, 2000; Musa *et al*, 2006; Al-Sobayil, 2008) and 402.22 ± 11.5 days in bactrian camels (Zhao *et al*, 1994).

Camels are able to produce large amount of saliva (Haberoová *et al*, 2012; Wemmer and Murtaugh, 1980; von Engelhardt *et al*, 2006) and monitoring of reproduction through saliva has been already confirmed in other ungulate species, e.g. in buffaloes (*Bubalus bubalis*) (Qureshi *et al*, 1999), black rhinoceros (*Diceros bicornis*) (Czekala and Callison, 1996) or Indian rhinoceros (*Rhinoceros unicornis*) (Gomez *et al*, 2004).

The aim of the study was to verify the potential of steroid hormones monitoring in camels from saliva as a new non-invasive method in bactrian camels and evaluate the seasonal changes in non-pregnant female camels kept in European zoological garden.

## Materials and Methods

### Animals and husbandry

Five adult female bactrian camels (*Camelus bactrianus*) housed in the Prague zoological garden, Czech Republic, aged between 14-24 years were included in the research. Females were kept together

SEND REPRINT REQUEST TO T. FEDOROVA [email: fedorova@ftz.czu.cz](mailto:fedorova@ftz.czu.cz)

with male for the whole time and these were able to mate freely, so the time of conception was not known. Each third of pregnancy was defined retrospectively from parturition and it is represented according to respective days based on overall gestation length. The term of parturition was used for data analysis. At least  $\frac{1}{3}$  out of every detected pregnancy ( $n=5$ ) was monitored during the sampling period. Non-pregnant stages were noticed in 4 females.

Animals were fed by meadow hay, fresh vegetable, pellets for herbivores, and oats for the majority of year. Fresh fodder was provided during some parts of spring. Animals were trained for human contact, enabling touching and feeding from hand but another type of manipulation was not possible.

### **Sample collection and hormonal assay**

Samples were collected from 3<sup>rd</sup> February 2011 to 27<sup>th</sup> February 2012. The interval between samplings was 2 or 3 days during the period between December and March, 7 days during April, May and November, and 10 days between June and October.

The whole herd was lured close to the fence by showing a bucket with dried bread. Usually, whole herd ran up to the sampling place immediately and animals were showing their willingness to cooperate. The sampling was carried out through the bar fence. In selected animal, the production of saliva was stimulated by feeding a piece of dried wheat or wheat-rye bread without any pieces of salt or cereals. When the camel was trying to get to the bread, saliva was wiped off from the tongue by the handle of a disposable plastic spoon. Saliva was collected in the crease of the spoon handle and was removed to the polypropylene centrifuge tubes of volume 15 ml. The process was repeated several times until the amount of foamy saliva reached 5 ml. The animal was rewarded by piece of bread.

The test tubes with saliva were stored in temperature between 4 - 8°C and transported to a freezer (-20°C) within 12 hours after the sampling. The concentrations of  $P_4$  and  $E_2$  were measured in the Institute of Endocrinology (Prague, Czech Republic). The total unconjugated  $17\beta$ -oestradiol was measured by the RIA Spectria Estradiol kit<sup>®</sup> (Orion Diagnostica Oy, Finland) and  $P_4$  by the Progesterone RIA kit (Immunotech, A Beckman Coulter Company, France). The salivary samples were processed according to the guidelines of the kits' producers and by the same methods as samples of serum.

All animals included in the research cooperated of their own free will and they were not forced

to sampling. The research was approved by the management of the Prague zoological garden and also by the veterinarian of this institution.

### **Data analysis**

Stages of pregnancy were counted from the term of parturition. One pregnancy ended by abortion and the term of possible parturition was counted from Crown Vertebral Rump Length (CVRL) of the aborted calf (Elwishy *et al*, 1981; Bello *et al*, 2012; Hena and Sonfada, 2012). The thirds of camel pregnancy were based on the study of Bello *et al* (2012): 130 days or less to parturition were classified as 3<sup>rd</sup>  $\frac{1}{3}$ , 131 to 260 days to parturition as 2<sup>nd</sup>  $\frac{1}{3}$  and 261 days or more as 1<sup>st</sup>  $\frac{1}{3}$ .

Periods of the year were classified according to meteorological reckoning: spring (March, April, May), summer (June, July, August), autumn (September, October, November), and winter (December, January, February) (Barnett and Dobson, 2010).

The statistics were performed using software Statistica CZ 12 (Stat Soft, Inc.). The data distribution was not normal ( $p < 0.01$ ; Lilliefors test and Shapiro-Wilk test) and the non-parametric statistics (Kruskal-Wallis test, Multiple comparison of p-values, Mann-Whitney U test) were used to analyse the results. The significance level was accepted at  $p < 0.05$ .

Low, undetectable concentrations of  $P_4$  ( $n=16$ ) were obtained mainly (75.0%) in non-pregnant animals.  $E_2$  concentration under detectable level was noticed only in one sample. These samples were analysed as zero concentrations. Two samples overreached a maximum measurable  $E_2$  concentration and these samples were excluded from the analyses.

### **Results**

312 samples of camel saliva were collected and successfully analysed. The concentrations of  $P_4$  ( $n = 312$ ) and  $E_2$  ( $n = 310$ ) were both significantly ( $p < 0.0001$ ) affected by pregnancy status of animals (Kruskal-Wallis test:  $H(4, N = 313) = 27.1809$  and  $H(4, N = 311) = 39.3600$ , respectively). Mean  $P_4$  and  $E_2$  levels during different stages of reproduction are provided in Table 1.

### **Hormonal changes during pregnancy**

Mean  $P_4$  concentrations are shown in Fig 1.  $P_4$  concentrations fluctuated during the pregnancy and the peak, averaging  $6.06 \pm 2.90$  nmol/l, was reached between the 150<sup>th</sup> and 126<sup>th</sup> day before parturition (BP). The mean  $P_4$  concentrations fell to  $3.61 \pm 0.75$  nmol/l by the last 50 days of pregnancy.

As shown in Fig 2, the  $E_2$  concentrations increased rapidly 175 days BP and reached the 1<sup>st</sup>

peak, averaging  $0.115 \pm 0.060$  nmol/l, between the 150<sup>th</sup> and 126<sup>th</sup> day BP. The E<sub>2</sub> increased significantly (Mann-Whitney U Test,  $U = 447$ ,  $p < 0.01$ ) again 50 days BP and reached the values of  $0.120 \pm 0.022$  nmol/l in the last 25 days of pregnancy.


### Hormonal changes in non-pregnant animals during the year

In non-pregnant animals, the P<sub>4</sub> values were significantly affected by the season (Kruskal-Wallis test:  $H(3, N = 114) = 10.2520$ ,  $p = 0.0165$ ). P<sub>4</sub> concentrations were significantly higher in summer than in spring (Multiple comparison of p-values,  $p = 0.0345$ ) and as shown in Fig 3, similar trend visible for the winter season (Multiple comparison of p-values,  $p = 0.0522$ ). The concentrations during autumn and winter stayed always under 6.5 nmol/l.

P<sub>4</sub> concentrations differed significantly also between months (Kruskal-Wallis test:  $H(11, N = 114) = 27.0522$ ,  $p = 0.0045$ ) and the highest mean concentrations were reached in July and August (4.248 and 5.055 nmol/l). The lowest mean P<sub>4</sub> concentrations were noticed during February (1.159 nmol/l).

**Table 1.** Salivary progesterone and oestradiol concentrations in female camels during different phases of reproduction.

	Progesterone (nmol/l)		Oestradiol (nmol/l)	
	Mean ± SE	Range	Mean ± SE	Range
Non-pregnant (n = 114)	2.2339±0.2197 <sup>a</sup>	0.0000 - 14.4100	0.0374±0.0049 <sup>a</sup>	0.0010 - 0.3620
Post-partum period (n = 14)	1.5193±0.3057 <sup>a,b</sup>	0.0000 - 3.3600	0.0500±0.0099 <sup>a,b</sup>	0.0120 - 0.1630
1 <sup>st</sup> third of pregnancy (n = 66)	2.5527±0.3346 <sup>a,b</sup>	0.0000 - 11.6400	0.0564±0.0086 <sup>b</sup>	0.0000 - 0.3680
2 <sup>nd</sup> third of pregnancy (n = 29/28)	5.1045±0.8584 <sup>c</sup>	0.3600 - 23.3100	0.0693±0.0181 <sup>a,b</sup>	0.0060 - 1.6500
3 <sup>rd</sup> third of pregnancy (n = 90/89)	4.3357±0.5348 <sup>b,c</sup>	0.0000 - 26.2700	0.0978±0.0118 <sup>b</sup>	0.0078 - 0.6149



**Fig 1.** Mean ± SE progesterone concentrations in camel saliva during pregnancy.

However, no significant seasonal or monthly differences ( $p > 0.05$ ) were found in E<sub>2</sub> concentrations.

### Discussion

The study is the first report of salivary sex steroid hormone monitoring in camels and also the first study focused on hormone monitoring in camels bred in Europe. Despite the fact that the number of studies focused on hormonal analysis from saliva in animals is limited and concentrations of hormones in saliva are fluctuating (Hofman, 2001; Kobelt *et al*, 2003), the presented results showed that salivary P<sub>4</sub> and E<sub>2</sub> are suitable for monitoring of different reproductive stages in half-tamed female camels similar as in other ungulate species, e.g. study of Czekala and Callison (1996), Moriyoshi *et al* (1996), Qureshi *et al* (1999) and Gomez *et al* (2004).

Mean P<sub>4</sub> and E<sub>2</sub> concentration changes during the pregnancies in presented study had similar pattern as in study of Skidmore *et al* (1996a) or Zhao *et al* (1998) dealing with blood samples of camels. A positive correlation between salivary and serum




Fig 2. Mean ± SE oestradiol concentrations in camel saliva during pregnancy.




Fig 3. Mean ± SE progesterone concentrations in saliva of non-pregnant female camels during different periods of the year.

oestradiol was already reported by Qureshi *et al* (1999) in buffaloes and the fact that serum hormonal concentrations correlated with concentrations in saliva was well proved in human (Hofman, 2001).

Salivary sex steroid hormones concentrations seemed not to be suitable for early pregnancy diagnosis in camels because no significant differences

between pregnant and non-pregnant animals were found. However, with repeated sampling, it could be possible to determine pregnancy during the 2<sup>nd</sup> or 3<sup>rd</sup> third of pregnancy. Contrary to research of Volkery *et al* (2012) carried out on alpacas, this study confirmed the significant differences between P<sub>4</sub> concentrations at these stages of pregnancy and non-

pregnant animals. Some results evaluated in the first 2 months of pregnancy could also be misrepresented because it was not possible to determine exact time of conception. Higher E<sub>2</sub> concentrations noticed in the beginning of pregnancy (Fig 2), between 375<sup>th</sup> and 400<sup>th</sup> day before parturition, were also probably caused by the impossibility to know the exact day of conception and some animals could be in oestrus in that time.

Contrary to Hussein *et al* (2008), seasonal difference in P<sub>4</sub> concentrations were proved non-pregnant animals. Higher P<sub>4</sub> concentrations during the spring and summer could be caused by unrecognised early embryonic losses that are common in camels (Nagy and Juhasz, 2008); some animals could conceived during spring but they did not give a birth due to embryonic loss and they could be incorrectly assessed as non-pregnant.

The majority (89.4%) of parturitions of camels kept in the Prague zoological garden occurred between February and May (Haberová and Fedorov, 2012), so autumn or winter periods belongs usually to 2<sup>nd</sup> or 3<sup>rd</sup> third of pregnancy in camels. This fact, together with the finding that P<sub>4</sub> stayed at lower concentrations during autumn and winter in non-pregnant animals instigate to the conclusion that these 2 periods of the year seem to be the best for pregnancy diagnosis in camels in Europe or temperate regions of northern Hemisphere.

Present study did not prove any significant variation in mean values of E<sub>2</sub> in non-pregnant animals between different months or seasons similar like Hussein *et al* (2008) but the maximum fluctuations were reached during February, March, June and December. This feature could be caused to the presence of heat in camel females (Skidmore *et al*, 1996a) which was not possible to detect in presented study and selected breeding facility.

In conclusion, (1) hormonal analyses of saliva are suitable for reproductive status monitoring in female camels and more detailed research at this topic can be recommended. (2) Both salivary steroid hormones, oestradiol-17 $\beta$  and progesterone, can be recommended for pregnancy diagnosis during the 2<sup>nd</sup> and 3<sup>rd</sup> third of camel pregnancy. (3) Autumn or winter seems to be the best periods for pregnancy diagnosis in camels in Europe.

### Acknowledgements

The author would like to acknowledge the staff of the Prague Zoological Garden for their assistance, especially Petr Fedorov, Barbora Crkvová, Jan

Marek, Jaroslav Šimek, and Roman Vodička, and the team from the Department of Steroids Hormones and Proteohormones, Institute of Endocrinology, for kindly cooperation. We also thank J. Steven Brown for English proof reading. The research was funded by the Internal Grant Agency of the Faculty of Tropical AgriSciences CULS Prague in the year 2011 (project no. 51120/1312/3112) and by CULS Prague University Internal Grant Agency (project no. 20145009).

### References

- Abdel Rahim SEA (1989). The use of milk progesterone analysis to monitor reproductive activity in the camel (*Camelus dromedarius*). *British Veterinary Journal* 145(1):23-27.
- Abdel Rahim SEA and Elnazier AE (1987). Estimation of progesterone level in camels (*Camelus dromedarius*) milk and its application in pregnancy diagnosis. *British Veterinary Journal* 143(6):555-559.
- Agarwal SP and Khanna ND (1993). Endocrine profiles of the Indian camel under different phases of reproduction. *Etudes et Syntheses de l'EMVT* 41:77-101.
- Al-Hazmi MA (2000). Mating behavioural aspect of one humped camel (*Camelus dromedarius*) in Jeddah Province, Saudi Arabia. *Saudi Journal of Biological Sciences* 7(2):113-122.
- Ali S, Ahmad N, Akhtar N, Zia ur R and Noakes DE (2008). Metabolite contents of blood serum and fluid from small and large sized follicles in dromedary camels during the peak and the low breeding seasons. *Animal Reproduction Science* 108(3-4):446-456.
- Al-Qarawi AA, Abdel-Rahman HA, El-Belely MS and El-Mougy SA (2000). Age-related changes in plasma testosterone concentrations and genital organs content of bulk and trace elements in the male dromedary camel. *Animal Reproduction Science* 62(4):297-307.
- Al-Sobayil KBA (2008). The use of estrus synchronisation and timed artificial insemination in dromedary she-camels in Saudi Arabia. *Journal of Agricultural and Veterinary Sciences* 1(1):3-9.
- Ayoub MA, El-Khouly AA and Mohamed TM (2003). Some haematological and biochemical parameters and steroid hormone levels in the one-humped camel during different physiological conditions. *Emirates Journal of Food and Agriculture* 15(1):44-55.
- Barnett AG and Dobson AJ (2010). *Analysing Seasonal Health Data*. (1<sup>st</sup> ed). Springer-Verlag, Heidelberg, Germany. pp 58-59.
- Bello A, Onyeausi BI, Sonfada ML, Adeyanju JB, Umar AA, Umaru MA, Shehu SA and Hena SA (2012). Histomorphological studies of the prenatal development of oesophagus of one humped camel (*Camelus dromedarius*). *Scientific Journal of Agricultural* 1(4):100-104.
- Bono G, Dahir AM, Comin A and Jumale MA (1989). Plasma LH, corticoid and sex steroid variations in camels (*Camelus dromedarius*) in relation to seasonal climatic changes. *Animal Reproduction Science* 21(2):101-113.

- Czekala NM and Callison L (1996). Pregnancy diagnosis in the black rhinoceros (*Diceros bicornis*) by salivary hormone analysis. *Zoo Biology* 15(1):37-44.
- El-Harairy MA and Attia KA (2010). Effect of age, pubertal stage and season on testosterone concentration in male dromedary camel. *Saudi Journal of Biological Sciences* 17(3):227-230.
- El-Harairy MA, Zeidan AEB, Afify AA, Amer HA and Amer AM (2010). Ovarian activity, biochemical changes and histological status of the dromedary she-camel as affected by different seasons of the year. *Nature and Science* 8(5):54.
- Elias E, Bedrak E and Yagil R (1984). Estradiol concentration in the serum of the one-humped camel (*Camelus dromedarius*) during the various reproductive stages. *General and Comparative Endocrinology* 56(2):258-264.
- Elwishy AB, Hemeida NA, Omar MA, Mobarak AM and El Sayed MA (1981). Functional changes in the pregnant camel with special reference to foetal growth. *British Veterinary Journal* 137(5):527-537.
- Gomez A, Jewell E, Walker SL and Brown JL (2004). Use of salivary steroid analyses to assess ovarian cycles in an Indian rhinoceros at the National Zoological Park. *Zoo Biology* 23(6):501-512.
- Haberová T and Fedorov P (2012). Umělý odchov velblouda dvouhřbého (*Camelus bactrianus*) - Hand-rearing of bactrian camel (*Camelus bactrianus*). *Gazella* 39:67-75.
- Haberová T, Skálová I, Brandlová K, Kajzrová S and Lukešová D (2012). The usage of disposable catering supplies for sampling of saliva and urine in even-toed ungulates. *Proceedings 6<sup>th</sup> Scientific Conference of Institute of Tropics and Subtropics: Sustainable Use of Natural Resources in Tropics and Subtropics, Prague, Czech Republic*. pp 51.
- Hena SA and Sonfada ML (2012). Morphological studies on the prenatal development of calvaria in dromedarian camel. *Journal of Animal Science Advances* 2(6):551-557.
- Hofman LF (2001). Human saliva as a diagnostic specimen. *Journal of Nutrition* 131(5):1621-1625.
- Hussein MM, El-Agawany AA and Amin K (2008). Ovarian activity of she-camel (*Camelus dromedarius*) in relation to season, hormonal pattern, age and body condition scores. *Beni-Suef Veterinary Medical Journal* 18(2):1-9.
- Kobelt AJ, Hemsworth PH, Barnett JL and Butler KL (2003). Sources of sampling variation in saliva cortisol in dogs. *Research in Veterinary Science* 75(2):157-161.
- Moriyoshi M, Tamaki M, Nakao T and Kawata K (1996). Early pregnancy diagnosis in the sow by saliva progesterone measurement using a bovine milk progesterone qualitative test EIA kit. *Journal of Veterinary Medical Science* 58(8):737-741.
- Muhammad B, Aliyu D, Njidda A and Madigawa I (2011). Some haematological, biochemical and hormonal profile of pregnant and non-pregnant she-camels (*Camelus dromedarius*) raised in a Sudan Savanna zone of Nigeria. *Journal of Camel Practice and Research* 18(1):73-77.
- Musa HH, Shuiep ES, El Zubier Ibtisam EM and Chen GH (2006). Some reproductive and productive traits of camel (*Camelus dromedarius*) in Western Sudan. *Journal of Animal and Veterinary Advances* 5(7):590-592.
- Nagy P and Juhasz J (2008). Reproductive efficiency of a non-traditional milk producing animal: The dromedary camel. *Proceedings WBC/ICAR, 2008 Satellite Meeting on Camelid Reproduction, Budapest, Hungary*. pp 66-69.
- Nowshari MA and Ali SA (2005). Effect of season and gonadotropins on the superovulatory response in camel (*Camelus dromedarius*). *Theriogenology* 64(7):1526-1535.
- Qureshi ZI, Lodhi LA, Bajwa MA, Muhammad G and Ahmad I (1999). Estradiol-17b concentration in different biological fluids during the oestrous cycle in buffalo. *Pakistan Journal of Agricultural Sciences* 36(3-4):106-108.
- Skidmore JA, Billah M and Allen WR (1996a). Patterns of hormone secretion throughout pregnancy in the one-humped camel (*Camelus dromedarius*). *Reproduction Fertility and Development* 8(5):863-869.
- Skidmore JA, Billah M and Allen WR (1996b). The ovarian follicular wave pattern and induction of ovulation in the mated and non-mated one-humped camel (*Camelus dromedarius*). *Journal of Reproduction and Fertility* 106(2):185-192.
- Skidmore JA, Billah M, Allen WR and Short RV (2000). Using modern reproductive technologies such as embryo transfer and artificial insemination to improve the reproductive potential of dromedary camels. *Revue d'élevage et de médecine vétérinaire des pays tropicaux* 53(2):97-100.
- Volkery J, Gottschalk J, Sobiraj A, Wittek T and Einspanier A (2012). Progesterone, pregnanediol-3-glucuronide, relaxin and oestrone sulphate concentrations in saliva, milk and urine of female alpacas (*Vicugna pacos*) and their application in pregnancy diagnosis. *Veterinary Record* 171(8):195-195.
- von Engelhardt W, Haarmeyer P and Lechner-Doll M (2006). Feed intake, forestomach fluid volume, dilution rate and mean retention of fluid in the forestomach during water deprivation and rehydration in camels (*Camelus* sp.). *Comparative biochemistry and physiology. Part A, Molecular and Integrative Physiology* 143(4):504-507.
- Wemmer C and Murtaugh J (1980). Olfactory aspects of rutting behaviour in the bactrian camel (*Camelus bactrianus ferus*). In: *Chemical Signals*. Müller-Schwarze D and Silverstein R) eds. 1<sup>st</sup> Springer US, California, USA. pp 107-124.
- Zhao XX, Huang YM and Chen BX (1994). Artificial insemination and pregnancy diagnosis in the bactrian camel (*Camelus bactrianus*). *Journal of Arid Environments* 26(1):61-65.
- Zhao XX, Zhang Y and Chen BX (1998). Serum progesterone and 17  $\beta$ -estradiol concentrations during pregnancy of bactrian camel (*Camelus bactrianus*). *Theriogenology* 50(4):595-604.